



Contribution ID: 136

Type: not specified

Improved automated lattice perturbation theory in background field gauge

Tuesday, 15 June 2010 18:00 (5 minutes)

We present an algorithm to automatically derive Feynman rules for lattice perturbation theory in background field gauge. Vertices with an arbitrary number of both background and quantum legs can be derived automatically from both gluonic and fermionic actions. The algorithm is a generalisation of our earlier algorithm based on prior work by Lüscher and Weisz. We also present techniques allowing for the parallelisation of the evaluation of the often rather complex lattice Feynman rules that should allow for efficient implementation on GPUs, but also give a significant speed-up when calculating the derivatives of Feynman diagrams with respect to external momenta.

Please, insert your presentation type (talk, poster)

poster

Primary author: VON HIPPEL, Georg (Institut f. Kernphysik, Uni Mainz)**Co-authors:** Dr HART, Alistair (University of Edinburgh); Mr MONAHAN, Chris (DAMTP, University of Cambridge); Dr MUELLER, Eike (University of Edinburgh); Prof. HORGAN, Ron (DAMTP, University of Cambridge); Mr HAMMANT, Thomas (DAMTP, University of Cambridge)**Presenter:** VON HIPPEL, Georg (Institut f. Kernphysik, Uni Mainz)**Session Classification:** Poster session**Track Classification:** Algorithms and machines