



Contribution ID: 135

Type: **not specified**

non-locality of the nucleon-nucleon potential from Lattice QCD

Thursday, 17 June 2010 17:40 (20 minutes)

Recently, a new approach to calculate the nuclear potential from lattice QCD has been proposed. In this approach, the nuclear potential is constructed from Bethe-Salpeter (BS) wave functions through the Schroedinger equation. The procedure leads to non-local but energy independent potential $U(r, r')$, which can be expressed in terms of local functions by derivative expansion. In several recent applications of this method, local potentials, which correspond to the leading order (LO) terms of the derivative expansion, are calculated from the BS wave function at $E \simeq 0$ MeV, where E is the center of mass energy. It is therefore important to estimate the validity of the LO potential and size of contribution from higher order terms, which can be done by examining the energy and angular momentum dependence of the LO potential.

In this report, we will estimate the energy dependence of LO potential by comparing two LO potentials obtained at two energies in quenched lattice QCD with $m_\pi = 530$ MeV. We will also examine the possible L dependence of $S=0$ (spin singlet) potential, by comparing 1S_0 and 1D_2 central potentials. We find that, in each comparison, difference of the LO potentials is negligible within the statistical errors. This implies that the contribution from NLO terms are less significant, so that the LO potential obtained at $E \simeq 0$ ($L \leq 2$) is valid up to (at least) 45 MeV ($L=2$) to describe the NN interaction in quenched QCD with $m_\pi = 530$ MeV.

Please, insert your presentation type (talk, poster)

talk

Primary author: MURANO, Keiko (High Energy Accelerator Research Organization, KEK)

Co-authors: Prof. NEMURA, Hidekatsu (Tohoku University); Dr ISHII, Noriyoshi (Department of Physics, University of Tokyo); Prof. AOKI, Sinya (Graduate School of Pure and Applied Sciences, University of Tsukuba); Prof. HATSUDA, Tetsuo (Department of Physics, University of Tokyo)

Presenter: MURANO, Keiko (High Energy Accelerator Research Organization, KEK)

Session Classification: Parallel 38: Hadronic structure and interactions

Track Classification: Hadronic structure and interactions