



Contribution ID: 89

Type: not specified

## Renormalisation of composite operators in lattice QCD: perturbative versus nonperturbative

*Monday, 14 June 2010 17:20 (20 minutes)*

The perturbative and nonperturbative renormalisation of quark-antiquark operators in lattice QCD with two flavours of clover fermions is investigated within the research programme of the QCDSF collaboration. Operators with up to three derivatives are considered. The nonperturbative results based on the RI-MOM scheme are compared with estimates from one- and two-loop lattice perturbation theory.

### Please, insert your presentation type (talk, poster)

talk

**Primary author:** GÖCKELER, Meinulf (Institute for Theoretical Physics, University of Regensburg)

**Co-authors:** SCHÄFER, Andreas (Institute for Theoretical Physics, University of Regensburg); SCHILLER, Arwed (Institute for Theoretical Physics, University of Leipzig); PLEITER, Dirk (DESY Zeuthen); SCHIERHOLZ, Gerrit (DESY Hamburg); STÜBEN, Hinnerk (ZIB Berlin); PERLT, Holger (Institute for Theoretical Physics, University of Leipzig); ZANOTTI, James (School of Physics and Astronomy, University of Edinburgh); RAKOW, Paul (Dept. of Mathematical Sciences, University of Liverpool); HORSLEY, Roger (School of Physics and Astronomy, University of Edinburgh); NAKAMURA, Yoshifumi (Institute for Theoretical Physics, University of Regensburg)

**Presenter:** GÖCKELER, Meinulf (Institute for Theoretical Physics, University of Regensburg)

**Session Classification:** Parallel 08: Standard model parameters and renormalization

**Track Classification:** Standard model parameters and renormalization