



Contribution ID: 122

Type: **not specified**

## The Aoki phase revisited

*Monday, 14 June 2010 14:30 (20 minutes)*

Continuing with the work exposed in the last lattice conference [Phys.Rev.D79:014509,2009, PoS(LAT2009)068], and in order to distinguish between the two different scenarios proposed by us and Sharpe [Phys.Rev.D79:054503,2009], we present data of simulations of QCD with two flavours of dynamical Wilson fermions, inside the Aoki phase and without an external source  $\bar{\psi}\gamma_5\tau_3\psi$ . Two different algorithms, MFA and the hybrid montecarlo, were used, and a consistency check was performed. During the simulations we measured, using the P.D.F. (probability distribution function) formalism, the second moment of the relevant order parameters of the Aoki phase, namely  $\langle(i\bar{\psi}\gamma_5\psi)^2\rangle$  and  $\langle(i\bar{\psi}\gamma_5\tau_3\psi)^2\rangle$ . These numerical results allow us to improve our understanding of the Aoki phase.

### Please, insert your presentation type (talk, poster)

Talk

**Primary authors:** VAQUERO, Alejandro (Universidad de Zaragoza); Dr FOLLANA, Eduardo (Universidad de Zaragoza); Dr DI CARLO, Giuseppe (INFN Laboratorio Gran Sasso); Prof. AZCOITI, Vicente (Universidad de Zaragoza)

**Presenter:** VAQUERO, Alejandro (Universidad de Zaragoza)

**Session Classification:** Parallel 11: Chiral symmetry

**Track Classification:** Chiral symmetry