



Contribution ID: 283

Type: **not specified**

Kaon and D meson masses with $N_f = 2+1+1$ twisted mass lattice QCD

Tuesday, 15 June 2010 18:00 (5 minutes)

We discuss the computation of the K and D meson masses in the $N_f = 2+1+1$ twisted mass lattice QCD setup, where explicit heavy flavour and parity breaking occurs at finite lattice spacing. We present three methods suitable in this context, and verify their consistency.

Please, insert your presentation type (talk, poster)

poster

Primary author: PALLANTE, Elisabetta (University of Groningen)

Co-authors: URBACH, Carsten (University of Bonn); MICHAEL, Chris (University of Liverpool); FARCHIONI, Federico (University of Munster); HERDOIZA, Gregorio (NIC, DESY); MONTVAY, Istvan (DESY, Hamburg); CARBONELL, Jaume (Laboratoire de Physique Subatomique et Cosmologie, Grenoble); JANSEN, Karl (NIC, DESY); WAGNER, Marc (Humboldt University, Berlin); PENE, Olivier (Universite de Paris XI, Orsay); BOUCAUD, Philippe (Universite de Paris XI, Orsay); BARON, Remi (CEA, Centre de Saclay, France); REKER, Siebren (University of Groningen); WENGER, Urs (University of Bern); DRACH, Vincent (Laboratoire de Physique Subatomique et Cosmologie, Grenoble)

Presenter: PALLANTE, Elisabetta (University of Groningen)

Session Classification: Poster session

Track Classification: Hadron spectroscopy