



Contribution ID: 222

Type: **not specified**

## Search for the IR fixed point in the Twisted Polyakov Loop scheme (II)

*Tuesday, 15 June 2010 11:30 (20 minutes)*

We present our updated study of the nonperturbative running coupling constant in Twisted Polyakov Loop (TPL) scheme.

We study the larger lattice size and the lower  $\beta$  region ( $\beta \geq 4.5$ ) in the case of SU(3) Nf=12.

We discuss in detail the step scaling function, the running coupling constant, and the nonperturbative beta function using these updated data.

We also discuss the comparison with the perturbative two-loop results.

### Please, insert your presentation type (talk, poster)

talk

**Primary author:** ITOU, Etsuko (Osaka University)

**Co-authors:** LIN, C.-J.David (National Chiao-Tung University, and National Center for Theoretical Sciences); SHINTANI, Eigo (RIKEN-BNL Research Center); MATSUFURU, Hideo (High Energy Accelerator Research Organization (KEK)); OHKI, Hiroshi (Osaka University); KURACHI, Masafumi (Tohoku University); YAMAZAKI, Takeshi (University of Tsukuba); AOYAMA, Tatsumi (Nagoya University); ONOGI, Tetsuya (Osaka University)

**Presenter:** ITOU, Etsuko (Osaka University)

**Session Classification:** Parallel 14: Applications beyond QCD

**Track Classification:** Applications beyond QCD