

## nazario tantalo

nazario.tantalo@roma2.infn.it

flavour physics workshop, rome, 17-02-2020

# QED corrections to hadronic decays on the lattice

▲□▶ ▲圖▶ ▲臣▶ ▲臣▶ 三臣 - のへで

FLAG, arXiv:1902.08191 PDG review, j.rosner, s.stone, r.van de water, 2016

v.cirigliano et al., Rev.Mod.Phys. 84 (2012)





$$\delta_{QED}\Gamma[\pi^- \to \ell\bar{\nu}(\gamma)] = 1.8\%$$
,

$$\delta_{QED} \Gamma[K^- \to \ell \bar{\nu}(\gamma)] = 1.1\%$$
,

$$\delta_{QED}\Gamma[K \to \pi \ell \bar{\nu}(\gamma)] = [0.5, 3]\%$$

A D > A P > A B > A B >

• at this level of precision QED radiative corrections must be included!



• from the last FLAG review we have

$$f_{\pi\pm} = 130.2(0.8) \text{ MeV} , \qquad \delta = 0.6\%$$
 .

$$f_{K^{\pm}} = 155.7(0.3) \text{ MeV} , \qquad \delta = 0.2\% ,$$

$$f_+(0) = 0.9706(27)$$
,  $\delta = 0.3\%$ 

≡ ∽ < (~

- the most precise value of  $V_{ud}$  comes from super-allowed nuclear  $\beta$ -decays (j.hardy, i.towner, Phys.Rev. C91 (2015)) and the associated QED radiative corrections have an impact on the first-row CKM unitarity check
- by using lattice data for  $f_K/f_{\pi}$  and the phenomenological estimate of w.marciano, a.sirlin, Phys.Rev.Lett. 96 (2006)

$$\sum_{=d,s,b} |V_{uf}|^2 = 0.9999(5)$$

by using c-y.seng et al., Phys.Rev.Lett. 121 (2018)

f

$$\sum_{f=d,s,b} |V_{uf}|^2 = 0.9988(4)$$

by using a.czarnecki et al., Phys.Rev. D100 (2019)

$$\sum_{f=d,s,b} |V_{uf}|^2 = 0.9992(4)$$

• a first-principles calculation is needed here!





・ロト ・ 同ト ・ ヨト ・ ヨト

< □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > <



- presently there are tensions between SM-theory and experiment in observables checking lepton-flavour universality, see f.archilli and m.rotondo talks
- the bulk of the hadronic uncertainties cancel in the ratios but QED radiative corrections are sensitive to the lepton mass and new hadronic quantities are needed at  $O(\alpha)$
- QED effects are taken into account by using PHOTOS but it is not excluded that an improved treatment can have an
  impact, s.de Boer et al PRL 120 (2018), s.cali et al EPJ C79 (2019)
- the analysis of s.de Boer et al PRL 120 (2018) used what in the following is called the point-like effective theory

including QED radiative corrections in a non-perturbative lattice calculation is a challenging problem!

- QED is a long-range unconfined interaction that needs to be consistently defined on a finite volume
- finite-volume effects are potentially very large, e.g. of  $O(L^{-1})$  in the case of the masses of stable hadrons
- in the case of decay rates the problem is much more involved because of the appearance of infrared divergences, O(log(L)), at intermediate stages of the calculation: the infrared problem!
- from the numerical point of view, it is difficult to disentangle QED radiative corrections from the leading QCD contributions that, b.t.w., needs to be properly defined
- as for any other observable on the lattice, QED radiative corrections have to be extracted from euclidean correlators



◆□ > ◆□ > ◆豆 > ◆豆 > □ = − のへで

RM123, JHEP 1204 (2012) RM123, PRD 87 (2013)

• once QCD has been defined, QED radiative corrections can be calculated directly or by expanding the lattice path-integral with respect to  $\alpha \sim (m_d - m_u)/\Lambda_{QCD}$ 

$$\mathcal{O}(\boldsymbol{g}_{s}) = \frac{\langle e^{-S^{full}} O \rangle}{\langle e^{-S^{full}} \rangle} = \frac{\langle e^{-S^{QCD}} \left( e^{-\Delta S} O \right) \rangle}{\langle e^{-S^{QCD}} \left( e^{-\Delta S} \right) \rangle} = \mathcal{O}(\boldsymbol{g}_{s}^{0}) + \Delta \mathcal{O}$$

 the building-blocks for the graphical notation, used as a device to do calculations, are the corrections to the quark propagator

$$\Delta \longrightarrow \pm =$$

$$(e_{f}e)^{2} \xrightarrow{\mathcal{O}} + (e_{f}e)^{2} \xrightarrow{\mathcal{O}} - [m_{f} - m_{f}^{0}] \longrightarrow \mp [m_{f}^{cr} - m_{0}^{cr}] \longrightarrow$$

$$-e^{2}e_{f} \sum_{f_{1}} e_{f_{1}} \xrightarrow{\mathcal{O}} - e^{2} \sum_{f_{1}} e_{f_{1}}^{2} \xrightarrow{\mathcal{O}} - e^{2} \sum_{f_{1}} e_{f_{1}}^{2} \xrightarrow{\mathcal{O}} + e^{2} \sum_{f_{1}} e_{f_{1}} e_{f_{$$

RM123, JHEP 1204 (2012) RM123, PRD 87 (2013)

• once QCD has been defined, QED radiative corrections can be calculated directly or by expanding the lattice path-integral with respect to  $\alpha \sim (m_d - m_u)/\Lambda_{QCD}$ 

$$\mathcal{O}(\boldsymbol{g}_s) = \frac{\langle e^{-S^{full}} O \rangle}{\langle e^{-S^{full}} \rangle} = \frac{\langle e^{-S^{QCD}} \left( e^{-\Delta S} O \right) \rangle}{\langle e^{-S^{QCD}} \left( e^{-\Delta S} \right) \rangle} = \mathcal{O}(\boldsymbol{g}_s^0) + \Delta \mathcal{O}(\boldsymbol{g}_s^0$$

· vacuum polarization effects are the numerical issue with our method

$$\begin{split} \Delta & \longrightarrow \pm = \\ (e_f e)^2 & \underbrace{\sum \bigvee}_{f_1} + (e_f e)^2 & \underbrace{\sum \bigvee}_{f_1} - [m_f - m_f^0] \longrightarrow \mp [m_f^{cr} - m_0^{cr}] & \bullet \\ & -e^2 e_f \sum_{f_1} e_{f_1} & \underbrace{\sum \bigvee}_{f_1} - e^2 \sum_{f_1} e_{f_1}^2 & \underbrace{\sum \bigvee}_{f_1} - e^2 \sum_{f_1} e_{f_1}^2 & \underbrace{\sum \bigvee}_{f_1} + e^2 \sum_{f_1 f_2} e_{f_1} e_{f_2} & \underbrace{\sum \bigvee}_{f_1 f_2} + e^2 \sum_{f_1 f_2} e_{f_1} e_{f_2} & \underbrace{\sum \bigvee}_{f_1} + e^2 \sum_{f_1} e_{f_1} e_{f_2} & \underbrace{\sum \bigvee}_{f_1} + e^2 \sum_{f_1} e_{f_1} e_{f_2} & \underbrace{\sum \bigvee}_{f_1} + e^2 \sum_{f_1} e_{f_1} e_{f_2} & \underbrace{\sum \bigvee}_{f_2} + e^2 \sum_{f_1} e_{f_2} e_{f_1} e_{f_2} & \underbrace{\sum \bigvee}_{f_2} + e^2 \sum_{f_1} e_{f_2} e_{f_1} e_{f_2} & \underbrace{\sum \bigvee}_{f_2} + e^2 \sum_{f_1} e_{f_2} e_{f_1} e_{f_2} & \underbrace{\sum \bigvee}_{f_2} + e^2 \sum_{f_2} e_{f_2} e_{f_2} & \underbrace{\sum \bigvee}_{f_2} & \underbrace{\sum \bigvee}_{f_2} + e^2 \sum_{f_2} e_{f_2} & \underbrace{\sum \bigvee}_{f_2} & \underbrace{\sum \bigcup}_{f_2} & \underbrace{\sum \bigvee}_{f_2} & \underbrace{\sum \bigvee}_{f_2} & \underbrace{\sum \bigvee}_{f_2} & \underbrace{\sum \bigvee}$$

RM123+SOTON collaboration: m.di carlo, d.giusti, v.lubicz, g.martinelli, c.t.sachrajda, f.sanfilippo, s.simula, c.tarantino, n.t.



- i'm now going to show some results of our non-perturbative lattice calculation of the O(α) QED radiative corrections to the decay rates P → ℓν(γ)
- both the theoretical and numerical results discussed below are the outcome of a big effort of the RM123+SOTON collaboration started in 2015 with contributions from other colleagues (m.testa, ...)
- the problem is particularly involved (much more than in the case of the spectrum) because of the appearance of infrared divergences that cancel in physical observables by summing virtual and real photon contributions

f.bloch, a.nordsieck, Phys.Rev. 52 (1937) t.d.lee, m.nauenberg, Phys.Rev. 133 (1964) p.p.kulish, l.d.faddeev, Theor.Math.Phys. 4 (1970)

RM123+SOTON, PRD 91 (2015)

▲□▶ ▲□▶ ▲目▶ ▲目▶ ▲目▶ ● のへで

• let's consider the infrared-safe observable: at  $O(\alpha)$  this is obtained by considering the real contributions with a single photon in the final state

$$\Gamma(E) = \Gamma_0 + e^2 \lim_{L \to \infty} \left\{ \Gamma_V(L) + \Gamma_R(L, E) \right\}$$

• the finite-volume calculation of the real contribution is an issue, momenta are quantized! more to say later on this...

#### RM123+SOTON, PRD 91 (2015)

▲□▶▲□▶▲□▶▲□▶ □ のの⊙

• let's consider the infrared-safe observable: at  $O(\alpha)$  this is obtained by considering the real contributions with a single photon in the final state

$$\Gamma(E) = \Gamma_0 + e^2 \lim_{L \to \infty} \left\{ \Gamma_V(L) + \Gamma_R(L, E) \right\}$$

- the finite-volume calculation of the real contribution is an issue, momenta are quantized! more to say later on this...
- for this reason, by relying on the universality of infrared divergences, it is convenient to rewrite the previous formula as

$$\Gamma(E) = \Gamma_0 + e^2 \lim_{L \to \infty} \left\{ \Gamma_V(L) \underbrace{-\Gamma_V^{pt}(L) + \Gamma_V^{pt}(L) + \Gamma_R^{pt}(L, E) - \Gamma_R^{pt}(L, E)}_{= -\Gamma_R(L, E)} + \Gamma_R(L, E) \right\}$$

where  $\Gamma_{V,R}^{pt}$  are evaluated in the point-like effective theory: these have the same infrared behaviour of  $\Gamma_{V,R}$ 

RM123+SOTON, PRD 91 (2015)

(日) (日) (日) (日) (日) (日) (日) (日)

• let's consider the infrared-safe observable: at  $O(\alpha)$  this is obtained by considering the real contributions with a single photon in the final state

$$\Gamma(E) = \Gamma_0 + e^2 \lim_{L \to \infty} \left\{ \Gamma_V(L) + \Gamma_R(L, E) \right\}$$

- the finite-volume calculation of the real contribution is an issue, momenta are quantized! more to say later on this...
- for this reason, by relying on the universality of infrared divergences, it is convenient to rewrite the previous formula as

$$\Gamma(E) = \Gamma_0 + e^2 \lim_{L \to \infty} \Gamma_V^{SD}(L) + e^2 \lim_{m_\gamma \to 0} \left\{ \Gamma_V^{pt}(m_\gamma) + \Gamma_R^{pt}(m_\gamma, E) \right\} + e^2 \lim_{m_\gamma \to 0} \Gamma_R^{SD}(m_\gamma, E)$$

where  $\Gamma_{V,R}^{pt}$  are evaluated in the point-like effective theory: these have the same infrared behaviour of  $\Gamma_{V,R}$ 

• in the limit of very small photon energies  $\Gamma_R^{SD}(E)$  is negligible because very soft photons cannot resolve the internal structure of an hadron

RM123+SOTON, PRL 120 (2018)

with this method, our result for

$$\Gamma_P(E) = \Gamma_P^0 \left\{ 1 + \delta R_P(E) \right\} \;,$$

$$\delta R_{K\pi} = \delta R_K (E_K^{max}) - \delta R_\pi (E_\pi^{max})$$

is the following

## $\delta \mathbf{R}_{\mathbf{K}\pi}$

$$= -0.0122(10)^{st}(2)^{tun}(8)^{\chi}(5)^{L}(4)^{a}(6)^{qQED}$$

= -0.0122(16)

 this can (with a caveat concerning the definition of QCD) be compared with the result currently quoted by the PDG and obtained in v.cirigliano and h.neufeld, PLB 700 (2011)

$$\delta R_{K\pi} = -0.0112(21)$$



### ETMC gauge configurations

▲□▶ ▲□▶ ▲□▶ ▲□▶ = □ - つへで

$$n_f = 1 + 1 + 1 + 1$$
  
 $a \ge 0.0619(18) \text{ fm}$   
 $m_\pi \ge 223(6) \text{ MeV}$   
 $m_\pi L \le 5.8$ 

#### RM123+SOTON, PRL 120 (2018), PRD 100 (2019)

by defining

$$\Gamma_P(E) = \Gamma_P^0 \left\{ 1 + \delta R_P(E) \right\} \, ,$$

our result are

 $\delta \mathbf{R}_{\mathbf{K}}(\mathbf{E}_{\mathbf{K}}^{\mathbf{max}}) = \mathbf{0.0024}(\mathbf{10})$ 

 $\delta \mathbf{R}_{\pi}(\mathbf{E}_{\pi}^{\mathbf{max}}) = 0.0153(19)$ 

 this can (remember the caveat concerning the definition of QCD) be compared with the result currently quoted by the PDG

$$\delta R_K(E_K^{max}) = 0.0064(24)$$

$$\delta R_{\pi}(E_{\pi}^{max}) = 0.0176(21)$$



◆□ > ◆□ > ◆豆 > ◆豆 > ̄豆 \_ のへで

• i'll now show some results of our on-going non-perturbative lattice calculation of the radiative leptonic decay rates for the processes  $P \mapsto \ell \bar{\nu}_{\ell} \gamma$ 



the RM123+SOTON collaboration:

g.martinelli, University of Rome La Sapienza f.mazzetti, University of Rome La Sapienza m.di carlo, University of Rome La Sapienza

g.m.de divitiis, University of Rome Tor Vergata a.desiderio, University of Rome Tor Vergata r.frezzotti, University of Rome Tor Vergata m.garofalo, INFN of Rome Tor Vergata

> d.giusti, University of Roma Tre v.lubicz, University of Roma Tre f.sanfilippo, INFN of Roma Tre s.simula, INFN of Roma Tre

c.t.sachrajda, University of Southampton

・ロト ・ 同 ト ・ 三 ト ・ 三 ・ うへつ



in our calculation, by using the  $\chi {\rm pt}$  results (v.cirigiliano and i.rosell, PRL 99 (2007)), we estimated the structure dependent real contribution to be, nowadays, phenomenologically irrelevant for  $P=\{\pi,K\}$  and  $\ell=\mu$ 

$$\Gamma_R^{SD}(E) < 0.002 \frac{\Gamma(E) - \Gamma_0}{e^2}$$

▲□▶ ▲□▶ ▲□▶ ▲□▶ = □ - ○○○

m.beneke, j.rohrwild Eur.Phys.J. C71 (2011)



leptonic decay rates for the processes 
$$P\mapsto \ell\bar\nu_\ell\gamma$$

• i'll now show some results of our on-going non-perturbative lattice calculation of the radiative

- as we have seen, in the region of small (soft) photon energies these are needed to properly define the measurable infrared-safe purely leptonic decay rates  $P \mapsto \ell \bar{\nu}_{\ell}(\gamma)$
- in the region of experimentally detectable (hard) photon energies these represent important probes of the internal structure of mesons and alternative (non helicity suppressed) channels for the extraction of CKM matrix elements

$$p = (m_B, \mathbf{0}) , \qquad E_\gamma = O(m_B)$$

$$F_{A,V}(E_{\gamma}) = \frac{q_u f_B m_B}{2E_{\gamma} \lambda_B(\mu)} R(E_{\gamma}, \mu) + O\left(\frac{1}{m_B}\right)$$

$$\frac{1}{\lambda_B(\mu)} = \int_0^\infty d\omega \, \frac{\phi^B_+(\omega,\mu)}{\omega}$$

▲□▶ ▲□▶ ▲豆▶ ▲豆▶ □ 豆 - のへで

 the non-perturbative information needed to compute the radiative decay-rates is encoded into the decay constant of the meson and into two form-factors

$$\epsilon^r_\mu(k)\,\int d^4y\,e^{ik\cdot y}\,\mathrm{T}\langle 0|j^\alpha_W(0)j^\mu_{em}(y)|P(p)\rangle=$$

$$\epsilon^{r}_{\mu}(k) \Bigg\{ -iF_{V} \frac{\varepsilon^{\mu\alpha\gamma\beta} k_{\gamma} p_{\beta}}{m_{P}} \Bigg]$$

$$+ \left[F_A + \frac{m_P f_P}{p \cdot k}\right] \frac{(p \cdot k g^{\mu\alpha} - p^{\mu} k^{\alpha})}{m_P}$$
$$+ \frac{m_P f_P}{p \cdot k} \frac{p^{\mu} p^{\alpha}}{m_P} \right\}$$

these can be expressed as functions of x<sub>γ</sub> (and of m<sub>P</sub>)

$$F_{A,V}(x_{\gamma})$$
,  $0 \le x_{\gamma} = \frac{2p \cdot k}{m_P^2} \le 1$ 

 the infrared divergent contribution (in red) is universal: it is proportional to the amplitude with no photons (f<sub>P</sub>)



 the non-perturbative information needed to compute the radiative decay-rates is encoded into the decay constant of the meson and into two form-factors

$$\epsilon^r_\mu(k)\,\int d^4y\,e^{ik\cdot y}\,\mathrm{T}\langle 0|j^\alpha_W(0)j^\mu_{em}(y)|P(p)\rangle=$$

$$\epsilon^{r}_{\mu}(k) \Biggl\{ -iF_{V} \frac{\varepsilon^{\mu\alpha\gamma\beta} k_{\gamma} p_{\beta}}{m_{P}} \Biggr\}$$

$$\begin{split} &+ \left[F_A + \frac{m_P f_P}{p \cdot k}\right] \frac{(p \cdot k \, g^{\mu \alpha} - p^{\mu} k^{\alpha})}{m_P} \\ &+ \frac{m_P f_P}{p \cdot k} \frac{p^{\mu} p^{\alpha}}{m_P} \bigg\} \end{split}$$

these can be expressed as functions of x<sub>γ</sub> (and of m<sub>P</sub>)

$$F_{A,V}(x_{\gamma})$$
,  $0 \le x_{\gamma} = \frac{2p \cdot k}{m_P^2} \le 1$ 

 the infrared divergent contribution (in red) is universal: it is proportional to the amplitude with no photons (f<sub>P</sub>)



 the non-perturbative information needed to compute the radiative decay-rates is encoded into the decay constant of the meson and into two form-factors

$$\epsilon^r_\mu(k)\,\int d^4y\,e^{ik\cdot y}\,\mathrm{T}\langle 0|j^\alpha_W(0)j^\mu_{em}(y)|P(p)\rangle=$$

$$\epsilon^{r}_{\mu}(k) \Bigg\{ -iF_{V} \frac{\varepsilon^{\mu\alpha\gamma\beta} k_{\gamma} p_{\beta}}{m_{P}} \Bigg]$$

$$\begin{split} &+ \left[F_A + \frac{m_P f_P}{p \cdot k}\right] \frac{(p \cdot k \, g^{\mu \alpha} - p^{\mu} k^{\alpha})}{m_P} \\ &+ \frac{m_P f_P}{p \cdot k} \frac{p^{\mu} p^{\alpha}}{m_P} \right\} \end{split}$$

these can be expressed as functions of x<sub>γ</sub> (and of m<sub>P</sub>)

$$F_{A,V}(x_{\gamma})$$
,  $0 \le x_{\gamma} = \frac{2p \cdot k}{m_P^2} \le 1$ 

 the infrared divergent contribution (in red) is universal: it is proportional to the amplitude with no photons (f<sub>P</sub>)









- the calculation of the QED corrections to (radiative) leptonic decays in the case of B mesons doesn't present any conceptual issue
- cutoff effects are the problem there but strategies to cope with *b*-physics on the lattice exist and can be applied

▲□▶ ▲□▶ ▲□▶ ▲□▶ ▲□ ● ● ●

• we also studied the case in which the **photon is off-shell** (relevant for  $P^- \mapsto \ell^+ \ell^- \ell^- \bar{\nu}$  decays): two more form factors enter the game that, at small virtuality, can be calculated with small modifications of the procedure already followed

- the problem is more challenging in the case of semileptonic decays because, for generic kinematical configurations, the physical observable cannot be extracted from euclidean correlators by the leading exponential contributions (the maiani-testa problem)
- this is a big issue, particularly in the case of B decays, because of the presence of many internal multi-hadron states that can go on-shell with energies smaller than the energy of the external meson-lepton



イロト 不得 とくほ とくほとう

Э

- the problem is more challenging in the case of semileptonic decays because, for generic kinematical configurations, the physical observable cannot be extracted from euclidean correlators by the leading exponential contributions (the maiani-testa problem)
- this is a big issue, particularly in the case of B decays, because of the presence of many internal multi-hadron states that can go on-shell with energies smaller than the energy of the external meson-lepton
- on the other hand, the RM123+SOTON method to cope with infrared divergences can be extended to the case of semileptonic decays, we have already analyzed the problem in great detail



・ロト ・ 理 ト ・ ヨ ト ・ ヨ ト

ъ

- the problem is more challenging in the case of semileptonic decays because, for generic kinematical configurations, the physical observable cannot be extracted from euclidean correlators by the leading exponential contributions (the maiani-testa problem)
- this is a big issue, particularly in the case of B decays, because of the presence of many internal multi-hadron states that can go on-shell with energies smaller than the energy of the external meson-lepton
- on the other hand, the RM123+SOTON method to cope with infrared divergences can be extended to the case of semileptonic decays, we have already analyzed the problem in great detail
- moreover, the problem does not arise at the point (on the boundary of the allowed phase-space)

$$s_{\nu} = (p_B - p_{\nu})^2 = (p_D + p_{\ell})^2 = (m_D + m_{\ell})^2$$

• at (around) this particular kinematical configuration, by calling  $s_D = (p_B - p_D)^2$ , the calculation of the QED radiative corrections to the double-differential decay rate  $d\Gamma/ds_D ds_\nu$  might be feasible!





# backup material

▲□▶ ▲□▶ ▲□▶ ▲□▶ □ ● ● ●

• at  $O(\alpha)$  the systematics associated with the quenching of the zero modes can be understood, for example

$$\begin{split} & \int_{\mu}^{t^{\mu}N^{\mu}u_{1}} = \int_{a}^{\frac{1}{a}} \frac{dk^{0}}{2\pi} \frac{1}{L^{3}} \sum_{k} \frac{1 - \delta_{k,0}}{k^{2}} H^{\mu\alpha}(k) L_{\mu\alpha}(k) , \\ & H^{\mu\alpha}(k) = \int_{a}^{b} d^{4}x e^{ikx} T\langle 0|j_{em}^{\mu}(x) j_{W}^{\alpha}(0)|P(p)\rangle , \\ & L^{\mu\alpha}(k) = \bar{v}_{\nu\ell}\gamma^{\alpha} \frac{1}{i(\not{p}_{\ell} + k) + m_{\ell}} \gamma^{\mu}u_{\ell} \end{split}$$



• at  $O(\alpha)$  the systematics associated with the quenching of the zero modes can be understood, for example

$$\begin{split} & = \int \frac{1}{\alpha} \frac{dk^0}{2\pi} \frac{1}{L^3} \sum_{\mathbf{k}} \frac{1 - \delta_{\mathbf{k},\mathbf{0}}}{k^2} H^{\mu\alpha}(k) L_{\mu\alpha}(k) , \\ H^{\mu\alpha}(k) &= \int d^4x \, e^{ikx} \, T\langle 0 | j^{\mu}_{em}(x) \, j^{\alpha}_W(0) | P(p) \rangle , \\ L^{\mu\alpha}(k) &= \bar{v}_{\nu_{\ell}} \gamma^{\alpha} \frac{1}{i(\not{p}_{\ell} + k) + m_{\ell}} \gamma^{\mu} u_{\ell} \end{split}$$



▲□▶ ▲□▶ ▲□▶ ▲□▶ ▲□ ● ● ●

• the ultraviolet behaviour of this object is given by

~NM-

 at O(\alpha) the systematics associated with the quenching of the zero modes can be understood, for example

$$\begin{split} & \int_{a}^{b^{\mu N \vee h_{a}}} = \int_{a}^{\frac{1}{a}} \frac{dk^{0}}{2\pi} \frac{1}{L^{3}} \sum_{k} \frac{1 - \delta_{k,0}}{k^{2}} H^{\mu\alpha}(k) L_{\mu\alpha}(k) , \\ & H^{\mu\alpha}(k) = \int_{a}^{b} d^{4}x e^{ikx} T\langle 0| j_{em}^{\mu}(x) j_{W}^{\alpha}(0) | P(p) \rangle , \\ & L^{\mu\alpha}(k) = \bar{v}_{\nu\ell} \gamma^{\alpha} \frac{1}{i(\not{p}_{\ell} + k) + m_{\ell}} \gamma^{\mu} u_{\ell} \end{split}$$



▲ロ ▶ ▲ 理 ▶ ▲ 国 ▶ ▲ 国 ■ ● ● ● ● ●

• the ultraviolet behaviour of this object is given by

0000

in the local theory the diagram has a logarithmic divergence (absent with a propagating W) that renormalizes G<sub>F</sub>; the
effect of the zero-modes subtraction is a term

$$\frac{1}{L^3} \int^{\frac{1}{a}} \frac{dk^0}{(k^0)^4} \sim \frac{a^3}{L^3}$$

no new ultraviolet divergences but tricky interplay between cutoff and finite volume effects!

RM123+SOTON, PRL 120 (2018), PRD 100 (2019)



・ロト ・ 日 ・ ・ 日 ・ ・ 日 ・ ・ 日 ・ ・ つ へ ()・

• let's look a bit more in details to the master formula

$$\Gamma(E) = \Gamma_0 + e^2 \lim_{m_\gamma \to 0} \left\{ \Gamma_V^{pt}(m_\gamma) + \Gamma_R^{pt}(m_\gamma, E) \right\} + e^2 \lim_{m_\gamma \to 0} \Gamma_R^{SD}(m_\gamma, E) + e^2 \lim_{L \to \infty} \Gamma_V^{SD}(L)$$



▲□▶▲□▶▲□▶▲□▶ □ のの⊙

RM123+SOTON, PRD 91 (2015), PRD 95 (2017), arXiv:1612.00199

• infrared divergences can be computed in the so called point-like effective theory

$$\mathcal{L}_{pt} = \phi_P^{\dagger} \left\{ -D_{\mu}^2 + m_P^2 \right\} \phi_P + f_P \left\{ 2iG_F V_{CKM} D_{\mu} \phi_P^{\dagger} \bar{\ell} \gamma^{\mu} \nu + \text{h.c.} \right\} \;, \qquad D_{\mu} = \partial_{\mu} - ieA_{\mu}$$

• properly matched effective field theories have, by definition, the same infrared behaviour of the fundamental theory: at leading order the matching is obtained by using  $\Gamma_0$ 

$$\Gamma_0^{pt} = \Gamma_0 = \frac{G_F^2 |V_{CKM}|^2 f_F^2}{8\pi} m_P^3 r_\ell^2 \left(1 - r_\ell^2\right)^2 , \qquad r_\ell = \frac{m_\ell}{m_P} , \qquad D_\mu \mapsto \partial_\mu$$

(日) (日) (日) (日) (日) (日) (日) (日)

RM123+SOTON, PRD 91 (2015), PRD 95 (2017), arXiv:1612.00199

infrared divergences can be computed in the so called point-like effective theory

. . .

$$\mathcal{L}_{pt} = \phi_P^{\dagger} \left\{ -D_{\mu}^2 + m_P^2 \right\} \phi_P + f_P \left\{ 2iG_F V_{CKM} D_{\mu} \phi_P^{\dagger} \bar{\ell} \gamma^{\mu} \nu + \text{h.c.} \right\} \;, \qquad D_{\mu} = \partial_{\mu} - ieA_{\mu}$$

• properly matched effective field theories have, by definition, the same infrared behaviour of the fundamental theory: at leading order the matching is obtained by using  $\Gamma_0$ 

$$\Gamma_0^{pt} = \Gamma_0 = \frac{G_F^2 |V_{CKM}|^2 f_P^2}{8\pi} m_P^3 r_\ell^2 \left(1 - r_\ell^2\right)^2 , \qquad r_\ell = \frac{m_\ell}{m_P} , \qquad D_\mu \mapsto \partial_\mu$$

structure-dependent terms can also be understood in the effective field theory language, e.g.

$$\mathcal{O}_V(x) = F_V \, \epsilon^{\mu\nu\rho\sigma} D_\mu \phi_P \, F_{\nu\rho} \, \bar{\ell}\gamma_\sigma \nu \;, \qquad F_{\nu\rho} = \frac{\partial_\nu}{\partial_\rho} A_\rho - \frac{\partial_\rho}{\partial_\rho} A_\nu \;, \qquad \text{subleading in} \; \frac{E\gamma}{m_\pi}$$

• by exploiting the full set of constraints coming from the WIs and from the e.o.m one can rigorously show that in the expansion around vanishing photon energies both the leading (infrared divergent) and the next-to-leading terms are universal: this implies that  $O(L^{-1})$  finite volume effects are universal (see next slide and backup)

$$\Gamma(E) = \Gamma_0 + e^2 \lim_{m_\gamma \to 0} \left\{ \Gamma_V^{pt}(m_\gamma) + \Gamma_R^{pt}(m_\gamma, E) \right\} + e^2 \lim_{m_\gamma \to 0} \Gamma_R^{SD}(m_\gamma, E) + e^2 \lim_{L \to \infty} \Gamma_V^{SD}(L)$$

• concerning the point-like calculation in infinite volume, we have generalized the results obtained in the early days of quantum field theory by berman 58, kinoshita 59

RM123+SOTON, PRD 91 (2015)

▲□▶▲□▶▲□▶▲□▶ □ のの⊙

$$\Gamma(E) = \Gamma_0 + e^2 \lim_{m_\gamma \to 0} \left\{ \Gamma_V^{pt}(m_\gamma) + \Gamma_R^{pt}(m_\gamma, E) \right\} + e^2 \lim_{m_\gamma \to 0} \Gamma_R^{SD}(m_\gamma, E) + e^2 \lim_{L \to \infty} \Gamma_V^{SD}(L)$$

 concerning the point-like calculation in infinite volume, we have generalized the results obtained in the early days of quantum field theory by berman 58, kinoshita 59
 RM123+SOTON, PRD 91 (2015)

concerning the real SD contribution, we have used χpt results, v.cirigliano and i.rosell, PRL 99 (2007), to show (see backup for non-perturbative results!)

$$\Gamma_R^{SD}(E) < 0.002 \frac{\Gamma(E) - \Gamma_0}{e^2} , \qquad E = E^{max} , \qquad P = \{\pi, K\} , \quad \ell = \mu$$

$$\Gamma(E) = \Gamma_0 + e^2 \lim_{m_\gamma \to 0} \left\{ \Gamma_V^{pt}(m_\gamma) + \Gamma_R^{pt}(m_\gamma, E) \right\} + e^2 \lim_{m_\gamma \to 0} \Gamma_R^{SD}(m_\gamma, E) + e^2 \lim_{L \to \infty} \Gamma_V^{SD}(L)$$

 concerning the point-like calculation in infinite volume, we have generalized the results obtained in the early days of quantum field theory by berman 58, kinoshita 59
 RM123+SOTON, PRD 91 (2015)

concerning the real SD contribution, we have used \(\chi pt\) results, v.cirigliano and i.rosell, PRL 99 (2007), to show (see backup for non-perturbative results!)

$$\Gamma_R^{SD}(E) < 0.002 \frac{\Gamma(E) - \Gamma_0}{e^2} , \qquad E = E^{max} , \qquad P = \{\pi, K\} , \quad \ell = \mu$$

• concerning the point-like finite volume contribution we have calculated the universal infrared logs but also the  $O(L^{-1})$  terms:  $\Gamma_V^{DD}(L)$  has  $O(L^{-2})$  finite volume effects!

RM123+SOTON, PRD 95 (2017), arXiv:1612.00199

$$\left(\frac{1}{L^3}\sum_{\boldsymbol{k}} -\int \frac{d^3k}{(2\pi)^3}\right) \int \frac{dk^0}{2\pi} \frac{1}{k^\beta} \sim O\left(\frac{1}{L^{4-\beta}}\right)$$

◆□ > ◆□ > ◆三 > ◆三 > ・三 ・ のへで

### RM123+SOTON, PRL 120 (2018), PRD 100 (2019)

A D > A P > A B > A B >



- we have performed the lattice calculation by using the RM123 method, i.e. by expanding the lattice path-integral with respect to α and the up-down quark mass difference
- by using this method we managed to obtain excellent numerical signals for the correlators corresponding to the diagrams shown in the figure and for the associated counter-terms
- we have computed non-perturbatively the required renormalization constants in the Rl<sup>'</sup>-MOM scheme and matched them perturbatively with the so-called W-scheme (a.sirlin, NPB 196 (1982): e.braaten and c.s.li PRD 42 (1990)) in which G<sub>F</sub> is defined
RM123+SOTON, PRL 120 (2018), PRD 100 (2019)

化白豆 化硼医 化黄豆 化黄豆 一声



- we have performed the lattice calculation by using the RM123 method, i.e. by expanding the lattice path-integral with respect to α and the up-down quark mass difference
- by using this method we managed to obtain excellent numerical signals for the correlators corresponding to the diagrams shown in the figure and for the associated counter-terms
- we have computed non-perturbatively the required renormalization constants in the Rl<sup>'</sup>-MOM scheme and matched them perturbatively with the so-called W-scheme (a.sirlin, NPB 196 (1982): e.braaten and c.s.li PRD 42 (1990)) in which G<sub>F</sub> is defined
- we have not computed the contributions corresponding to charged sea-quarks; this is the so called electroquenched approximation: although we have estimated the associated uncertainty by using \(\chi pt\), there is certainly room for improvement here...

- it is impossible to have a net electric charge in a periodic box
- this is a consequence of gauss's law

$$S = \int_{L^3} d^4x \left\{ \frac{1}{4} F_{\mu\nu} F_{\mu\nu} + \bar{\psi}_f \left( \gamma_\mu D^f_\mu + m_f \right) \psi_f \right\}$$

$$\partial_k \underbrace{F_{0k}(x)}_{E_k(x)} - \underbrace{ieq_f \bar{\psi}_f \gamma_0 \psi_f(x)}_{e\rho(x)} = 0$$

$$Q = \int_{L^3} d^3 x \, \rho(x) \; = \; \frac{1}{e} \int_{L^3} d^3 x \, \partial_k E_k(x) = 0$$



• one may think to overcome this problem by gauge fixing but large gauge transformations survive a local gauge fixing procedure ( $n \in \mathbb{Z}^4$ )

$$\begin{split} \psi(x) &\mapsto e^{2\pi i \sum_{\mu} \frac{x_{\mu} n_{\mu}}{L_{\mu}}} \psi(x) , \qquad \qquad A_{\mu}(x) \mapsto A_{\mu}(x) + \frac{2\pi n_{\mu}}{L_{\mu}} \\ \psi(x) \bar{\psi}(0) &\mapsto e^{2\pi i \sum_{\mu} \frac{x_{\mu} n_{\mu}}{L_{\mu}}} \psi(x) \bar{\psi}(0) , \qquad \langle \psi(x) \bar{\psi}(0) \rangle = 0 , \qquad x \neq 0 \end{split}$$

in order to study charged particles in a periodic box it has been suggested long ago (duncan et al. 96) to quench (a set of) the zero momentum modes of the gauge field, for example

$$\langle \mathcal{O} \rangle = \int_{\text{pbc in space}} \mathcal{D} \psi \mathcal{D} \bar{\psi} \, \mathcal{D} A_{\mu} \prod_{\mu} \delta \left\{ \int_{TL^3} d^4 x \, A_{\mu}(x) \right\} e^{-S} \, \mathcal{O}$$

- by using this procedure one is also quenching large gauge transformations that are no longer a symmetry and charged particles can propagate
- the assumption is that the induced modifications on the infrared dynamics of the theory should disappear once the infinite volume limit is taken

the point to note is that the resulting finite volume theory, although it may admit an hamiltonian description, is non-local m.havakawa, s.uno Prog.Theor.Phys. 120 (2008) BMW, Science 347 (2015), Phys.Lett. B755 (2016) z.davoudi, m.j.savage PRD90 (2014)

$$\text{QED}_{\text{L}}: \quad \prod_{\mu,t} \delta\left\{ \int_{L^3} d^3x \, A_{\mu}(t, \textbf{\textit{x}}) \right\} \quad \mapsto \quad \int_{\text{pbc in space}} \mathcal{D}\alpha_{\mu}(t) \, e^{-\int_{L^3} d^4x \, \alpha_{\mu}(t) \, A_{\mu}(t, \textbf{\textit{x}})}$$





• QED+QCD isospin breaking effects can be calculated by expanding the lattice path-integral w.r.t.  $e^2$  and  $m_d - m_u$ 

$$M_{\pi^+} - M_{\pi^0} = \frac{(e_a - e_d)^2}{2} e^2 \partial_t \frac{\overleftarrow{\nabla \mathcal{V}} - \overleftarrow{\nabla}}{\overleftarrow{\nabla}}$$

the numerical issue here are quark-disconnected diagrams



BMW, Science 347 (2015), PRL 117 (2016)

- one can also perform simulations of QED+QCD at all orders in  $e^2$  and eventually fit leading isospin breaking effects
- the numerical issue here is that the very small isospin breaking effects come together with the big isosymmetric QCD contributions



b.lucini, a.patella, a.ramos, n.t, JHEP 1602(2016)

• consider C\* boundary conditions (first suggested by wise and polley 91)

$$\psi_f(x+L\mathbf{k}) = C^{-1}\bar{\psi}_f^T(x)$$

$$\bar{\psi}_f(x + L\mathbf{k}) = -\psi_f^T(x)C$$

$$A_{\mu}(x + L\mathbf{k}) = -A_{\mu}(x) , \qquad U_{\mu}(x + L\mathbf{k}) = U_{\mu}^{*}(x) ,$$

- the gauge field is anti-periodic (  $|m{p}| \ge \pi/L)$ : no zero modes by construction!
- · this means no large gauge transformations and

$$Q = \int_{L^3} d^3 x \, \rho(x) \; = \; \frac{1}{e} \int_{L^3} d^3 x \, \partial_k E_k(x) \neq 0$$

a fully gauge invariant formulation is possible: technically this is a consequence of the fact that the electrostatic
potential is unique with anti-periodic boundary conditions

$$\partial_k \partial_k \Phi(\boldsymbol{x}) = \delta^3(\boldsymbol{x}) , \qquad \Phi(\boldsymbol{x} + L\boldsymbol{k}) = -\Phi(\boldsymbol{x})$$



▲□▶▲□▶▲□▶▲□▶ □ のの⊙

• electrically charged states can be probed by considering (Dirac's factor)

$$\Psi_f(t, \boldsymbol{x}) = \underbrace{e^{-iq_f \int d^3 y \, \Phi(\boldsymbol{y} - \boldsymbol{x}) \partial_k A_k(t, \boldsymbol{y})}}_{\Theta(t, \boldsymbol{x})} \, \psi_f(t, \boldsymbol{x}) \,, \qquad \partial_k \partial_k \Phi(\boldsymbol{x}) = \delta^3(\boldsymbol{x})$$

these interpolating operators are invariant under U(1) local gauge transformations

$$\begin{split} \psi_f(x) &\mapsto e^{iq_f\alpha(x)} \psi_f(x) , \qquad A_\mu(x) \mapsto A_\mu(x) + \partial_\mu\alpha(x) , \\ \Theta(t, \boldsymbol{x}) &\mapsto e^{-iq_f \int d^3 y \, \Phi(\boldsymbol{y}-\boldsymbol{x})\partial_k \partial_k \alpha(t, \boldsymbol{y})} \Theta(t, \boldsymbol{x}) = e^{-iq_f \alpha(t, \boldsymbol{x})} \, \Theta(t, \boldsymbol{x}) \end{split}$$

· the gauge factor is not unique, for example one can consider

$$\Psi_f(t, \mathbf{x}) = e^{-iq_f \int_{-\infty}^{x_1} dy \, A_1(t, y, x_2, x_3)} \, \psi_f(t, \mathbf{x}) ,$$

- for any consistent gauge-fixing condition one can build the Dirac factor that provides the unique gauge-invariant extension of matter fields in that gauge
- notice though: interpolating operators can be non-local in space but must be localized in time!

m.hansen et al. JHEP 1805 (2018)



- besides being an attractive theoretical possibility, we have recently shown that  $QED_C$  can be profitably used in numerical applications
- hadron masses can be computed in a fully gauge invariant and local setup with good numerical accuracy
- the RC\* collaboration has developed an open-source code, openQ\*D, that allows to perform full-simulations of QED+QCD with a wide variety of temporal and spatial boundary conditions

## https://gitlab.com/rcstar/openQxD



- QED<sub>L</sub>: very attractive for its formal simplicity; generally, at O(α) the systematics associated with non-localities can be understood
- QED<sub>m</sub>: formally, the simplest way to solve the problem in a local framework is to give a mass to the photon; the L → ∞ limit must be taken before restoring gauge invariance (m<sub>γ</sub> → 0)

m.endres et al. PRL 117 (2016)

 QED<sub>C</sub> a local and fully gauge invariant solution, formally a bit cumbersome, flavour symmetries reduced to discrete subgroups (no spurious operator mixings though) and fully recovered in the infinite volume limit •  $QED_{\infty}$ : at any fixed order in  $\alpha$  radiative corrections can be represented as the convolution of hadronic correlators with QED kernels, e.g.

x.feng et al PRD 100 (2019), LATTICE19

$$O(L) = \int_{L^3} d^4 x \, H^L_{QCD}(x) \, D^L_{\gamma}(x)$$
  
$$\mapsto \int d^4 x \, H_{QCD}(x) \, D_{\gamma}(x)$$

the subtle issue here is the parametrization of the long-distance tails of the hadronic part;

in fact the proposal is an extension of the spectacular applications of the convolution approach to the  $g_{\mu}$  - 2,



.., n.asmussen et al. arXiv:1911.05573

- $QED_L$ : very attractive for its formal simplicity; generally, at  $O(\alpha)$  the systematics associated with non–localities can be understood
- $QED_m$ : formally, the simplest way to solve the problem in a local framework is to give a mass to the photon; the  $L \mapsto \infty$  limit must be taken before restoring gauge invariance  $(m_\gamma \mapsto 0)$

m.endres et al. PRL 117 (2016)

 QED<sub>C</sub> a local and fully gauge invariant solution, formally a bit cumbersome, flavour symmetries reduced to discrete subgroups (no spurious operator mixings though) and fully recovered in the infinite volume limit

• which is the best approach?

•  $QED_{\infty}$ : at any fixed order in  $\alpha$  radiative corrections can be represented as the convolution of hadronic correlators with QED kernels, e.g.

x.feng et al PRD 100 (2019), LATTICE19

$$O(L) = \int_{L^3} d^4 x \, H^L_{QCD}(x) \, D^L_{\gamma}(x)$$
  
$$\mapsto \int d^4 x \, H_{QCD}(x) \, D_{\gamma}(x)$$

the subtle issue here is the parametrization of the long-distance tails of the hadronic part;

in fact the proposal is an extension of the spectacular applications of the convolution approach to the  $g_{\mu}$  – 2,



..., n.asmussen et al. arXiv:1911.05573

- QED<sub>L</sub>: very attractive for its formal simplicity; generally, at  $O(\alpha)$  the systematics associated with non–localities can be understood
- $QED_m$ : formally, the simplest way to solve the problem in a local framework is to give a mass to the photon; the  $L \mapsto \infty$  limit must be taken before restoring gauge invariance  $(m_\gamma \mapsto 0)$

m.endres et al. PRL 117 (2016)

 QED<sub>C</sub> a local and fully gauge invariant solution, formally a bit cumbersome, flavour symmetries reduced to discrete subgroups (no spurious operator mixings though) and fully recovered in the infinite volume limit •  $QED_{\infty}$ : at any fixed order in  $\alpha$  radiative corrections can be represented as the convolution of hadronic correlators with QED kernels, e.g.

x.feng et al PRD 100 (2019), LATTICE19

$$O(L) = \int_{L^3} d^4 x \, H^L_{QCD}(x) \, D^L_{\gamma}(x)$$
  
$$\mapsto \int d^4 x \, H_{QCD}(x) \, D_{\gamma}(x)$$

the subtle issue here is the parametrization of the long-distance tails of the hadronic part;

in fact the proposal is an extension of the spectacular applications of the convolution approach to the  $g_{\mu}$  – 2,



., n.asmussen et al. arXiv:1911.05573

(日) (日) (日) (日) (日) (日) (日) (日)

- which is the best approach?
- in my opinion this is not the relevant point: what really matters is that one must be able to estimate reliably the systematic uncertainties associated with the chosen approach!

▲ロ ▶ ▲周 ▶ ▲ 国 ▶ ▲ 国 ▶ ● の Q @

- in the compact formulation the path-integral is well defined without gauge fixing
- by choosing an unconventional normalization for the U(1) gauge field (action),



$$S = \frac{1}{g^2} \sum_{x,\mu\nu} \operatorname{tr} \left\{ 1 - V_{\mu\nu}(x) \right\} + \frac{36}{2e^2} \sum_{x,\mu\nu} \left\{ 1 - U_{\mu\nu}(x) \right\} + \sum_{f,x} \bar{\psi}_f(x) D[U^{6q}f V] \psi_f(x)$$

$$abla_{\mu}[U^{6q_f}V]\psi_f(x) = U^{6q_f}_{\mu}(x)V_{\mu}(x)\psi_f(x+\mu) - \psi_f(x) , \qquad U_{\mu}(x) = 1 + rac{i}{6}A_{\mu}(x) + \cdots$$

• Dirac's interpolating operators can then be implemented as analytical functions of the link variables, e.g.

$$\Psi_f(x) = \prod_{s=-x_k}^{-1} U_k^{3q_f}(x+sk) \,\psi_f(x) \prod_{s=0}^{L-x_k-1} U_k^{-3q_f}(x+sk)$$

• the mass of, say, the charged kaon can be extracted from the fully gauge invariant correlator

$$\sum_{\mathbf{x}} \left\langle \bar{S}\gamma_5 U(t, \mathbf{x}) \, \bar{U}\gamma_5 S(0) \right\rangle = \frac{Z_K^+(L)}{2M_K^+(L)} e^{-M_K^+(L)t} + \mathcal{O}\left[e^{-\Delta(L)t}\right]$$

 power-law finite volume effects arise when internal states can go on-shell, e.g.

$$k = \frac{2\pi n + \theta}{L}$$
,

$$\Delta \mathcal{O}(p,L) = \mathcal{O}(p,L) - \mathcal{O}(p,\infty)$$

$$=\left(\frac{1}{L^3}\sum_{k}-\int\,\frac{d^3k}{(2\pi)^3}\right)\int\,\frac{dk^0}{2\pi}f_{\mathcal{O}}(p,k)$$



 power-law finite volume effects arise when internal states can go on-shell, e.g.

$$oldsymbol{k} = rac{2\pioldsymbol{n}+oldsymbol{ heta}}{L} \;, \qquad lpha > 0 \;,$$

$$\Delta \mathcal{O}(p,L) = \mathcal{O}(p,L) - \mathcal{O}(p,\infty)$$

$$= \left(\frac{1}{L^3}\sum_{\boldsymbol{k}} -\int \frac{d^3k}{(2\pi)^3}\right) \int \frac{dk^0}{2\pi} f_{\mathcal{O}}(p,k)$$
$$= \left(\frac{1}{L^3}\sum_{\boldsymbol{k}} -\int \frac{d^3k}{(2\pi)^3}\right) \left\{\frac{g_{\mathcal{O}}(p) + O(\boldsymbol{k})}{(\boldsymbol{k} \cdot \boldsymbol{p})^{\alpha}}\right\}$$





▲□▶ ▲□▶ ▲ 臣▶ ★ 臣▶ 三臣 - のへぐ

• power-law finite volume effects arise when internal states can go on-shell, e.g.

$$oldsymbol{k} = rac{2\pioldsymbol{n}+oldsymbol{ heta}}{L} \;, \qquad lpha > 0 \;,$$

$$\Delta \mathcal{O}(p,L) = \mathcal{O}(p,L) - \mathcal{O}(p,\infty)$$

$$= \left(\frac{1}{L^3}\sum_{\boldsymbol{k}} -\int \frac{d^3k}{(2\pi)^3}\right) \int \frac{dk^0}{2\pi} f_{\mathcal{O}}(\boldsymbol{p}, \boldsymbol{k})$$

$$= \left(\frac{1}{L^3}\sum_{\boldsymbol{k}} -\int \frac{d^3k}{(2\pi)^3}\right) \left\{\frac{g_{\mathcal{O}}(p) + O(\boldsymbol{k})}{(\boldsymbol{k} \cdot \boldsymbol{p})^{\alpha}}\right\}$$

$$= \frac{g_{\mathcal{O}}(p)\xi(p,\boldsymbol{\theta})}{\boldsymbol{L}^{3-\alpha}} + O\left(\frac{1}{\boldsymbol{L}^{4-\alpha}}\right) \;,$$





▲ロト ▲圖 ▶ ▲ 国 ▶ ▲ 国 ▶ ● のへで

 power-law finite volume effects arise when internal states can go on-shell, e.g.

$$oldsymbol{k} = rac{2\pioldsymbol{n}+oldsymbol{ heta}}{L} \;, \qquad lpha > 0 \;,$$

$$\Delta \mathcal{O}(p,L) = \mathcal{O}(p,L) - \mathcal{O}(p,\infty)$$

$$= \left(\frac{1}{L^3}\sum_{\boldsymbol{k}} -\int \frac{d^3k}{(2\pi)^3}\right) \int \frac{dk^0}{2\pi} f_{\mathcal{O}}(\boldsymbol{p}, \boldsymbol{k})$$

$$= \left(\frac{1}{L^3}\sum_{\boldsymbol{k}} -\int \frac{d^3k}{(2\pi)^3}\right) \left\{\frac{g_{\mathcal{O}}(p) + O(\boldsymbol{k})}{(\boldsymbol{k} \cdot \boldsymbol{p})^{\alpha}}\right\}$$

$$= \frac{g_{\mathcal{O}}(p)\xi(p,\theta)}{L^{3-\alpha}} + O\left(\frac{1}{L^{4-\alpha}}\right) \;,$$

$$\xi(p, \boldsymbol{\theta}) = \left\{ \sum_{\boldsymbol{n}} - \int \frac{d^3 n}{(2\pi)^3} \right\} \frac{1}{(2\pi \boldsymbol{n} \cdot \boldsymbol{p} + \boldsymbol{\theta} \cdot \boldsymbol{p})^{\alpha}}$$





$$\begin{split} & \left(\frac{1}{L^3}\sum_{k} - \int \frac{d^3k}{(2\pi)^3}\right) \int \frac{dk^0}{2\pi} \frac{1}{k^{\beta}} \\ & \sim O\left(\frac{1}{L^{4-\beta}}\right) \end{split}$$



## • the key point of our method is the universality of infrared divergences

• to see how this works, let's consider the contribution to the decay rate coming from the diagrams shown in the figure

$$\Gamma_V^{P\ell} = \int \frac{d^4k}{(2\pi)^4} \, H^{\alpha\mu}(k,p) \, \frac{1}{k^2} \, \frac{\mathcal{L}_{\alpha\mu}(k)}{2p_\ell \cdot k + k^2}$$

- infrared divergences (and power-law finite volume effects) come from the singularity at  $k^2 = 0$  of the integrand
- the tensor L<sub>αμ</sub> is a regular function, it contains the numerator of the lepton propagator and the appropriate normalization factors

$$\mathcal{L}_{\alpha\mu}(k) \equiv \mathcal{L}_{\alpha\mu}(k, p_{\nu}, p_{\ell}) = O(1)$$

• the hadronic tensor is a QCD quantity

$$H^{\alpha\mu}(k,p) = i \int d^4x \, e^{ik \cdot x} \, T\langle 0 | J^{\alpha}_W(0) \, j^{\mu}(x) \, | P \rangle$$

it satisfies the WIs coming from QED gauge invariance, e.g.

$$k_{\mu} H^{\alpha \mu}(k, p) = -f_P p^{\alpha} ,$$

• and, given the kinematics of the process, it is singular *only* at the single-meson pole





▲□▶ ▲□▶ ▲□▶ ▲□▶ ▲□ ● ● ●

• the hadronic tensor is a QCD quantity

$$H^{\alpha\mu}(k,p) = i \int d^4x \, e^{ik \cdot x} \, T\langle 0 | J^{\alpha}_W(0) \, j^{\mu}(x) \, | P \rangle$$

it satisfies the WIs coming from QED gauge invariance, e.g.

$$k_{\mu} H^{\alpha \mu}(k, p) = -f_P p^{\alpha} ,$$

- and, given the kinematics of the process, it is singular only at the single-meson pole
- the singularity can be isolated by considering the *point-like* tensor, built in such a way to satisfy the same WIs of the full theory

$$H_{pt}^{\alpha\mu}(k,p) = f_P \left\{ \delta^{\alpha\mu} - \frac{(p+k)^{\alpha} \left(2p+k\right)^{\mu}}{2p \cdot k + k^2} \right\},\,$$

$$H_{SD}^{\alpha\mu}(k,p) = H^{\alpha\mu}(k,p) - H_{pt}^{\alpha\mu}(k,p) , \qquad k_{\mu} H_{pt}^{\alpha\mu}(k,p) = -f_{P} p^{\alpha} , \qquad k_{\mu} H_{SD}^{\alpha\mu}(k,p) = 0$$





• the hadronic tensor is a QCD quantity

$$H^{\alpha\mu}(k,p) = i \int d^4x \, e^{ik \cdot x} \, T\langle 0 | J^{\alpha}_W(0) \, j^{\mu}(x) \, | P \rangle$$

it satisfies the WIs coming from QED gauge invariance, e.g.

$$k_{\mu} H^{\alpha \mu}(k, p) = -f_P p^{\alpha} ,$$

- and, given the kinematics of the process, it is singular only at the single-meson pole
- the singularity can be isolated by considering the *point-like* tensor, built in such a way to satisfy the same WIs of the full theory

$$H_{pt}^{\alpha\mu}(k,p) = f_P \left\{ \delta^{\alpha\mu} - \frac{(p+k)^{\alpha} (2p+k)^{\mu}}{2p \cdot k + k^2} \right\},\,$$

$$H_{SD}^{\alpha\mu}(k,p) = H^{\alpha\mu}(k,p) - H_{pt}^{\alpha\mu}(k,p) , \qquad k_{\mu} H_{pt}^{\alpha\mu}(k,p) = -f_{P} p^{\alpha} , \qquad k_{\mu} H_{SD}^{\alpha\mu}(k,p) = 0$$

• the structure dependent contributions are regular and, since there is no constant two-index tensor orthogonal to k,

$$H_{SD}^{\alpha\mu}(k,p) = \left(p \cdot k \,\delta^{\alpha\mu} - k^{\alpha} p^{\mu}\right) F_A + \epsilon^{\alpha\mu\rho\sigma} p_\rho k_\sigma F_V + \dots = O(k)$$





• at  $O(e^2)$  with massive charged particles, singularities arise only at

$$k^{2} = (\pm i|\mathbf{k}|)^{2} + \mathbf{k}^{2} = 0$$

• the blobs on the right are QCD vertexes, e.g.  $\Delta(n + k) \Gamma^{\mu}(n - k) \Delta(n) = -$ 

$$iN(p) \int d^4x d^4y e^{-ipy-ikx} T\langle 0|P(y)j^{\mu}(x)P^{\dagger}(0)|0\rangle ,$$

$$\Delta(p) = N(p) \int d^4 y e^{-ipy} T\langle 0|P(y)P^{\dagger}(0)|0\rangle ,$$

$$N^{-1}(p) = |\langle P(\mathbf{p}) | P^{\dagger}(0) | 0 \rangle|^{2}$$
,

• gauge WIs constrain the first two terms in the expansion, e.g.

$$k_{\mu}\Gamma^{\mu}(p,k) = \Delta^{-1}(p+k) - \Delta^{-1}(p)$$

$$\Gamma^{\mu}(p,k) = 2p^{\mu} + k^{\mu} + O(k^2)$$



• at  $O(e^2)$  with massive charged particles, singularities arise only at

$$k^{2} = (\pm i|\mathbf{k}|)^{2} + \mathbf{k}^{2} = 0$$

• the blobs on the right are QCD vertexes, e.g.

$$\Delta(p+k)\Gamma^{\mu}(p,k)\Delta(p) = \frac{1}{p} \int_{p+k}^{q} d^{4}x d^{4}y e^{-ipy-ikx} T\langle 0|P(y)j^{\mu}(x)P^{\dagger}(0)|0\rangle ,$$
$$\Delta(p) = N(p) \int d^{4}y e^{-ipy} T\langle 0|P(y)P^{\dagger}(0)|0\rangle ,$$
$$N^{-1}(p) = |\langle P(\mathbf{p})|P^{\dagger}(0)|0\rangle|^{2} ,$$

• gauge WIs constrain the first two terms in the expansion, e.g.

$$k_{\mu}\Gamma^{\mu}(p,k) = \Delta^{-1}(p+k) - \Delta^{-1}(p) ,$$
  
$$\Gamma^{\mu}(p,k) = 2p^{\mu} + k^{\mu} + O(k^{2})$$



the first two terms in 1/L are universal!!

$$\left\{\Gamma_V - \Gamma_V^{pt}\right\}(L) = \Gamma_V^{SD}(\infty) + O\left(\frac{1}{L^2}\right)$$

▲□▶ ▲□▶ ▲□▶ ▲□▶ = □ - ○○○

- in order to compare results for QED radiative corrections we must first agree on what we call QCD...
- indeed, when electromagnetic interactions are taken into account the physical theory is QCD+QED
- the QCD action is no longer expected to reproduce physics and, consequently, its renormalization becomes prescription dependent
- a natural matching prescription is to use again physical experimental inputs to set the QCD parameters
- another prescription (j.gasser, a.rusetsky and i.scimemi, EPJ C32 (2003)) consists in imposing the condition that the renormalized couplings of the full theory and QCD are the same, say in the MS scheme at µ = 2 GeV
- in RM123+SOTON, PRL 120 (2018), PRD 100 (2019) we have compared the two approaches and found that the difference, nowadays, is smaller than the statistical uncertainties
- this will rapidly became an important issue on which we should find an agreement



◆ロト ◆昼 → ◆ 臣 ト ◆ 臣 - の へ ()

- it is always a good idea to address the issue of analytical continuation by starting from correlators, it is usually more cumbersome to locate singularities in the amplitudes
- the reason is that correlators (Schwinger's functions) can always be Wick rotated without any problem
- euclidean reduction formulae work straightforwardly only for the lightest states, i.e. the leading exponentials appearing in the correlators, because the corresponding integrals are convergent
- problems arise when one is interested in processes corresponding to non-leading exponentials (notice that at finite L the spectrum of H is discrete)
- the first step in a lattice calculation of a new observable is to understand if the leading exponentials correspond to the external states for the process of interest
- the lightest state appearing in a correlator is readily found by using the quantum numbers of the theory (in p.t. by using the quantum numbers of the *full* theory)

in minkowsky time:

$$C(t) = \mathsf{T}\langle 0| \cdots \bar{O}(t) O(0)|0\rangle$$
  
=  $\langle 0| \cdots e^{-it(H-i\epsilon)} O|0\rangle + \text{o.t.o.}$ 

$$\mathcal{A}(E) = 2E(p^0 - E) \int_0^\infty dt \, e^{ip^0 t} \, C(t) + \text{o.t.o.}$$

in euclidean time:

$$C_E(\tau) = \langle 0 | \cdots e^{-\tau H} O | 0 \rangle + \text{o.t.o}$$

$$\mathcal{A}(E) = -2iE(p^0 - E) \int_0^\infty d\tau \ e^{p^0 \tau} \ C_E(\tau) + \text{o.t.o.}$$

▲ロト ▲帰 ト ▲ ヨ ト ▲ ヨ ト ・ ヨ ・ の Q ()

from the spectral decomposition of correlators at  $O(\alpha)$  one gets expressions that are rather involved but their structure is easy to understand and somehow illuminating

$$C(t) = e^{-tE(p)} \int \frac{d^4q}{(2\pi)^4} A^{virt}(q)$$

$$+\int \frac{d^3q}{(2\pi)^3} A^{real}(q) e^{-t[E(\boldsymbol{p}-\boldsymbol{q})+E_{\gamma}(\boldsymbol{q})]}$$

|q|

m



when the spatial momentum  $\boldsymbol{q}$  of the photon goes to zero we have

$$|\mathbf{q}| \mapsto 0$$
  
 $E(\mathbf{p} - \mathbf{q}) + E_{\gamma}(\mathbf{q}) \mapsto E(\mathbf{p})$   
 $A^{virt}(\mathbf{q}) \mapsto c^{virt} - c_{IR} \log$ 

$$A^{real}(q) \mapsto c^{real} + c_{IR} \log \frac{|q|}{m}$$



for each charged particle emitting a photon one has the exponential corresponding to the charged particle itself as an external state (the virtual photon contribution)

but also the exponential corresponding to the external states with the photon on-shell (the real photon contribution)

since

$$|q| + \sqrt{M^2 + |p - q|^2} \ge \sqrt{M^2 + |p|^2}$$

▲□▶ ▲□▶ ▲□▶ ▲□▶ = □ - ○○○

with an infrared regulator the blue exponentials are sub-leading and, if one is interested in the virtual contribution, there is **no problem of analytical continuation**  in the case of the  $O(e^2)$  QED radiative corrections to the leptonic decays of pseudoscalar mesons



-----

since as we have seen

$$|m{q}| + \sqrt{M^2 + |m{p} - m{q}|^2} \ge \sqrt{M^2 + |m{p}|^2}$$

here there is a problem of analytical continuation! but this diagram can be factorized and the leptonic part can be computed analytically at fixed total momentum and with an infrared regulator the pseudoscalar meson is the lightest state in QED+QCD with the given quantum numbers

therefore, **no problems of analytical continuation** arise in the self-energy diagrams and in the diagram in which the real photon is emitted from the meson!

notice that this is true for a pion but also in the case of flavoured pseudoscalar mesons such as K, B, D!

▲□▶ ▲□▶ ▲□▶ ▲□▶ = □ - ○○○

- problems of analytical continuation do arise in the case of semileptonic decays because of electromagnetic final state interactions
- the internal meson-lepton pair, and eventually multi-hadrons-lepton internal states, can be lighter than the external meson-lepton state
- this is a big issue, particularly in the case of B decays because of the presence of many kinematically-allowed multi-hadron states



▲ロ ▶ ▲周 ▶ ▲ 国 ▶ ▲ 国 ▶ ● の Q @

- problems of analytical continuation do arise in the case of semileptonic decays because of electromagnetic final state interactions
- the internal meson-lepton pair, and eventually multi-hadrons-lepton internal states, can be lighter than the external meson-lepton state
- this is a big issue, particularly in the case of B decays because of the presence of many kinematically-allowed multi-hadron states
- the problem does not arise at the point (on the boundary of the allowed phase-space)

$$s_{\nu} = (p_B - p_{\nu})^2 = (p_D + p_{\ell})^2 = (m_D + m_{\ell})^2$$

• in this particular kinematical configuration, by calling  $s_D = (p_B - p_D)^2$ , the calculation of the QED radiative corrections to the double-differential decay rate  $d\Gamma/ds_D ds_\nu$  might be feasible!





・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・

• the starting point is the hadronic tensor  $(p^2 = m_P^2)$ 

$$H^{\mu\alpha}(k,p) = \int d^4y \, e^{ik \cdot y} \, \mathrm{T}\langle 0 | j^{\alpha}_W(0) j^{\mu}_{em}(y) | P(p) \rangle$$

this can be conveniently decomposed in terms of form-factors as follows

$$H^{\mu\alpha}(k,p) = H^{\mu\alpha}_{SD}(k,p) + H^{\mu\alpha}_{pt}(k,p)$$

$$\begin{split} H^{\mu\alpha}_{SD}(k,p) = &H_1\left[k^2g^{\mu\alpha} - k^{\mu}k^{\alpha}\right] + H_2\left[(p\cdot k - k^2)k^{\mu} - k^2(p-k)^{\mu}\right](p-k)^{\alpha}\\ &- i\frac{F_V}{m_P}\varepsilon^{\mu\alpha\gamma\beta}k_{\gamma}p_{\beta} + \frac{F_A}{m_P}\left[(p\cdot k - k^2)g^{\mu\alpha} - (p-k)^{\mu}k^{\alpha}\right] \end{split}$$

$$H_{pt}^{\mu\alpha}(k,p) = f_P \left[ g^{\mu\alpha} + \frac{(2p-k)^{\mu}(p-k)^{\alpha}}{2p \cdot k - k^2} \right]$$

• the choice of the basis is of course not unique and, moreover, the separation of the point-like contribution can also depend upon the conventions: our definition of  $H_{pt}^{\mu\alpha}(k, p)$  is consistent with the point-like effective lagrangian and it is what we used to compute  $\Gamma_{pt}^{pt}(E)$ ; notice that

$$k_{\mu}H^{\mu\alpha}(k,p) = f_{P} p^{\alpha} , \qquad k_{\mu}H^{\mu\alpha}_{pt}(k,p) = f_{P} p^{\alpha} , \qquad k_{\mu}H^{\mu\alpha}_{SD}(k,p) = 0$$

i.e.  $H_{pt}^{\mu\alpha}(k,p)$  satisfies the same ward identity of the full-theory tensor

• in the case of real photons,  $k^2=0,$  the previous expressions simplify as follows

$$H^{\mu\alpha}(k,p) = H^{\mu\alpha}_{SD}(k,p) + H^{\mu\alpha}_{pt}(k,p)$$

$$\begin{split} H^{\mu\alpha}_{SD}(k,p) = &k^{\mu} \left\{ -H_{1} \, k^{\alpha} + H_{2} \, p \cdot k(p-k)^{\alpha} \right\} \\ &- i \frac{F_{V}}{m_{P}} \varepsilon^{\mu\alpha\gamma\beta} k_{\gamma} p_{\beta} + \frac{F_{A}}{m_{P}} \left[ p \cdot k g^{\mu\alpha} - (p-k)^{\mu} k^{\alpha} \right] \end{split}$$

$$H_{pt}^{\mu\alpha}(k,p) = f_P \left[ g^{\mu\alpha} + \frac{(2p-k)^{\mu}(p-k)^{\alpha}}{2p \cdot k} \right]$$

• the form factors  $H_{1,2}$  do not enter into the physical decay rate for  $P \mapsto \ell \bar{\nu} \gamma$  and can be conveniently separated by considering the projector onto the transverse (and therefore physical) degrees of freedom of the photon that is attached to the vector current

$$n = (1, \mathbf{0}) , \qquad P^{\mu\nu}(k, n) = -g^{\mu\nu} + n^{\mu}n^{\nu} + \frac{[k^{\mu} - n \cdot kn^{\mu}][k^{\nu} - n \cdot kn^{\nu}]}{k^2 - (n \cdot k)^2}$$

▲□▶ ▲圖▶ ▲臣▶ ▲臣▶ 三臣 - のへで

▲□▶▲□▶▲□▶▲□▶ □ のの⊙

the projector P<sup>μν</sup>(k, n) is such that

$$P^{\mu\nu}(k,n)k_{\nu} = P^{\mu\nu}(k,n)n_{\nu} = 0 , \qquad P^{\mu\beta}(k,n)P^{\nu}_{\beta}(k,n) = P^{\mu\nu}(k,n) ,$$

$$P^{\mu\nu}(k,n) = P^{\nu\mu}(k,n) , \qquad \qquad P^{00}(k,n) = P^{0i}(k,n) = 0 ,$$

$$P^{ij}(k,n) = \delta^{ij} - \frac{k^i k^j}{k^2}$$

 in fact P<sup>µν</sup>(k, n) is nothing but the numerator of the photon propagator in the Coulomb's gauge that forbids the propagation of unphysical degrees of freedom; we have

$$P_{\nu\mu}(k,n) H_{SD}^{\mu\alpha}(k,p) = P_{\nu\mu}(k,n) \left\{ -i \frac{F_V}{m_P} \varepsilon^{\mu\alpha\gamma\beta} k_\gamma p_\beta + \frac{F_A}{m_P} \left[ p \cdot k g^{\mu\alpha} - (p-k)^{\mu} k^{\alpha} \right] \right\}$$

• by introducing the polarization vectors as follows (that depend upon n and k)

$$\epsilon_0 = n = (1, \mathbf{0}), \quad \epsilon_{1,2} = (0, \epsilon_{1,2}), \quad \epsilon_3 = (0, \mathbf{k}/|\mathbf{k}|),$$

$$\epsilon_r \cdot \epsilon_s = g_{rs} , \qquad g^{rs} \epsilon_r^\mu \epsilon_s^\nu = g^{\mu\nu}$$

• the projector  $P^{\mu
u}(k,n)$  can be rewritten in terms of the transverse polarization vectors  $\epsilon_{1,2}$  as follows

$$\sum_{r=1,2} \epsilon_r^{\mu} \epsilon_r^{\nu} = P^{\mu\nu}(k,n) \;, \qquad \epsilon_{r,\mu} \; P^{\mu\nu}(k,n) = -\epsilon_r^{\nu} \;, \qquad r=1,2$$

• explicit expressions for the transverse polarization vectors are given below

$$\begin{split} \epsilon_1^{\mu}(\mathbf{k}) &= \left(0, \frac{-k_1 k_3}{|\mathbf{k}| \sqrt{k_1^2 + k_2^2}}, \frac{-k_2 k_3}{|\mathbf{k}| \sqrt{k_1^2 + k_2^2}}, \frac{\sqrt{k_1^2 + k_2^2}}{|\mathbf{k}|}\right) \;,\\ \epsilon_2^{\mu}(\mathbf{k}) &= \left(0, \frac{k_2}{\sqrt{k_1^2 + k_2^2}}, -\frac{k_1}{\sqrt{k_1^2 + k_2^2}}, 0\right) \end{split}$$

▲□▶▲□▶▲□▶▲□▶ □ のの⊙

- in light of the previous discussion, one can either use the (formally) covariant expressions given above for P<sup>μν</sup>(k, n) or the explicit expressions for the transverse polarizations ε<sub>1,2</sub> in order to isolate the physical contributions appearing into H<sup>μα</sup>(k, p)
- in particular, since the axial and vector part of the weak current can be computed separately, we have

$$\epsilon_{r,\mu} H_A^{\mu\alpha}(k,p) = \frac{p \cdot k \, \epsilon_r^\alpha - \epsilon_r \cdot p \, k^\alpha}{m_P} \left\{ F_A + \frac{m_P f_P}{p \cdot k} \right\} + p^\alpha \, \epsilon_r \cdot p \frac{f_P}{p \cdot k} \; ,$$

$$\epsilon_{r,\mu} H_V^{\mu\alpha}(k,p) = i \frac{F_V}{m_P} \varepsilon^{\alpha\mu\gamma\beta} \epsilon_{r,\mu} k_\gamma p_\beta ,$$

r = 1, 2

f.bloch, a.nordsieck, Phys.Rev. 52 (1937) t.d.lee, m.nauenberg, Phys.Rev. 133 (1964) p.p.kulish, l.d.faddeev, Theor.Math.Phys. 4 (1970)

- the infrared problem has been analyzed by many authors over the years
- electrically-charged asymptotic states are not eigenstates of the photon-number operator
- the perturbative expansion of decay-rates and cross-sections with respect to α is cumbersome because of the infinitely many degenerate states
- the block & nordsieck approach consists in lifting the degeneracies by introducing an infrared regulator, say m<sub>γ</sub>, and in computing infrared-safe observables
- at any fixed order in α, infrared-safe observables are obtained by adding the appropriate number of photons in the final states and by integrating over their energy in a finite range, say [0, E]
- in this framework, infrared divergences appear at intermediate stages of the calculations and cancel in the sum of the so-called virtual and real contributions



$$(p+k)^2 + m_P^2 = 2p \cdot k + k^2 \sim 2p \cdot k$$
,

$$\int \frac{d^4k}{(2\pi)^4} \frac{1}{(k^2 + m_{\gamma}^2) \left(2p \cdot k\right) \left(2p_{\ell} \cdot k\right)} \sim c_{IR} \log\left(\frac{m_P}{m_{\gamma}}\right)$$

$$c_{IR}\left\{\log\left(\frac{m_P}{m_\gamma}\right) + \log\left(\frac{m_\gamma}{E}\right)\right\} = c_{IR}\,\log\left(\frac{m_P}{E}\right)$$

▲□ > ▲圖 > ▲目 > ▲目 > → 目 - のへで

## RM123+SOTON, PRD 91 (2015)

 concerning the perturbative point-like calculation in infinite volume, we have generalized the results obtained in the early days of quantum field theory by berman 58, kinoshita 59

$$\Gamma^{pt}(E) = e^2 \lim_{m_\gamma \to \infty} \left\{ \Gamma^{pt}_V(m_\gamma) + \Gamma^{pt}_R(m_\gamma, E) \right\}$$

$$\begin{split} &= \Gamma_0 \; \frac{\alpha_{em}}{4\pi} \; \left\{ 3 \log \left( \frac{m_P^2}{m_W^2} \right) + \log(r_\ell^2) - 4 \log(r_E^2) + \frac{2 - 10 r_\ell^2}{1 - r_\ell^2} \log(r_\ell^2) \\ &\quad -2 \frac{1 + r_\ell^2}{1 - r_\ell^2} \log(r_E^2) \log(r_\ell^2) - 4 \frac{1 + r_\ell^2}{1 - r_\ell^2} \operatorname{Li}_2(1 - r_\ell^2) - 3 \\ &\quad + \frac{3 + r_E^2 - 6 r_\ell^2 + 4 r_E(-1 + r_\ell^2)}{(1 - r_\ell^2)^2} \; \log(1 - r_E) + \frac{r_E(4 - r_E - 4 r_\ell^2)}{(1 - r_\ell^2)^2} \; \log(r_\ell^2) \\ &\quad - \frac{r_E(-22 + 3 r_E + 28 r_\ell^2)}{2(1 - r_\ell^2)^2} - 4 \frac{1 + r_\ell^2}{1 - r_\ell^2} \operatorname{Li}_2(r_E) \; \right\} \,, \end{split}$$

where

$$r_E = \frac{2E}{m_P} , \qquad r_\ell = \frac{m_\ell}{m_P}$$

(日) (日) (日) (日) (日) (日) (日) (日)

- notice that  $\Gamma_V(L)$  and  $\Gamma_V^{pt}(L)$  are ultraviolet divergent in the Fermi theory
- the divergence can be reabsorbed into a renormalization of  $G_F$ , both in the full theory and in the point-like effective theory
- we have analyzed the renormalization of the four-fermion weak operator on the lattice in details and calculated non-perturbatively the renormalization constants in the RI-MOM scheme
- we have then matched the non-perturbative results to the so-called W-regularization at O(α) (a.sirlin, NPB 196 (1982);
   e.braaten and c.s.li PRD 42 (1990))

$$\frac{1}{k^2} \mapsto \frac{1}{k^2} - \frac{1}{k^2 + m_W^2} \;, \quad H_W = \frac{G_F V_{CKM}}{\sqrt{2}} \left\{ 1 + \frac{\alpha}{\pi} \log \frac{m_Z}{m_W} \right\} O_1^{\text{W-reg}} \;,$$

$$O_1^{\mathsf{W}\text{-}\mathsf{reg}} = \sum_{i=1}^5 Z_{1i} O_i^{latt}(a)$$

• indeed, this is the scheme conventionally used to extract  $G_F$  from the muon decay

$$\frac{1}{\tau_{\mu}} = \frac{G_F^2 m_{\mu}^5}{192\pi^3} \left[ 1 - \frac{8m_e^2}{m_{\mu}^2} \right] \left[ 1 + \frac{\alpha}{2\pi} \left( \frac{25}{4} - \pi^2 \right) \right]$$

RM123+SOTON, PRD 91 (2015)



concerning the real structure dependent contributions, the relevant hadronic quantity is

that can be expressed in terms of (two if  $\varepsilon \cdot k = 0$ ) hadronic form-factors (see below)

by using the χpt results (v.cirigliano and i.rosell, PRL 99 (2007)) for these quantities, we have estimated the structure dependent real contribution to be, nowadays, phenomenologically irrelevant for P = {π, K} and ℓ = μ

$$\Gamma_R^{SD}(E) = \lim_{m_\gamma \to 0} \left\{ \Gamma_R(m_\gamma, E) - \Gamma_R^{pt}(m_\gamma, E) \right\} < 0.002 \frac{\Gamma(E) - \Gamma_0}{e^2}$$
RM123+SOTON, PRD 91 (2015)

▲□▶ ▲□▶ ▲□▶ ▲□▶ = □ - ○○○



• concerning the real structure dependent contributions, the relevant hadronic quantity is

that can be expressed in terms of (two if  $\varepsilon \cdot k = 0$ ) hadronic form-factors (see below)

• in the last part of the talk I will show the preliminary results of a fully non-perturbative calculation of the structure dependent real contribution: these confirm the phenomenological analysis for  $P = \{\pi, K\}$  and open the possibility of calculating  $D_{(s)} \mapsto \ell \bar{\nu} \gamma$  and  $B \mapsto \ell \bar{\nu} \gamma$ 

• we performed an analytical calculation of  $\Gamma_V^{pt}(L)$ 

RM123+SOTON, PRD 95 (2017), arXiv:1612.00199

$$\frac{\Gamma_V^{pt}(L) - \Gamma_V^{\ell\ell}(L)}{\Gamma_0} = c_{IR} \log(L^2 m_P^2) + c_0 + \frac{c_1}{(m_P L)} + O\left(\frac{1}{L^2}\right)$$

where

$$c_{IR} = \frac{1}{8\pi^2} \left\{ \frac{(1+r_\ell^2)\log(r_\ell^2)}{(1-r_\ell^2)} + 1 \right\} \;,$$

$$c_0 = \frac{1}{16\pi^2} \left\{ 2 \log \left( \frac{m_P^2}{m_W^2} \right) + \frac{(2 - 6r_\ell^2) \log(r_\ell^2) + (1 + r_\ell^2) \log^2(r_\ell^2)}{1 - r_\ell^2} - \frac{5}{2} \right\} + \frac{\zeta_C(\mathbf{0}) - 2\zeta_C(\boldsymbol{\beta}_\ell)}{2},$$

$$c_1 = -\frac{2(1+r_{\ell}^2)}{1-r_{\ell}^2}\zeta_B(\mathbf{0}) + \frac{8r_{\ell}^2}{1-r_{\ell}^4}\zeta_B(\boldsymbol{\beta}_{\ell})$$

and we have shown that  $c_{IR}$ ,  $c_0$  and  $c_1$  are universal, i.e. they are the same in the point-like and in the full theories! this means that in  $\Gamma_V^{SD}(L) = \Gamma_V(L) - \Gamma_V^{pt}(L)$  we subtract exactly, together with the infrared divergence, the leading O(1/L) terms and we have  $O(1/L^2)$  finite size effects

- notice: the lepton wave-function contribution,  $\Gamma_V^{\ell\ell}(L),$  does not contribute to  $\Gamma_V^{SD}(L)$ 



- the numerical results presented in this talk have been obtained by using the gauge configurations generated and made publicly available by the ETM collaboration
- after the inclusion of QED radiative corrections with the RM123 method, these have  $n_f=1+1+1+1$  dynamical flavours
- 3 different lattice spacings with  $a \ge 0.0619(18)$  fm
- several sea quark masses and volumes with  $m_\pi \geq 223(6)~{\rm MeV}$  and  $m_\pi L \leq 5.8$

ensemble	β	$V/a^4$	$N_{\rm cfg}$	$a\mu_{sea} = a\mu_{ud}$	$a\mu_{\sigma}$	$a\mu_{\delta}$	$a\mu_s$	$M_{\pi}(\text{MeV})$	$M_K(MeV)$	$M_{\pi}L$
A40.40	1.90	$40^3 \times 80$	100	0.0040	0.15	0.19	0.02363	317 (12)	576 (22)	5.7
A30.32	1	$32^3 \times 64$	150	0.0030				275 (10)	568 (22)	3.9
A40.32			100	0.0040				316(12)	578(22)	4.5
A50.32			150	0.0050				350 (13)	586(22)	5.0
A40.24	]	$24^3 \times 48$	150	0.0040				322 (13)	582 (23)	3.5
A60.24			150	0.0060				386 (15)	599(23)	4.2
A80.24			150	0.0080				442 (17)	618 (14)	4.8
A100.24			150	0.0100				495 (19)	639(24)	5.3
A40.20		$20^3 \times 48$	150	0.0040				330(13)	586(23)	3.0
B25.32	1.95	$32^3 \times 64$	150	0.0025	0.135	0.170	0.02094	259 (9)	546 (19)	3.4
B35.32			150	0.0035				302 (10)	555 (19)	4.0
B55.32			150	0.0055				375 (13)	578 (20)	5.0
B75.32			80	0.0075				436 (15)	599 (21)	5.8
B85.24	]	$24^3 \times 48$	150	0.0085				468(16)	613(21)	4.6
D15.48	2.10	$48^3 \times 96$	100	0.0015	0.1200	0.1385	0.01612	223 (6)	529 (14)	3.4
D20.48			100	0.0020				256 (7)	535(14)	3.9
D30.48			100	0.0030				312 (8)	550 (14)	4.7

◆□▶ ◆□▶ ◆臣▶ ◆臣▶ 三臣 - のへで