

A 3D likelihood based approach for extended VHE source extraction



A. Sinha^{1*}, M. Lemoine-Goumard², Y. Gallant¹, V. Marandon³, A. Jardin-Blicq^{4,5}, P. Chambéry²

¹LUPM, Université de Montpellier, IN2P3/CNRS, ²CENBG, Université de Bordeaux, IN2P3/CNRS, ³Max-Planck-Institut für Kernphysik, Heidelberg, ⁴Chulalongkorn University, Bangkok, Thailand, ⁵NARIT, Chiang Mai, Thailand

Efficiency of 3D FoV Background Models for IACTs

Context


Rejection of the residual hadronic background is a key challenge in data analysis for present generation Imaging Atmospheric Cherenkov Telescopes (IACTs). Since this is statistically removed using source free regions in the field of view (FoV), it becomes particularly relevant in the studies of large sources, where the extension is comparable to the FoV of the instrument. The use of FoV background models (eg: as used in [1] to detect extended sources in the galactic plane with H.E.S.S.) is expected to overcome this limitation and yield significant improvement in the sensitivity of IACT to large scale emission.

Aim

Using the IRFs from the [First HESS DL3 Data Release \[2\]](#) and the associated background models supplied in [3], we characterise here the efficiency of a 3D FoV likelihood minimisation as implemented in Gammapy [4] as compared with a traditional ring background estimation for a range of source sizes.


Results

- Consistent results for up to 0.6° source size for all analyses
- Ring estimator fails with increasing source size
- Good stability of reconstructed flux, size for both stacked and joint fits
- Joint fitting convergence time increases non-linearly with the number of observations
- Better stability of reconstructed flux for larger wobble offsets
- Reconstructed background norms correlate well with injected ones



Set-up

- Simulate 20 hrs of observation - 44 runs * 28 mins each
- 4 different wobble positions around the source, different wobble offsets
- Add 15% Gaussian fluctuations on the background normalisation to account for typical background uncertainties.
- Poisson fluctuate source + background
- Simulate sources of different sizes - Gaussian morphology with varying sigma
- Analyse simulated observations using
 - a) Ring Background estimation (radius=1.5°, width=0.3°)
 - b) Stacked analysis - all datasets stacked, 1 free background parameter in total
 - c) Joint analysis - all datasets fitted simultaneously - 1 free background parameter per run - computationally heavy



References

1. Jardin-Blicq et al, *PoS ICRC2019* (2020) 706
2. H.E.S.S. Collaboration [arXiv:1810.04516]
3. L. Mohrmann et al *Astron. Astrophys.* 632 (2019) A72
4. <https://docs.gammapy.org/>
5. H.E.S.S. Collaboration *A&A* 612, A8 (2018)

Acknowledgements

This work is partly supported by ANR 19-CE31-0014-01GAMALO project

*Contact info: atreyee.sinha@umontpellier.fr

Energy dependent morphology

Context



Pulsar wind nebulae and halos are expected to have an energy dependent morphology. However, such signatures can be complicated to detect even using 3D FoV likelihood minimisation

Set-up

- Implement a 3D Energy Dependent Gaussian model (PWNGaussian), with a power law dependence of the size on the energy $\sigma \sim \sigma_0 E^{-\alpha}$
- Simulate HESS observations using the above model and a PowerLaw Spectral dependence for the total spectrum
- Present case, $\alpha = 1$ as expected for a constant velocity advection driven energy loss scenario
- Fit the simulated observation using
 - (A)PWNGaussian Model - reconstructed parameters match injected ones
 - (B)A double Gaussian
 - i) A normal gaussian with a power law spectrum
 - ii) Residuals not flat - add a second gaussian
 - iii) End up with flat residuals!!!
- Compare (A) and (B)

To verify if the observations point to (A) A true energy dependent morphology or (B) two overlapping sources in the same region, a likelihood ratio test cannot be performed since these are non-nested models with different number of free parameters. In this case, we suggest to use the Akaike Information Criterion (AIC)(eg: see, [5], eq 3) to probe the improvement of the fit. In our present case, the AIC yields a significant change in the test statistic ($\Delta \text{AIC} = 98.5$, $p\text{-value} = 3e-22$), correctly pointing to the simulated case of (A).

Investigating a fit with a double Gaussian



Attention!!!
Two overlapping gaussians can mimic an underlying energy dependent morphology