


# Capabilities of the DUNE Near Detector Complex

Tanaz Angelina Mohayai for the DUNE Collaboration  
XIX International Workshop on Neutrino Telescopes  
February 22, 2021

# Outline

- Outlook on Deep Underground Neutrino Experiment, DUNE
  - ★ Overview of its rich physics program
- Near Detector Complex of DUNE
  - ★ Its overall design
  - ★ Overview of the physics that it enables
  - ★ Its components and examples of their capabilities
- Summary



# Deep Underground Neutrino Experiment, a Long-baseline Neutrino Experiment



- 1.2 MW, upgradable to 2.4 MW high-intensity, wide-band neutrino beam
  - ★ Produced at Fermilab and sent to Sanford Underground Research Facility, 1300 km away
- 40 kT liquid Argon time projection chamber far detector
- Highly capable near detector complex:
  - ★ Precise neutrino cross-section measurements and characterization of the spectrum and flavor composition of the beam

# DUNE's Rich Physics Program


- Oscillation physics program:
  - ★ Measurement of the leptonic CP violation
  - ★ Determining the neutrino mass hierarchy
  - ★ Precise measurement of PMNS parameters






# DUNE's Rich Physics Program

- Supernova physics program:
  - ★ Characterization of the time and flavor profile of supernova neutrinos for insight into collapse and evolution of supernova
  - ★ Take advantage of LArTPC's unique sensitivity to  $\nu_e$  flavor




# DUNE's Rich Physics Program

- Beyond standard model program, e.g. baryon number violation:
  - ★ LAr TPC technology well-suited to certain proton decay channels




# DUNE's Rich Physics Program


- Beyond standard model program, e.g. baryon number violation:
  - ★ LAr TPC technology well-suited to certain proton decay channels



For more details on DUNE and its rich physics program,  
see the talk by Georgia Karagiorgi titled **DUNE** on  
February 25



# DUNE Near Detector Complex





- Near detector hall houses various near detector components and enables the DUNE PRISM program:
  - ★ ND-LAr ArgonCube, Liquid Argon time projection chamber
  - ★ ND-GAr, magnetized gaseous Argon time projection chamber surrounded by ECAL calorimeter
  - ★ SAND, system for on-axis neutrino detection

# Physics Enabled by Near Detector Complex – Overview

- Primary goal of the near detector complex:


$$N_{\nu_e}^{FD}(E_{reco}) = \int P_{\nu_\mu \rightarrow \nu_e}(E_\nu) \times \Phi_{\nu_e}(E_\nu) \times \sigma_{\nu_e}(E_\nu) \times \epsilon_{\nu_e}^{FD}(E_\nu) \times S_{\nu_e}^{FD}(E_\nu \rightarrow E_{reco}) dE_\nu$$

- ★ Constraining uncertainties in near to far extrapolation + measure flux,  $\Phi$ , cross section,  $\sigma$ , and  $\nu$ -energy (migration matrix S)



# ND-LAr ArgonCube – Design

- Key design features:
  - ★ Same target nucleus as the far detector, 50t fiducial mass
  - ★ Designed to mitigate high event rates:
    - ▶ Modular design with 35 1m x 1m x 3.5m modules
    - ▶ Pixelated charge readout – LArPix



prototype module 0  
field cage




# ND-LAr ArgonCube – Capabilities


- Key capabilities:

- ★ Collects a high stat independent  $\nu_\mu$  CC interactions on Ar
- ▶ e.g. precisely measures absolute and relative flux using the  $\nu$ -e elastic scattering data

FHC	mode	total	accepted	0.5 GeV to 4.0 GeV	accepted
	$\nu_\mu$ CC	$8.2 \times 10^7$	$3.0 \times 10^7$	$5.9 \times 10^7$	$2.4 \times 10^7$
	$\bar{\nu}_\mu$ CC	$3.6 \times 10^6$	$1.4 \times 10^6$	$1.1 \times 10^6$	$4.6 \times 10^5$
	NC total	$2.8 \times 10^7$	$1.6 \times 10^7$	$1.9 \times 10^7$	$1.3 \times 10^7$
	$\nu_\mu$ CC0 $\pi$	$2.9 \times 10^7$	$1.6 \times 10^7$	$2.6 \times 10^7$	$1.3 \times 10^7$
	$\nu_\mu$ CC1 $\pi^\pm$	$2.0 \times 10^7$	$7.5 \times 10^6$	$1.7 \times 10^7$	$6.0 \times 10^6$
	$\nu_\mu$ CC1 $\pi^0$	$8.0 \times 10^6$	$2.9 \times 10^6$	$6.5 \times 10^6$	$2.2 \times 10^6$
	$\nu_\mu$ CC3 $\pi$	$4.6 \times 10^6$	$7.2 \times 10^5$	$1.7 \times 10^6$	$3.8 \times 10^5$
	$\nu_\mu$ CC other	$9.2 \times 10^6$	$7.4 \times 10^5$	$1.5 \times 10^6$	$3.1 \times 10^5$
	$\nu_e + \bar{\nu}_e$ CC	$1.4 \times 10^6$	$6.6 \times 10^5$	$4.5 \times 10^5$	$3.3 \times 10^5$
	$\nu + e$ elastic	$8.4 \times 10^3$	$7.2 \times 10^3$	$5.3 \times 10^3$	$4.2 \times 10^3$



without ND-LAr  
constrain




with ND-LAr  
constrain

10.1103/PhysRevD.101.032002

# ND-GAr – Design

- Key design features:
  - ★ Has a High Pressure Gas Argon TPC (HPgTPC) at its core; will be a copy of ALICE TPC (acquired ALICE's multiwire chambers will be re-purposed for HPgTPC)
  - ★ Ar-CH<sub>4</sub> 90-10 baseline gas mixture (97% Ar interactions) at 10 atm
  - ★ ECAL calorimeter & superconducting magnet surround the HPgTPC







# ND-GAr – Capabilities

- Key capabilities:

- ★ Designed to reconstruct & sign-tag ND-LAr exiting tracks + collect an independent sample of  $\nu$ -Ar interactions + lower detection threshold than ND-LAr
- ★ PEP-4-like PID & tracking:
  - ▶ e.g. can be used to select exclusive multi- $\pi$  final state samples for further constraining any uncertainties

FHC Beam		RHC Beam	
Process	Events/ton/yr	Process	Events/ton/yr
All $\nu_\mu$ -CC	$1.64 \times 10^6$	All $\bar{\nu}_\mu$ -CC	$5.26 \times 10^5$
CC $0\pi$	$5.85 \times 10^5$	CC $0\pi$	$2.36 \times 10^5$
CC $1\pi^\pm$	$4.09 \times 10^5$	CC $1\pi^\pm$	$1.51 \times 10^5$
CC $1\pi^0$	$1.61 \times 10^5$	CC $1\pi^0$	$4.77 \times 10^4$
CC $2\pi$	$2.10 \times 10^5$	CC $2\pi$	$5.21 \times 10^4$
CC $3\pi$	$9.28 \times 10^4$	CC $3\pi$	$1.66 \times 10^4$
CC $K_s$	$1.20 \times 10^4$	CC $K_s$	$2.72 \times 10^3$
CC $K^\pm$	$4.57 \times 10^4$	CC $K^\pm$	$4.19 \times 10^3$
CC other	$1.27 \times 10^5$	CC other	$1.62 \times 10^4$
All $\bar{\nu}_\mu$ -CC	$7.16 \times 10^4$	All $\nu_\mu$ -CC	$2.72 \times 10^5$
All NC	$5.52 \times 10^5$	All NC	$3.05 \times 10^5$
All $\nu_e$ -CC	$2.85 \times 10^4$	All $\nu_e$ -CC	$1.84 \times 10^4$
$\nu e \rightarrow \nu e$	170	$\nu e \rightarrow \nu e$	120




# ND-GAr – Capabilities

## Key capabilities:

- ★ Reconstructs & sign-tags ND-LAr exiting tracks + collects an independent sample of  $\nu$ -Ar interactions
- ★ PEP-4-like PID & tracking + lower detection threshold than ND-LAr:
  - e.g. can be used to select exclusive multi- $\pi$  final state samples for further constraining



FHC Beam		RHC Beam	
Process	Events/ton/yr	Process	Events/ton/yr
All $\nu_\mu$ -CC	$1.64 \times 10^6$	All $\bar{\nu}_\mu$ -CC	$5.26 \times 10^5$
CC $0\pi$	$5.85 \times 10^5$	CC $0\pi$	$2.36 \times 10^5$
CC $1\pi^\pm$	$4.09 \times 10^5$	CC $1\pi^\pm$	$1.51 \times 10^5$
CC $1\pi^0$	$1.61 \times 10^5$	CC $1\pi^0$	$4.77 \times 10^4$
CC $2\pi$	$2.10 \times 10^5$	CC $2\pi$	$5.21 \times 10^4$
CC $3\pi$	$9.28 \times 10^4$	CC $3\pi$	$1.66 \times 10^4$
CC $K_s$	$1.20 \times 10^4$	CC $K_s$	$2.72 \times 10^3$
CC $K^\pm$	$4.57 \times 10^4$	CC $K^\pm$	$4.19 \times 10^3$
CC other	$1.27 \times 10^5$	CC other	$1.62 \times 10^4$
All $\bar{\nu}_\mu$ -CC	$7.16 \times 10^4$	All $\nu_\mu$ -CC	$2.72 \times 10^5$
All NC	$5.52 \times 10^5$	All NC	$3.05 \times 10^5$
All $\nu_e$ -CC	$2.85 \times 10^4$	All $\nu_e$ -CC	$1.84 \times 10^4$

For more details, see the talk by Federico Battisti on **Physics potential with the DUNE ND-GAr detector** on February 26






# DUNE PRISM

- Move ND-LAr and ND-GAr to various off-axis positions to collect off-axis flux data:
  - ★ Can predict oscillated neutrino event spectra at FD with reduced model dependencies
  - ★ Provides a handle for de-convolving flux & cross section uncertainties



# SAND – Design


- Key design features:
  - ★ Designed to measure the on-axis beam
  - ★ KLOE magnet + ECAL making up the outer layers
  - ★ Central tracking options:
    - ▶ 3D segmented plastic scintillator (3DST) + TPCs
    - ▶ 3DST + Straw Tube Tracker (STT)
    - ▶ STT-only
  - ★ Design is being finalized



# SAND – Capabilities


- With ND-LAr and ND-GAr moving to off-axis positions, SAND remains on-axis to measure any changes in the beam parameters:
  - ★ e.g. can measure the spectral shift in the reconstructed neutrino energy for different horn shifts

## shifted significance



# Summary

- DUNE near detector components and program consist of:
  - ★ ND-LAr
  - ★ ND-GAr
  - ★ SAND
  - ★ DUNE PRISM
- Near detector components and program enable a very precise measurement of oscillation parameters
- The design of the various near detector components add unique and important capabilities to DUNE's overall physics program



Thank you!

Questions are welcome,  
now or on Slack or via  
email ([mtanaz@fnal.gov](mailto:mtanaz@fnal.gov))



# Additional Slides



# ND-GAr – Capabilities

- Key capabilities:

- ★ Lower density ( $\rho_{\text{LAr}}/\rho_{\text{GAr}} \approx 85$  for 10 atm GAr) compared with ND-LAr, more sensitivity to lower energy charged particles that may not be seen in ND-LAr
- ★ Reveals discrepancies between different neutrino event generators for choosing a more accurate  $\nu$ -N interaction model @ lower energies





# ND-GAr – Capabilities

- Key capabilities:

- ★ More specifically, the excellent PID and tracking can help select exclusive multi- $\pi$  final state samples for further reducing any bias in  $\delta_{\text{CP}}$

DUNE Preliminary

