



Contribution ID: 156

Type: **Parallel Contributed Talk**

The day before Betelgeuse dies: identifying a near-earth pre-supernova star from its neutrinos

Thursday, 25 February 2021 18:10 (20 minutes)

I discuss the potential of current and future liquid scintillator neutrino detectors of $\mathcal{O}(10)$ kt mass to localize a presupernova neutrino signal in the sky. In the hours preceding the core collapse of a nearby star (at distance D less than 1 kpc), tens to hundreds of inverse beta decay events will be recorded, and their reconstructed topology in the detector can be used to estimate the direction to the star. Although the directionality of inverse beta decay is weak ($\sim 8\%$ forward-backward asymmetry for currently available liquid scintillators), we find that for a fiducial signal of 200 events (which is realistic for Betelgeuse), a positional error of $\sim 60^\circ$ can be achieved, resulting in the possibility to narrow the list of potential stellar candidates to less than ten, typically. For a configuration with improved forward-backward asymmetry ($\sim 40\%$, as expected for a lithium-loaded liquid scintillator), the angular sensitivity improves to $\sim 15^\circ$, and – when a distance upper limit is obtained from the overall event rate – it is in principle possible to uniquely identify the progenitor star. Any localization information accompanying an early supernova alert will be useful to multi-messenger observations, dark matter detection experiments and to particle physics tests using collapsing stars.

Collaboration name

Primary author: MUKHOPADHYAY, Mainak (Arizona State University)

Co-authors: LUNARDINI, Cecilia; Prof. TIMMES, F.X. (Arizona State University); Prof. ZUBER, Kai (Institute for Nuclear and Particle Physics)

Presenter: MUKHOPADHYAY, Mainak (Arizona State University)

Session Classification: Multimessenger Investigations

Track Classification: Neutrino Telescopes and Multimessenger