



ID contributo: 70

Tipo: non specificato

## Shape and connectivity of clusters: Impact of dynamical state and accretion history

*giovedì 1 luglio 2021 14:40 (25 minuti)*

Matter distribution around clusters is highly anisotropic from their being the nodes of the cosmic web. Clusters' shape and the number of filaments they are connected to, i.e., their connectivity, should reflect the level of anisotropy in the matter distribution and must be, in principle, related to their physical properties.

In this presentation, I investigate the influence of the dynamical state and the formation history on both the shape and local connectivity of about 2400 groups and clusters of galaxies from the large hydrodynamical simulation IllustrisTNG at  $z=0$ . I find that the mass of groups and clusters mainly influences the geometry of the matter distribution: massive halos are significantly more elliptical, and more connected to the cosmic web than low-mass ones.

Beyond the mass-driven effect, ellipticity and connectivity appear to trace different dynamical states, and this is the sign of different accretion histories.

Relaxed groups and clusters are mostly formed long time ago, and slowly accreting matter at the present time. They are rather spherical and weakly connected to their environment. This is mostly because they had enough time to relax and, hence, lost the connection with their preferential directions of accretion and merging. In contrast, late-formed unrelaxed groups and clusters are highly anisotropic with large connectivities and ellipticities. These objects are in formation phase and must be strongly affected by the infalling of materials from filaments.

**Autore principale:** Dr. GOUIN, Celine (Institut d'Astrophysique Spatiale)

**Coautore:** Sig. BONNAIRE, Tony (Institut d'Astrophysique Spatiale); Prof. AGHANIM, Nabila (Institut d'Astrophysique Spatiale)

**Relatore:** Dr. GOUIN, Celine (Institut d'Astrophysique Spatiale)