




MANGO DATA ANALYSES

G. Dho, E. Baracchini, A. Cortez

MANGO SETUP




GEOMETRY STUDY

GEOMETRY

sc_xmean:sc_ymean {(sc_xmean>650 && sc_xmean<1750) && (sc_ymean>370 && sc_ymean<2100)}


GEMs @ 420V

$E_m = 0$ kV/cm




GEOMETRY

sc_xmean:sc_ymean {(sc_xmean>650 && sc_xmean<1750) && (sc_ymean>370 && sc_ymean<2100)}





GEOMETRY: INTEGRAL

sc_integral {sc_xmean>650 && sc_xmean<1750 && sc_ymean>370 && sc_ymean<2100}




GEOMETRY: INTEGRAL

sc_integral (sc_xmean>650 && sc_xmean<1750 && sc_ymean>370 && sc_ymean<2100)




GEOMETRY: ENERGY RESOLUTION AND STATISTICS

Iron peak resolution




Res=peak/sigma

Events in the selection



CURVE GAIN


CURVE GAIN: CUTS




- In the end a simple geometrical cut was used to take the spots in the very centre, also considering the geometrical deformations

CURVE GAIN: CUTS

sc_integral:sc_integral/sc_nhits {sc_integral>5 && sc_integral/sc_nhits<60}




sc_integral:sc_integral/sc_nhits {sc_integral>5 && sc_integral/sc_nhits<60 && (sc_ymean>366 && sc_ymean<2000) && (sc_xmean>900 && sc_xmean<1520)}




CURVE GAIN: CUTS

sc_xmean:sc_ymean {(sc_xmean>650 && sc_xmean<1750) && (sc_ymean>370 && sc_ymean<2100)}




CURVE GAIN: GAIN

- GEM voltages changed together from 400 V to 435 V
- Only a couple of points were analysed with the code after the new Friday's patch (red points)





CURVE GAIN: GAIN

- GEM voltages changed together from 400 V to 435 V
- Only a couple of points were analysed with the code after the new Friday's patch (red points)



CURVE GAIN: ENERGY AND SPATIAL RESOLUTION


Energy Resolution




Consistent value with the geometry study with data taken in **different weeks**

CURVE GAIN: ENERGY AND SPATIAL RESOLUTION

Energy Resolution




Spatial Resolution Using $(l_{gauss}\sigma + t_{gauss}\sigma)/2$



ELECTROLUMINESCENCE


EL: LONG EXPOSURE

- Central region is taken and, after pedestal subtraction, all the light is integrated.
- Being close with the camera and the source not as intense as the one in Frascati, the outcome is fairly dependent on how many iron spots are present in that part of the picture




EL: SHORT EXPOSURE WITH IRON SPOTS ANALYSIS

- Same cuts applied as in the gain part




EL: SHORT EXPOSURE WITH IRON SPOTS ANALYSIS

- Same cuts applied as in the gain part




EL: SHORT EXPOSURE WITH IRON SPOTS ANALYSIS

- Same cuts applied as in the gain part




EL: SHORT EXPOSURE WITH IRON SPOTS ANALYSIS

- Same cuts applied as in the gain part




EL: SHORT EXPOSURE WITH IRON SPOTS ANALYSIS

- Same cuts applied as in the gain part



EL: SHORT EXPOSURE WITH IRON SPOTS ANALYSIS



sc_integral {sc_integral<35000 && sc_integral>0 &&(sc_ymean>950 && sc_ymean<1350) && (sc_xmean>900 && sc_xmean<1540)}



EL: SHORT EXPOSURE WITH IRON SPOTS ANALYSIS


- Are we increasing the EF in the hole by adding a field between GEM and Mesh?

$$E_h = \frac{V_{GEM}}{50\mu m} + E_M$$





EL: SHORT EXPOSURE WITH IRON SPOTS ANALYSIS

- Are we increasing the EF in the hole by adding a field between GEM and Mesh?




EL: SHORT EXPOSURE ENERGY AND SPATIAL RESOLUTION

- Are we increasing the EF in the hole by adding a field between GEM and Mesh?




EL: SHORT EXPOSURE PMT

- Waveforms are integrated from the oscilloscope and the iron peak is found also here




EL: SHORT EXPOSURE PMT

- Waveforms are integrated from the oscilloscope and the iron peak is found also here




PMT more sensitive to UV




Electronic excitation (dissociation into neutral fragments)†	12.5 (10)	12.5 (10)
Dissociative ionization†	15.9	15.9

EL: CHARGE ANALYSIS


0 kV



0,3 kV



12 kV




EL: CHARGE ANALYSIS

- We have the integral of the waveforms calculated by the oscilloscope
- We can select the integral related to the iron spots
- After discussing with Luciano we think that:
Due to the RC circuit we do not read the whole charge but probably only the fast component

Electron time for crossing the GEM–mesh gap	~ 40 ns
Ion time for crossing the GEM–mesh gap	~ ms
RC constant	~ 200 us

EL: ALL TOGETHER




CONCLUSION


- Data was taken with MANGO with iron 55 source at LNGS
- We observe a geometrical dependency of the light output
- Some analysis on the energy and spatial resolution were performed as a function of GEM voltage and induction field
- With GEM at 435 V we do not see clear signs of saturation
- The electroluminescence study is consistent with data taken in Frascati and points at a clear increase in light output with reduced increase in charge without affecting the energy resolution

BACKUP


EL: PMT LIGHT



EL: PMT RESOLUTION



EL: MESH CHARGE



EL: GEM CHARGE

