

# DIFFUSE $\gamma$ RAY CONSTRAINTS ON ANNIHILATING OR DECAYING DM AFTER FERMI

Paolo Panci


Supported by Marie Curie Early Stage Research Training  
(MRTN-CT-2006-035863 - UniverseNet)

UNIVERSITÀ DEGLI STUDI DE L'AQUILA & UNIVERSITÉ PARIS 7




14<sup>th</sup> September 2010, Università del Salento, Lecce

[NPB 840:284-303, 2010](#) in collaboration with M. Cirelli and P.D. Serpico



- Dark Matter Indirect Detection with  $\gamma$  rays
  - Inverse Compton  $\gamma$  rays
  - Prompt  $\gamma$  rays

# PLAN OF THE TALK




- Dark Matter Indirect Detection with  $\gamma$  rays
  - Inverse Compton  $\gamma$  rays
  - Prompt  $\gamma$  rays




- Constraints on Annihilating or Decaying Dark Matter by using the diffuse  $\gamma$  rays measured by the FERMI satellite

# INDIRECT DETECTION ( $\gamma$ RAY CONSTRAINTS)




# INDIRECT DETECTION ( $\gamma$ RAY CONSTRAINTS)




- 1 Prompt  $\gamma$  rays from DM annihilations/decays in the **Galactic Center**, in Dwarf Galaxies and Satellites

# INDIRECT DETECTION ( $\gamma$ RAY CONSTRAINTS)




- 1 Prompt  $\gamma$  rays from DM annihilations/decays in the **Galactic Center**, in Dwarf Galaxies and Satellites
- 2 Radio wave from synchrotron radiation of  $e^+e^-$  produced by DM annihilations/decays in the **GC (very large magnetic field)**


# INDIRECT DETECTION ( $\gamma$ RAY CONSTRAINTS)





- 1 Prompt  $\gamma$  rays from DM annihilations/decays in the **Galactic Center**, in Dwarf Galaxies and Satellites
- 2 Radio wave from synchrotron radiation of  $e^+e^-$  produced by DM annihilations/decays in the **GC (very large magnetic field)**
- 3  $\gamma$  rays from the Inverse Compton Scattering (ICS) of the  $e^+e^-$ , produced by DM annihilations/decays in the **Galactic Halo (GH)**, with the ISRF photons



# INDIRECT DETECTION ( $\gamma$ RAY CONSTRAINTS)




- 1 Prompt  $\gamma$  rays from DM annihilations/decays in the **Galactic Center**, in Dwarf Galaxies and Satellites
- 2 Radio wave from synchrotron radiation of  $e^+e^-$  produced by DM annihilations/decays in the **GC (very large magnetic field)**
- 3  $\gamma$  rays from the Inverse Compton Scattering (ICS) of the  $e^+e^-$ , produced by DM annihilations/decays in the **Galactic Halo (GH)**, with the ISRF photons






# DIFFUSE $\gamma$ RAY EMISSION (FERMI DATA POINTS)

3° lat × 3° lon



Talk by S. Digel

5° lat × 30° lon



Talk by S. Digel


## FERMI DATA (FERMISYMPOSIUM)

2 regions that surround the GC

- 3° latitude × 3° longitude
- 5° latitude × 30° longitude


# DIFFUSE $\gamma$ RAY EMISSION (FERMI DATA POINTS)

$3^\circ \text{ lat} \times 3^\circ \text{ lon}$




Talk by S. Digel

$5^\circ \text{ lat} \times 30^\circ \text{ lon}$




Talk by S. Digel

$10^\circ \text{ lat} - 20^\circ \text{ lat} \times 180^\circ \text{ lon}$



Talk by T. Porter

$60^\circ \text{ lat} - 90^\circ \text{ lat} \times 180^\circ \text{ lon}$



Talk by M. Ackerman

## FERMI DATA (FERMISYMPOSIUM)

2 regions that surround the GC

- $3^\circ$  latitude  $\times$   $3^\circ$  longitude
- $5^\circ$  latitude  $\times$   $30^\circ$  longitude


## FERMI DATA (FERMISYMPOSIUM)

2 regions outside the Galactic Plane

- $10^\circ$ - $20^\circ$  latitude  $\times$   $180^\circ$  longitude
- $60^\circ$ - $90^\circ$  latitude  $\times$   $180^\circ$  longitude


# DIFFUSE $\gamma$ RAY EMISSION (FERMI DATA POINTS)

3° lat × 3° lon




Talk by S. Digel

5° lat × 30° lon




Talk by S. Digel

10° lat – 20° lat × 180° lon



Talk by T. Porter

60° lat – 90° lat × 180° lon



Talk by M. Ackerman

## FERMI DATA (FERMISYMPOSIUM)

2 regions that surround the GC

- 3° latitude × 3° longitude
- 5° latitude × 30° longitude

## FERMI DATA (FERMISYMPOSIUM)

2 regions outside the Galactic Plane

- 10°-20° latitude × 180° longitude
- 60°-90° latitude × 180° longitude

The DM signals do not exceed more than  $3\sigma$  the data

# ICS FLUXES AT EARTH FROM DM ANN/DEC

$$\frac{d\Phi}{d\epsilon} = \frac{1}{\epsilon} \int_{\Delta\Omega} d\Omega \int_{\text{l.o.s.}} ds \frac{j_{\text{tot}}(\epsilon, r(s))}{4\pi}$$

$$\frac{d\Phi}{d\epsilon} = \frac{1}{\epsilon} \int_{\Delta\Omega} d\Omega \int_{1.o.s.} ds \frac{j_{\text{tot}}(\epsilon, r(s))}{4\pi}$$

- $\epsilon$  is the energy of the photon that we detect at Earth

$$\frac{d\Phi}{d\epsilon} = \frac{1}{\epsilon} \int_{\Delta\Omega} d\Omega \int_{\text{l.o.s.}} ds \frac{j_{\text{tot}}(\epsilon, r(s))}{4\pi}$$

- $\epsilon$  is the energy of the photon that we detect at Earth
- $s$  is the coordinate along the line of sight (l.o.s)

$$\frac{d\Phi}{d\epsilon} = \frac{1}{\epsilon} \int_{\Delta\Omega} d\Omega \int_{\text{l.o.s.}} ds \frac{j_{\text{tot}}(\epsilon, r(s))}{4\pi}$$

- $\epsilon$  is the energy of the photon that we detect at Earth
- $s$  is the coordinate along the line of sight (l.o.s)
- $j_{\text{tot}}(\epsilon, r) = j_{\text{p}\gamma}(\epsilon, r) + j_{\text{IC}}(\epsilon, r)$ : is the total emissivity of a cell located at distance  $r$  from the GC

# ICS FLUXES AT EARTH FROM DM ANN/DEC

$$\frac{d\Phi}{d\epsilon} = \frac{1}{\epsilon} \int_{\Delta\Omega} d\Omega \int_{\text{l.o.s.}} ds \frac{j_{\text{tot}}(\epsilon, r(s))}{4\pi}$$

- $\epsilon$  is the energy of the photon that we detect at Earth
- $s$  is the coordinate along the line of sight (l.o.s.)
- $j_{\text{tot}}(\epsilon, r) = j_{\text{p}\gamma}(\epsilon, r) + j_{\text{IC}}(\epsilon, r)$ : is the total emissivity of a cell located at distance  $r$  from the GC

$$j_{\text{p}\gamma}(\epsilon, r) = \epsilon Q_{\gamma}(\epsilon, r)$$

## DM ANNIHILATION

$$Q_{\gamma}^{\text{ann}}(\epsilon, r) = \frac{1}{2} \langle \sigma v \rangle n_{\chi}^2(r) \frac{dN_{\gamma}^{\text{ann}}}{d\epsilon}(\epsilon)$$

## DM DECAY

$$Q_{\gamma}^{\text{dec}}(\epsilon, r) = \Gamma_{\text{dec}} n_{\chi}(r) \frac{dN_{\gamma}^{\text{dec}}}{d\epsilon}(\epsilon)$$

$dN_{\gamma}/d\epsilon$  computed by using the **PYTHIA MonteCarlo code**



$$\frac{d\Phi}{d\epsilon} = \frac{1}{\epsilon} \int_{\Delta\Omega} d\Omega \int_{\text{l.o.s.}} ds \frac{j_{\text{tot}}(\epsilon, r(s))}{4\pi}$$

- $\epsilon$  is the energy of the photon that we detect at Earth
- $s$  is the coordinate along the line of sight (l.o.s)
- $j_{\text{tot}}(\epsilon, r) = j_{\text{p}\gamma}(\epsilon, r) + j_{\text{IC}}(\epsilon, r)$ : is the total emissivity of a cell located at distance  $r$  from the GC

$$j_{\text{IC}}(\epsilon, r) = 2 \int_{m_e}^{m_\chi} dE_e \mathcal{P}(\epsilon, E_e, r) n_e(E_e, r)$$

### DIFFERENTIAL POWER

The derivation is straightforward in terms of the well-known IC kinematics

$$\frac{d\Phi}{d\epsilon} = \frac{1}{\epsilon} \int_{\Delta\Omega} d\Omega \int_{\text{l.o.s.}} ds \frac{j_{\text{tot}}(\epsilon, r(s))}{4\pi}$$

- $\epsilon$  is the energy of the photon that we detect at Earth
- $s$  is the coordinate along the line of sight (l.o.s)
- $j_{\text{tot}}(\epsilon, r) = j_{\text{p}\gamma}(\epsilon, r) + j_{\text{IC}}(\epsilon, r)$ : is the total emissivity of a cell located at distance  $r$  from the GC

$$j_{\text{IC}}(\epsilon, r) = 2 \int_{m_e}^{m_\chi} dE_e \mathcal{P}(\epsilon, E_e, r) n_e(E_e, r)$$

## DIFFERENTIAL POWER

The derivation is straightforward in terms of the well-known IC kinematics

## ELECTRONS NUMBER DENSITY

The derivation can be done by solving the diffusion-loss equation

# DERIVATION OF THE ELECTRONS NUMBER DENSITY

$$\underbrace{-\frac{1}{r^2} \frac{\partial}{\partial r} \left[ r^2 D \frac{\partial f}{\partial r} \right]}_{\text{diffusion}} + \underbrace{v \frac{\partial f}{\partial r}}_{\text{advection}} - \underbrace{\frac{1}{3r^2} \frac{\partial}{\partial r} (r^2 v) \rho \frac{\partial f}{\partial \rho}}_{\text{convection}} + \underbrace{\frac{1}{p^2} \frac{\partial}{\partial p} \left[ \dot{p} p^2 f \right]}_{\text{radiative losses}} = \underbrace{\frac{Q_e(E, r)}{4\pi p^2}}_{\text{source}}$$

$f = n_e(E_e, r)/(4\pi p^2)$  with  $p$  electron momentum

# DERIVATION OF THE ELECTRONS NUMBER DENSITY

$$\underbrace{-\frac{1}{r^2} \frac{\partial}{\partial r} \left[ r^2 D \frac{\partial f}{\partial r} \right]}_{\text{diffusion}} + \underbrace{v \frac{\partial f}{\partial r}}_{\text{advection}} - \underbrace{\frac{1}{3r^2} \frac{\partial}{\partial r} (r^2 v) \rho \frac{\partial f}{\partial p}}_{\text{convection}} + \underbrace{\frac{1}{p^2} \frac{\partial}{\partial p} \left[ \dot{p} p^2 f \right]}_{\text{radiative losses}} = \underbrace{\frac{Q_e(E, r)}{4\pi p^2}}_{\text{source}}$$

$$f = n_e(E_e, r) / (4\pi p^2) \text{ with } p \text{ electron momentum}$$

## FERMI REGIONS

Big Regions of the sky, well outside the GC

$$\theta = 1^\circ \Rightarrow \lambda^\circ = r_\odot \theta \simeq 0.15 \text{ kpc}$$

- 1 0.45 kpc  $\times$  0.45 kpc
- 2 0.74 kpc  $\times$  4.44 kpc
- 3 1.48 kpc - 2.96 kpc  $\times$  26.65 kpc
- 4 8.88 kpc - 13.33 kpc  $\times$  26.65 kpc

# DERIVATION OF THE ELECTRONS NUMBER DENSITY

$$\underbrace{-\frac{1}{r^2} \frac{\partial}{\partial r} \left[ r^2 D \frac{\partial f}{\partial r} \right]}_{\text{diffusion}} + \underbrace{v \frac{\partial f}{\partial r}}_{\text{advection}} - \underbrace{\frac{1}{3r^2} \frac{\partial}{\partial r} (r^2 v) \rho \frac{\partial f}{\partial p}}_{\text{convection}} + \underbrace{\frac{1}{p^2} \frac{\partial}{\partial p} \left[ \dot{p} p^2 f \right]}_{\text{radiative losses}} = \underbrace{\frac{Q_e(E, r)}{4\pi p^2}}_{\text{source}}$$

$$f = n_e(E_e, r) / (4\pi p^2) \text{ with } p \text{ electron momentum}$$

## FERMI REGIONS

Big Regions of the sky, well outside the GC

$$\theta = 1^\circ \Rightarrow \lambda^\circ = r_\odot \theta \simeq 0.15 \text{ kpc}$$

- 1 0.45 kpc  $\times$  0.45 kpc
- 2 0.74 kpc  $\times$  4.44 kpc
- 3 1.48 kpc - 2.96 kpc  $\times$  26.65 kpc
- 4 8.88 kpc - 13.33 kpc  $\times$  26.65 kpc

- Are only important in a region close to the BH accretion disk ( $r_{\text{acc}} \simeq 0.04 \text{ pc}$ )

# DERIVATION OF THE ELECTRONS NUMBER DENSITY

$$\underbrace{-\frac{1}{r^2} \frac{\partial}{\partial r} \left[ r^2 D \frac{\partial f}{\partial r} \right]}_{\text{diffusion}} + \underbrace{v \frac{\partial f}{\partial r}}_{\text{advection}} - \underbrace{\frac{1}{3r^2} \frac{\partial}{\partial r} (r^2 v) \rho \frac{\partial f}{\partial p}}_{\text{convection}} + \underbrace{\frac{1}{p^2} \frac{\partial}{\partial p} \left[ \dot{p} p^2 f \right]}_{\text{radiative losses}} = \underbrace{\frac{Q_e(E, r)}{4\pi p^2}}_{\text{source}}$$

$$f = n_e(E_e, r) / (4\pi p^2) \text{ with } p \text{ electron momentum}$$

## FERMI REGIONS

Big Regions of the sky, well outside the GC

$$\theta = 1^\circ \Rightarrow \lambda^\circ = r_\odot \theta \simeq 0.15 \text{ kpc}$$

- 1 0.45 kpc  $\times$  0.45 kpc
- 2 0.74 kpc  $\times$  4.44 kpc
- 3 1.48 kpc - 2.96 kpc  $\times$  26.65 kpc
- 4 8.88 kpc - 13.33 kpc  $\times$  26.65 kpc

- Are only important in a region close to the BH accretion disk ( $r_{\text{acc}} \simeq 0.04 \text{ pc}$ )
- Approximation for the inner part of our Galaxy ( $\tau_{\text{diff}} \sim \tau_{\text{rad}}$ )  
True outside the Galactic Plane ( $\tau_{\text{diff}} \gg \tau_{\text{rad}}$ )

# DERIVATION OF THE ELECTRONS NUMBER DENSITY

$$\underbrace{-\frac{1}{r^2} \frac{\partial}{\partial r} \left[ r^2 D \frac{\partial f}{\partial r} \right]}_{\text{diffusion}} + \underbrace{v \frac{\partial f}{\partial r}}_{\text{advection}} - \underbrace{\frac{1}{3r^2} \frac{\partial}{\partial r} (r^2 v)}_{\text{convection}} p \frac{\partial f}{\partial p} + \underbrace{\frac{1}{p^2} \frac{\partial}{\partial p} [\dot{p} p^2 f]}_{\text{radiative losses}} = \underbrace{\frac{Q_e(E, r)}{4\pi p^2}}_{\text{source}}$$

$f = n_e(E_e, r)/(4\pi p^2)$  with  $p$  electron momentum

## FERMI REGIONS

Big Regions of the sky, well outside the GC

$$\theta = 1^\circ \Rightarrow \lambda^\circ = r_\odot \theta \simeq 0.15 \text{ kpc}$$

- 1 0.45 kpc  $\times$  0.45 kpc
- 2 0.74 kpc  $\times$  4.44 kpc
- 3 1.48 kpc - 2.96 kpc  $\times$  26.65 kpc
- 4 8.88 kpc - 13.33 kpc  $\times$  26.65 kpc

- Are only important in a region close to the BH accretion disk ( $r_{\text{acc}} \simeq 0.04 \text{ pc}$ )
- Approximation for the inner part of our Galaxy ( $\tau_{\text{diff}} \sim \tau_{\text{rad}}$ )  
True outside the Galactic Plane ( $\tau_{\text{diff}} \gg \tau_{\text{rad}}$ )
- Turn out to be dominated by the **ICS radiative process**

$$n_e(E_e, r) = \frac{1}{b_{\text{tot}}(E_e, r)} \int_{E_e}^{m_\chi} d\tilde{E}_e Q_e(\tilde{E}_e, r)$$



$$n_e(E_e, r) = \frac{1}{b_{\text{tot}}(E_e, r)} \int_{E_e}^{m_\chi} d\tilde{E}_e Q_e(\tilde{E}_e, r)$$

- $b_{\text{tot}}(E_e, r) = b_{\text{CMB}}(E_e) + b_{\text{IR}}(E_e, r) + b_{\text{SL}}(E_e, r)$

# DERIVATION OF THE ELECTRONS NUMBER DENSITY

$$n_e(E_e, r) = \frac{1}{b_{\text{tot}}(E_e, r)} \int_{E_e}^{m_\chi} d\tilde{E}_e Q_e(\tilde{E}_e, r)$$

- $b_{\text{tot}}(E_e, r) = b_{\text{CMB}}(E_e) + b_{\text{IR}}(E_e, r) + b_{\text{SL}}(E_e, r)$
- $Q_e(E_e, r)$ : Source term in the diffusion-loss differential equation

# DERIVATION OF THE ELECTRONS NUMBER DENSITY

$$n_e(E_e, r) = \frac{1}{b_{\text{tot}}(E_e, r)} \int_{E_e}^{m_\chi} d\tilde{E}_e Q_e(\tilde{E}_e, r)$$

- $b_{\text{tot}}(E_e, r) = b_{\text{CMB}}(E_e) + b_{\text{IR}}(E_e, r) + b_{\text{SL}}(E_e, r)$
- $Q_e(E_e, r)$ : Source term in the diffusion-loss differential equation

## DM ANNIHILATION

$$Q_e^{\text{ann}}(E_e, r) = \frac{1}{2} \langle \sigma v \rangle n_\chi^2(r) \frac{dN_e^{\text{ann}}}{dE_e}(E_e)$$

- $\langle \sigma v \rangle$ : Annihilation cross section
- $n_\chi = \rho/m_\chi$ : DM number density
- $dN_e^{\text{ann}}/dE_e$ : Electron spectrum produced by DM annihilation

## DM DECAY

$$Q_e^{\text{dec}}(E_e, r) = \Gamma_{\text{dec}} n_\chi(r) \frac{dN_e^{\text{dec}}}{dE_e}(E_e)$$

- $\Gamma_{\text{dec}} = 1/\tau_{\text{dec}}$ : Decay rate
- $n_\chi = \rho/m_\chi$ : DM number density
- $dN_e^{\text{dec}}/dE_e$ : Electron spectrum produced by DM decay

$dN_e/dE_e$  computed by using the **PYTHIA MonteCarlo code**


# DERIVATION OF THE ELECTRONS NUMBER DENSITY

$$n_e^i(E_e, r) \simeq \frac{1}{b_{\text{tot}}^i(E_e)} \int_{E_e}^{m_\chi} d\tilde{E}_e Q_e(\tilde{E}_e, r)$$


Constant  $b_{\text{tot}}$  over each of the observation regions that we consider

$$b_{\text{tot}}(E_e, r) \simeq b_{\text{tot}}^i(E_e)$$


$3^\circ \times 3^\circ$  &  $5^\circ \times 30^\circ$  Regions



$10^\circ - 20^\circ \times 180^\circ$  Region



$60^\circ - 90^\circ \times 180^\circ$  Region



# ICS FLUXES AT EARTH FROM DM ANN/DEC

$$\frac{d\Phi_i}{d\epsilon \Delta\Omega} = \begin{cases} \mathbf{j}_i^{\text{ann}} \frac{1}{2} \frac{\langle \sigma v \rangle}{4\pi} \frac{\rho_{\odot}^2}{m_{\chi}^2} r_{\odot} \left[ \left. \frac{dN_i^{\text{ann}}}{d\epsilon} \right|_{\text{IC}} + \left. \frac{dN^{\text{ann}}}{d\epsilon} \right|_{\text{p}\gamma} \right] & \text{(annihilation)} \\ \mathbf{j}_i^{\text{dec}} \frac{\Gamma_{\text{dec}}}{4\pi} \frac{\rho_{\odot}}{m_{\chi}} r_{\odot} \left[ \left. \frac{dN_i^{\text{dec}}}{d\epsilon} \right|_{\text{IC}} + \left. \frac{dN^{\text{dec}}}{d\epsilon} \right|_{\text{p}\gamma} \right] & \text{(decay)} \end{cases}$$

$$\left. \frac{dN_i^{\text{ann/dec}}}{d\epsilon} \right|_{\text{IC}} = \frac{2}{\epsilon} \int_{m_e}^{m_{\chi}} dE_e \frac{\mathcal{P}_i(E_e, \epsilon)}{b_{\text{tot}}^i(E_e)} \int_{E_e}^{m_{\chi}} d\tilde{E}_e \frac{dN_e^{\text{ann/dec}}}{d\tilde{E}_e}, \quad \left. \frac{dN^{\text{ann/dec}}}{d\epsilon} \right|_{\text{p}\gamma} = \frac{dN_{\gamma}^{\text{ann/dec}}}{d\epsilon}.$$

# ICS FLUXES AT EARTH FROM DM ANN/DEC

$$\frac{d\Phi_i}{d\epsilon \Delta\Omega} = \begin{cases} \bar{J}_i^{\text{ann}} \frac{1}{2} \frac{\langle\sigma v\rangle}{4\pi} \frac{\rho_\odot^2}{m_\chi^2} r_\odot \left[ \left. \frac{dN_i^{\text{ann}}}{d\epsilon} \right|_{\text{IC}} + \left. \frac{dN^{\text{ann}}}{d\epsilon} \right|_{\text{P}\gamma} \right] & \text{(annihilation)} \\ \bar{J}_i^{\text{dec}} \frac{\Gamma_{\text{dec}}}{4\pi} \frac{\rho_\odot}{m_\chi} r_\odot \left[ \left. \frac{dN_i^{\text{dec}}}{d\epsilon} \right|_{\text{IC}} + \left. \frac{dN^{\text{dec}}}{d\epsilon} \right|_{\text{P}\gamma} \right] & \text{(decay)} \end{cases}$$

$$\left. \frac{dN_i^{\text{ann/dec}}}{d\epsilon} \right|_{\text{IC}} = \frac{2}{\epsilon} \int_{m_e}^{m_\chi} dE_e \frac{\mathcal{P}_i(E_e, \epsilon)}{b_{\text{tot}}^i(E_e)} \int_{E_e}^{m_\chi} d\tilde{E}_e \frac{dN_e^{\text{ann/dec}}}{d\tilde{E}_e}, \quad \left. \frac{dN^{\text{ann/dec}}}{d\epsilon} \right|_{\text{P}\gamma} = \frac{dN_\gamma^{\text{ann/dec}}}{d\epsilon}.$$

## ANNIHILATION SCENARIO

$$\bar{J}_i^{\text{ann}} = \int \frac{ds}{r_\odot} \frac{\rho^2[r(s, b, l)]}{\rho_\odot^2}$$

$b \rightarrow$  Galactic latitude

$l \rightarrow$  Galactic longitude

## DECAY SCENARIO

$$\bar{J}_i^{\text{dec}} = \int \frac{ds}{r_\odot} \frac{\rho[r(s, b, l)]}{\rho_\odot}$$

$b \rightarrow$  Galactic latitude

$l \rightarrow$  Galactic longitude







# BOOST FACTORS IN OUR REGIONS

Region	latitude $b$ & longitude $l$	$\bar{j}_i^{\text{ann}}$			$\bar{j}_i^{\text{dec}}$		
		IsoT	NFW	Einasto	IsoT	NFW	Einasto
'3×3'	$0.25^\circ <  b  < 2.75^\circ$ $357.25^\circ < l < 359.75^\circ$ $0.25^\circ < l < 2.75^\circ$	35.1	325	595	8.20	14.6	18.5
'5×30'	$0.25^\circ <  b  < 4.75^\circ$ $330.25^\circ < l < 359.75^\circ$ $0.25^\circ < l < 29.75^\circ$	20.9	51.1	85.9	6.56	7.53	8.72
'10–20'	$10^\circ <  b  < 20^\circ$ $0^\circ < l < 360^\circ$	3.35	3.46	4.23	2.45	2.38	2.47
'Gal Poles'	$60^\circ <  b  < 90^\circ$ $0^\circ < l < 360^\circ$	0.92	0.96	0.94	1.74	1.69	1.67

In the **DM annihilation scenario** the signals from the inner part of our Galaxy are boosted compare to the **decay one**

More competitive  $\gamma$  ray constraints in the DM annihilation scenario

# SUMMARY & RESULTS (DM ANNIHILATION)





# DRAW THE EXCLUSION LINES $\langle \sigma_{\text{ann}} \mathbf{v} \rangle / m_\chi$ PLANE

# DRAW THE EXCLUSION LINES $\langle \sigma_{\text{ann}} \mathbf{v} \rangle / m_\chi$ PLANE


Recent Numerical Simulations (Einasto profile):

$$\rho_{\text{Ein}}(r) = \rho_s \exp \left[ -\frac{2}{\alpha} \left( \left( \frac{r}{r_s} \right)^\alpha - 1 \right) \right], \quad \alpha = 0.17.$$

Previously standard choices (NFW & IsoT):

$$\rho_{\text{NFW}}(r) = \rho_s \frac{r_s}{r} \left( 1 + \frac{r}{r_s} \right)^{-2}, \quad \rho_{\text{isoT}}(r) = \frac{\rho_s}{1 + (r/r_s)^2}$$

DM halo model	$r_s$ in kpc	$\rho_s$ in $\text{GeV}/\text{cm}^3$
NFW	20.0	0.26
Einasto	21.8	0.05
Isothermal	3.20	2.31



# DRAW THE EXCLUSION LINES $\langle\sigma_{\text{ann}}v\rangle/m_\chi$ PLANE


- Consider a Benchmark DM Halo profile
- Calculate the **ICS signal** and the **prompt signal** in each given primary annihilation channel spanning the DM mass in a range between **100 GeV up to 10 TeV**

# DRAW THE EXCLUSION LINES $\langle\sigma_{\text{ann}}v\rangle/m_\chi$ PLANE

- Consider a Benchmark DM Halo profile
- Calculate the **ICS signal** and the **prompt signal** in each given primary annihilation channel spanning the DM mass in a range between **100 GeV up to 10 TeV**
- Require that the DM signals do not exceed more than  $3\sigma$  the **FERMI experimental data**

# IC + PROMPT $\gamma$ CONSTRAINTS (DM ANNIHILATION)

DM DM  $\rightarrow \tau\tau$ , Einasto profile



P.P., M. Cirelli, P.D. Serpico


The PAMELA allowed region (green 95% C.L. and yellow 99.999% C.L.)

FERMI + HESS + PAMELA allowed region (red 95% C.L. and orange 99.999% C.L.)


are completely excluded by the IC + Prompt  $\gamma$  constraints !!!

# IC + PROMPT $\gamma$ CONSTRAINTS (DM ANNIHILATION)


DM DM  $\rightarrow ee$ , Einasto profile




DM DM  $\rightarrow \mu\mu$ , Einasto profile




DM DM  $\rightarrow \tau\tau$ , Einasto profile




DM DM  $\rightarrow ee$ , NFW profile



DM DM  $\rightarrow \mu\mu$ , NFW profile




DM DM  $\rightarrow \tau\tau$ , NFW profile



P.P., M. Cirelli, P.D. Serpico



# IC + PROMPT $\gamma$ CONSTRAINTS (DM ANNIHILATION)




P.P., M. Cirelli, P.D. Serpico

For the smooth isothermal profile, regions of the parameters space seem to be reopened.


The FERMI + HESS + PAMELA allowed region in the case of annihilation into muons is not excluded yet

# HADRONIC MODE CONSTRAINTS


DM DM  $\rightarrow$  bb, Einasto profile




DM DM  $\rightarrow$  tt, Einasto profile




DM DM  $\rightarrow$  WW, Einasto profile




DM DM  $\rightarrow$  bb, NFW profile



DM DM  $\rightarrow$  tt, NFW profile



DM DM  $\rightarrow$  WW, NFW profile




P.P., M. Cirelli, P.D. Serpico



# DRAW THE EXCLUSION LINES $\tau_{\text{dec}}/m_\chi$ PLANE

# DRAW THE EXCLUSION LINES $\tau_{\text{dec}}/m_\chi$ PLANE

## ■ IC + Prompt $\gamma$ Constraints from our Galaxy



P.P., M. Cirelli, P.D. Serpico

The exclusion lines lie below the FERMI+HESS+PAMELA allowed region

The constraints from our Galaxy are not so strong

# DRAW THE EXCLUSION LINES $\tau_{\text{dec}}/m_\chi$ PLANE

- IC + Prompt  $\gamma$  Constraints from the residual "Isotropic radiation"

# DRAW THE EXCLUSION LINES $\tau_{\text{dec}}/m_\chi$ PLANE

- IC + Prompt  $\gamma$  Constraints from the residual "Isotropic radiation"

$$\frac{d\Phi_{\text{cosm}}^{\text{dec}}}{d\epsilon} = \Gamma_{\text{dec}} \frac{\Omega_\chi \rho_{c,0}}{m_\chi} \frac{1}{H_0} \int_0^\infty dz \frac{e^{-\tau(\epsilon,z)}}{\sqrt{\Omega_M(1+z)^3 + \Omega_\Lambda}} \left[ \left. \frac{dN^{\text{dec}}}{d\epsilon} \right|_{\text{IC}} + \left. \frac{dN^{\text{dec}}}{d\epsilon} \right|_{\text{p}\gamma} \right]$$

# DRAW THE EXCLUSION LINES $\tau_{\text{dec}}/m_\chi$ PLANE


- IC + Prompt  $\gamma$  Constraints from the residual "Isotropic radiation"

$$\frac{d\Phi_{\text{isotropic}}^{\text{dec}}}{d\epsilon} = \frac{d\Phi_{\text{cosm}}^{\text{dec}}}{d\epsilon} + 4\pi \frac{d\Phi_{\text{halo}}^{\text{dec}}}{d\epsilon d\Omega} \Big|_{\text{Anti-GC}}$$

# DRAW THE EXCLUSION LINES $\tau_{\text{dec}}/m_\chi$ PLANE

- IC + Prompt  $\gamma$  Constraints from the residual "Isotropic radiation"

$$\frac{d\Phi_{\text{isotropic}}^{\text{dec}}}{d\epsilon} = \frac{d\Phi_{\text{cosm}}^{\text{dec}}}{d\epsilon} + 4\pi \frac{d\Phi_{\text{halo}}^{\text{dec}}}{d\epsilon d\Omega} \Big|_{\text{Anti-GC}}$$




Talk by M. Ackerman, FermiSymposium

The "Isotropic Signal" does not exceed more than  $3\sigma$  the FERMI data in the Anti-GC

- IC + Prompt  $\gamma$  Constraints from the residual "Isotropic radiation"

$$\frac{d\Phi_{\text{isotropic}}^{\text{dec}}}{d\epsilon} = \frac{d\Phi_{\text{cosm}}^{\text{dec}}}{d\epsilon} + 4\pi \frac{d\Phi_{\text{halo}}^{\text{dec}}}{d\epsilon d\Omega} \Big|_{\text{Anti-GC}}$$



Talk by M. Ackerman, FermiSymposium


The "Isotropic Signal" does not exceed more than  $3\sigma$  the FERMI data in the Anti-GC

## ANNIHILATION SCENARIO

- Stronger dependence on the angular distance from the GC is introduced in the galactic flux (no longer "Isotropic signal")
- Dependence on the DM profiles and the clumpiness of DM halos is introduced in the cosmological flux (strong dependence on  $M_{\text{min}}^{\text{Halo}}$ ,  $c_{\text{vir}}$ )

see e.g. Zaharijas et al. JCAP 1004:014,2010

# IC + PROMPT $\gamma$ CONSTRAINTS (DM DECAY)



P.P., M. Cirelli, P.D. Serpico

The residual "Isotropic radiation" measured by FERMI imposes the strongest constraints

It excludes the decay explanation of the FERMI+HESS+PAMELA anomalies for the  $\tau\tau$  channel and starts to exclude the decay explanation for the  $\mu\mu$  channel



# CONCLUSIONS

## Leptonic Annihilation modes:

- For the NFW or Einasto profiles, the current data exclude not only DM scenarios explaining the FERMI+HESS+PAMELA allowed regions, but also PAMELA regions alone to high confidence level
- For "cored" profiles, regions of the parameters space seem to be reopened (The annihilation into muons is not excluded yet)

## Leptonic Annihilation modes:

- For the NFW or Einasto profiles, the current data exclude not only DM scenarios explaining the FERMI+HESS+PAMELA allowed regions, but also PAMELA regions alone to high confidence level
- For "cored" profiles, regions of the parameters space seem to be reopened (The annihilation into muons is not excluded yet)

## Leptonic Decay modes:

- The residual isotropic radiation measured by Fermi imposes the strongest constraint and it is independent on the DM halo profiles
- It excludes the decay explanation of the FERMI+HESS+PAMELA anomalies for the  $\tau\tau$  channel and starts to exclude the decay explanation for the  $\mu\mu$  channel

## Leptonic Annihilation modes:

- For the NFW or Einasto profiles, the current data exclude not only DM scenarios explaining the FERMI+HESS+PAMELA allowed regions, but also PAMELA regions alone to high confidence level
- For "cored" profiles, regions of the parameters space seem to be reopened (The annihilation into muons is not excluded yet)

## Leptonic Decay modes:

- The residual isotropic radiation measured by Fermi imposes the strongest constraint and it is independent on the DM halo profiles
- It excludes the decay explanation of the FERMI+HESS+PAMELA anomalies for the  $\tau\tau$  channel and starts to exclude the decay explanation for the  $\mu\mu$  channel

## Tensions with other Constraints:

- Constraints from Synchrotron radiation (Bertone et al. arXiv:0811.3744)
- Constraints from Ionization and Heating of the InterGalactic Medium (Cirelli, Iocco, Panci arXiv:0907.0719), (Huetsi et al. arXiv:0906.4550), (Galli et al. arXiv:0905.0003)