



Contribution ID: 76

Type: Poster

## The Scaling of Electron Heating in Low-beta Reconnection Exhausts with Kinetic Reconnection Simulations

*Wednesday, 26 May 2021 13:00 (40 minutes)*

Particle heating in reconnection is essential to understand the heating in the solar corona, solar flares and the magnetotail. It plays an important role distributing magnetic energy into different species and between thermal and nonthermal components. Previous observational and theoretical studies on electron heating in reconnection exhausts within the  $\beta \sim 1$  regime suggest a simple linear scaling where the electron heating is proportional to the magnetic energy per particle. Using kinetic reconnection simulations in the low-beta regime (with  $\beta$  down to 0.005), we demonstrate that electron heating follows a sub-linear scaling below  $\beta \sim 0.01$ , with or without guide fields. As a result, the maximum heating is limited to only  $\sim 5$  times of upstream electron temperature. This electron heating scaling may be testable by MMS observations at the magnetotail. This new finding has strong implications for the efficiency of electron heating in reconnection at low-beta environments throughout heliosphysics.

**Primary authors:** ZHANG, Qile (Los Alamos National Lab); GUO, Fan (Los Alamos National Lab); Dr ØIEROSET, Marit (University of California Berkeley); Dr PHAN, Tai (University of California Berkeley); SHAY, Michael (University of Delaware); Dr DRAKE, James (University of Maryland College Park); Dr SWISDAK, Marc (University of Maryland College Park)

**Presenter:** ZHANG, Qile (Los Alamos National Lab)

**Session Classification:** Poster session: SQ3 and SQ6

**Track Classification:** Flare and Nanoflare heating