



Contribution ID: 42

Type: **Poster**

## Multi-instrument STIX microflare study

*Wednesday, 26 May 2021 13:00 (40 minutes)*

Solar flares are generally thought to be the impulsive release of magnetic energy giving rise to a wide range of solar phenomena that influence the heliosphere and in some cases even conditions of earth. Part of this liberated energy is used for particle acceleration and to heat up the solar plasma. The Spectrometer/Telescope for Imaging X-rays (STIX) instrument onboard the Solar Orbiter mission launched on February 10th 2020 promises advances in the study of solar flares of various sizes. It is capable of measuring X-ray spectra from 4 to 150 keV with 1 keV resolution binned into 32 energy bins before downlinking. With this energy range and sensitivity, STIX is capable of sampling thermal plasma with temperatures of  $\sim 10$  MK, and to diagnose the nonthermal bremsstrahlung emission of flare accelerated electrons. During the spacecraft commissioning phase in the first half of the year 2020, STIX observed 69 microflares. Of this set, 26 events could clearly be identified in at least two energy channels, all of which originated in an active region that was also visible from earth. These events provided a great opportunity to combine the STIX observations with the multi-band EUV imagery from the Atmospheric Imaging Assembly (AIA) instrument on board the earth orbiting Solar Dynamics Observatory (SDO). For two of the largest events (GOES class B6 and B2), it was possible to perform a spectroscopic analysis and to fit the spectra assuming thermal and nonthermal sources. These results are combined with plasma diagnostics derived from AIA EUV images. To this aim, the Differential Emission Measure (DEM) was reconstructed from AIA observations to infer maps of plasma temperatures and EM in the flaring regions. This allows us to supplement the information gained with STIX with high spatial resolution diagnostics of thermal plasma from the DEM reconstructed from AIA data.

**Primary authors:** SAQRI, Jonas (Institute of Physics, University of Graz); Prof. VERONIG, Astrid M. (Institute of Physics, Institute for Geophysics, Astrophysics and Meteorology (IGAM), University of Graz, Graz, Austria); Dr DICKSON, Ewan C. M. (Institute of Physics, Institute for Geophysics, Astrophysics and Meteorology (IGAM), University of Graz, Graz, Austria; University of Applied Sciences and Arts Northwestern Switzerland, Windisch, Switzerland); Dr WARMUTH, Alexander (Leibniz-Institut für Astrophysik Potsdam (AIP), Potsdam, Germany); Mr BATTAGLIA, Andrea Francesco (University of Applied Sciences and Arts Northwestern Switzerland, Windisch, Switzerland; ETH Zürich, Zürich, Switzerland); Dr BATTAGLIA, Marina (University of Applied Sciences and Arts Northwestern Switzerland, Windisch, Switzerland); Dr XIAO, Hualin (University of Applied Sciences and Arts Northwestern Switzerland, Windisch, Switzerland); Dr HURFORD, Gordon J. (University of Applied Sciences and Arts Northwestern Switzerland, Windisch, Switzerland); Prof. KRUCKER, Säm (University of Applied Sciences and Arts Northwestern Switzerland, Windisch, Switzerland; Space Sciences Laboratory, University of California, Berkeley, USA); THE STIX TEAM

**Presenter:** SAQRI, Jonas (Institute of Physics, University of Graz)

**Session Classification:** Poster session: SQ3 and SQ6

**Track Classification:** Flare and Nanoflare heating