


Self gravity affects quantum states

David Edward Bruschi

Institute for Quantum Computing Analytics (PGI-12)
Forschungszentrum Jülich
Germany

VIII.IX.MMXXI



Einstein gravity

We know that **ALL ENERGY GRAVITATES**.

Einstein gravity

We know that **ALL ENERGY GRAVITATES**.

Einstein equations

$$G_{\mu\nu} = \kappa T_{\mu\nu}.$$

These equations have been **highly successful** in providing many predictions.

Successes

- Precession of orbits;
- Bending of light;
- Black Holes;
- Penrose process;
- Gravitational waves;
- Cosmology.

Difficulties

- Rotation curves of galaxies;
- Nonlinearity of equations;
- Gravitation of quantum objects;
- Quantum nature of gravity;
- Fundamental or emergent theory?

Einstein gravity extended

We know that **ALL ENERGY GRAVITATES**.

Semiclassical gravity

$$G_{\mu\nu} = \kappa \langle : \hat{T}_{\mu\nu} : \rangle$$

$\hat{T}_{\mu\nu}$: **stress-energy tensor** for quantum field. $: \cdot :$ is **normal ordering**.

Successes

- Takes into account (somehow) backreaction;
- ...

Problems

- Fluctuations of stress energy tensor big/huge;
- Curved spacetime: inconsistent renormalisation procedures;
- “Strange” predictions for gravitational fields of superpositions;
- ...

Experiments at the overlap of relativity/quantum phys.

Planning

There are plans to try to test the gravitational field of small quantum objects that can be found in quantum states.

Experiments at the overlap of relativity/quantum phys.

Planning

There are plans to try to test the gravitational field of small quantum objects that can be found in quantum states.

Experiments of interest

- Spontaneous WF collapse;
- Gravitational decoherence;
- Superposition of masses;
- Optomechanical systems;
- Space based tests;
- Atom Interferometry;
- More?

One setup




Figure: Micius satellite CAS

Quantum Information (QI) and Thermodynamics (QT)

Quantum Information

Allows us to connect concepts such as entropy and quantum correlations.

Quantum Thermodynamics

QT investigates thermodynamics **far** from **thermodynamic limit**.

Regime of interest: where **fluctuations around the mean** are important;

Quantum Information (QI) and Thermodynamics (QT)

Quantum Information

Allows us to connect concepts such as entropy and quantum correlations.

Quantum Thermodynamics

QT investigates thermodynamics **far** from **thermodynamic limit**.

Regime of interest: where **fluctuations around the mean** are important;

Features

- Small (quantum) constituents;
- Few (e.g **ONE**) systems;
- Concepts of energy and work not unique;
- Fluctuation relations;
- ...

Quantum Information (QI) and Thermodynamics (QT)

Quantum Information

Allows us to connect concepts such as entropy and quantum correlations.

Quantum Thermodynamics

QT investigates thermodynamics **far** from **thermodynamic limit**.

Regime of interest: where **fluctuations around the mean** are important;

Features

- Small (quantum) constituents;
- Few (e.g **ONE**) systems;
- Concepts of energy and work not unique;
- Fluctuation relations;
- ...

Applications

- Quantum chemistry;
- Quantum refrigerators;
- Fundamental physics;
- Connections to Information Theory;
- ...

Quantum Thermodynamics (QT)

Resources


State $\hat{\rho}$. Unitaries \hat{U} . Then, exists \hat{U}_p : $\hat{\rho}_p = \hat{U}_p^\dagger \hat{\rho} \hat{U}_p$. And, $\hat{\rho}_p$ is “passive”.

Quantum Thermodynamics (QT)

Resources

State $\hat{\rho}$. Unitaries \hat{U} . Then, exists \hat{U}_p : $\hat{\rho}_p = \hat{U}_p^\dagger \hat{\rho} \hat{U}_p$. And, $\hat{\rho}_p$ is “passive”.

Features




Quantum Thermodynamics (QT)

Resources

State $\hat{\rho}$. Unitaries \hat{U} . Then, exists \hat{U}_p : $\hat{\rho}_p = \hat{U}_p^\dagger \hat{\rho} \hat{U}_p$. And, $\hat{\rho}_p$ is “passive”.

Features



Applications (PRE 91, 032118 (2015))


- Correlations \mathcal{I}_{AB} give work (W) \Leftrightarrow work (W) gives correlations \mathcal{I}_{AB} .

Quantum Thermodynamics (QT)

Resources

State $\hat{\rho}$. Unitaries \hat{U} . Then, exists \hat{U}_p : $\hat{\rho}_p = \hat{U}_p^\dagger \hat{\rho} \hat{U}_p$. And, $\hat{\rho}_p$ is “passive”.

Features



Applications (PRE 91, 032118 (2015))

- Correlations \mathcal{I}_{AB} give work (W) \Leftrightarrow work (W) gives correlations \mathcal{I}_{AB} .
- Bound on correlations (work):


$$\mathcal{I}_{AB} \leq \beta W.$$

Quantum Thermodynamics (QT)

Resources

State $\hat{\rho}$. Unitaries \hat{U} . Then, exists \hat{U}_p : $\hat{\rho}_p = \hat{U}_p^\dagger \hat{\rho} \hat{U}_p$. And, $\hat{\rho}_p$ is “passive”.

Features



Applications (PRE 91, 032118 (2015))

- Correlations \mathcal{I}_{AB} give work (W) \Leftrightarrow work (W) gives correlations \mathcal{I}_{AB} .
- Bound on correlations (work):

$$\mathcal{I}_{AB} \leq \beta W.$$


- Conclusion: Correlations must “carry energy”.

Role of correlations

On the weight of entanglement (Physics Letters B 54, 182-186 (2016))

Employing semiclassical gravity: “entanglement has a weight”.

We find $G_{\beta}^{(1)} \propto \beta$, where $G^{(1)}$ is correction to flat Einstein tensor.



A gedankenexperiment

The weight of a passive state


$$\hat{\rho} \xrightarrow{U} \hat{\rho}_p, \quad \text{Tr}(\hat{H}_0 \hat{\rho}_p) = E_0.$$

A gedankenexperiment

The weight of a passive state

$$\hat{\rho} \xrightarrow{U} \hat{\rho}_p, \quad \text{Tr}(\hat{H}_0 \hat{\rho}_p) = E_0.$$

Work from passive states




A gedankenexperiment

The weight of a passive state

$$\hat{\rho} \xrightarrow{U} \hat{\rho}_p, \quad \text{Tr}(\hat{H}_0 \hat{\rho}_p) = E_0.$$

Work from passive states



What happens


- Two passive states;

A gedankenexperiment

The weight of a passive state

$$\hat{\rho} \xrightarrow{U} \hat{\rho}_p, \quad \text{Tr}(\hat{H}_0 \hat{\rho}_p) = E_0.$$

Work from passive states



What happens


- Two passive states;
- (Also if GR dof in passive state);

A gedankenexperiment

The weight of a passive state

$$\hat{\rho} \xrightarrow{U} \hat{\rho}_p, \quad \text{Tr}(\hat{H}_0 \hat{\rho}_p) = E_0.$$

Work from passive states



What happens


- Two passive states;
- (Also if GR dof in passive state);
- Same state (e.g., temperature);

A gedankenexperiment

The weight of a passive state

$$\hat{\rho} \xrightarrow{U} \hat{\rho}_p, \quad \text{Tr}(\hat{H}_0 \hat{\rho}_p) = E_0.$$

Work from passive states



What happens


- Two passive states;
- (Also if GR dof in passive state);
- Same state (e.g., temperature);
- Attract each other;

A gedankenexperiment

The weight of a passive state

$$\hat{\rho} \xrightarrow{U} \hat{\rho}_p, \quad \text{Tr}(\hat{H}_0 \hat{\rho}_p) = E_0.$$

Work from passive states



What happens


- Two passive states;
- (Also if GR dof in passive state);
- Same state (e.g., temperature);
- Attract each other;
- Cannot extract energy from state with unitaries on both;

A gedankenexperiment

The weight of a passive state

$$\hat{\rho} \xrightarrow{U} \hat{\rho}_p, \quad \text{Tr}(\hat{H}_0 \hat{\rho}_p) = E_0.$$

Work from passive states



What happens


- Two passive states;
- (Also if GR dof in passive state);
- Same state (e.g., temperature);
- Attract each other;
- Cannot extract energy from state with unitaries on both;
- System interacts gravitationally;

A gedankenexperiment

The weight of a passive state

$$\hat{\rho} \xrightarrow{U} \hat{\rho}_p, \quad \text{Tr}(\hat{H}_0 \hat{\rho}_p) = E_0.$$

Work from passive states



What happens


- Two passive states;
- (Also if GR dof in passive state);
- Same state (e.g., temperature);
- Attract each other;
- Cannot extract energy from state with unitaries on both;
- System interacts gravitationally;
- grav. inter.=employable energy;

A gedankenexperiment

The weight of a passive state

$$\hat{\rho} \xrightarrow{U} \hat{\rho}_p, \quad \text{Tr}(\hat{H}_0 \hat{\rho}_p) = E_0.$$

Work from passive states



What happens


- Two passive states;
- (Also if GR dof in passive state);
- Same state (e.g., temperature);
- Attract each other;
- Cannot extract energy from state with unitaries on both;
- System interacts gravitationally;
- grav. inter.=employable energy;
- Employable energy= work;

A gedankenexperiment

The weight of a passive state

$$\hat{\rho} \xrightarrow{U} \hat{\rho}_p, \quad \text{Tr}(\hat{H}_0 \hat{\rho}_p) = E_0.$$

Work from passive states



What happens

- Two passive states;
- (Also if GR dof in passive state);
- Same state (e.g., temperature);
- Attract each other;
- Cannot extract energy from state with unitaries on both;
- System interacts gravitationally;
- grav. inter.=employable energy;
- Employable energy= work;
- Something is wrong.

A novel proposal (arXiv:1701.00699)

A novel proposal (arXiv:1701.00699)

Solution

Only **extractible energy** gravitates.

A novel proposal (arXiv:1701.00699)

Solution

Only **extractible energy** gravitates.

We suggest that only extractible work is the source of gravity. In this sense

A novel proposal (arXiv:1701.00699)

Solution

Only **extractible energy** gravitates.

We suggest that only extractible work is the source of gravity. In this sense

New proposal

The field equations of Nature need to take into account the fact that it is extractible work that determines gravity, not the “whole amount of energy”.

However: this might require to change other fundamental equations (i.e., Heisenberg eq.).

Features of the proposal I

Clarification

Passive state $\hat{\rho}_p$ is vacuum state of **full theory** (including gravity part).

Immediate consequences significantly different from standard GR/QFTCS?

Features of the proposal I

Clarification

Passive state $\hat{\rho}_p$ is vacuum state of **full theory** (including gravity part).

Immediate consequences significantly different from standard GR/QFTCS?

Prediction

- IF: Universe initially filled with thermal radiation;
- Thermal state is passive;
- $G_{\mu\nu} \equiv 0$. **No gravity** in such Universe.

Features of the proposal I

Clarification

Passive state $\hat{\rho}_p$ is vacuum state of **full theory** (including gravity part).

Immediate consequences significantly different from standard GR/QFTCS?

Prediction

- IF: Universe initially filled with thermal radiation;
- Thermal state is passive;
- $G_{\mu\nu} \equiv 0$. **No gravity** in such Universe.

Prediction

- Initial static (Schwarzschild) black hole;
- Choose: any initial vacuum;
- $G_{\mu\nu} \equiv 0 \Rightarrow$ **NO Hawking radiation**.
- (Same for Unruh effect).

Features of the proposal I

Clarification

Passive state $\hat{\rho}_p$ is vacuum state of **full theory** (including gravity part).

Immediate consequences significantly different from standard GR/QFTCS?

Prediction

- IF: Universe initially filled with thermal radiation;
- Thermal state is passive;
- $G_{\mu\nu} \equiv 0$. **No gravity** in such Universe.

Prediction

- Initial static (Schwarzschild) black hole;
- Choose: any initial vacuum;
- $G_{\mu\nu} \equiv 0 \Rightarrow$ **NO Hawking radiation**.
- (Same for Unruh effect).

N.B.

Classical states: little/vanishing zero point energy compared to total one.

Features of the proposal II

Considerations on known effects in QFT

- Unruh effect: theoretically requires infinite acceleration times (i.e., infinite fuel).
- Schwarzschild BH evaporation: evaporates forever as seen by somebody sitting at infinity (i.e., infinite energy).

Features of the proposal II

Considerations on known effects in QFT

- Unruh effect: theoretically requires infinite acceleration times (i.e., infinite fuel).
- Schwarzschild BH evaporation: evaporates forever as seen by somebody sitting at infinity (i.e., infinite energy).

Considerations on known effects in QFT

- Both cases: based on existence of different, non-equivalent vacua.
- Non-trivial Bogoliubov transf. between vacua (entangled states).

Features of the proposal II

Considerations on known effects in QFT

- Unruh effect: theoretically requires infinite acceleration times (i.e., infinite fuel).
- Schwarzschild BH evaporation: evaporates forever as seen by somebody sitting at infinity (i.e., infinite energy).

Considerations on known effects in QFT

- Both cases: based on existence of different, non-equivalent vacua.
- Non-trivial Bogoliubov transf. between vacua (entangled states).

The proposal...

... predicts that these highly ideal scenarios should be “trivial” because the unitary transformation involved (“change of observer”) does **not** add energy to the observed system (which is in the vacuum state).

Time evolution by work (AoP 394, 155-161 (2018))

Time evolution

- Time evolution is typically “driven” by Hamiltonian.
- The theory is based on the fact that not all energy gravitates.
- \Rightarrow time evolution cannot be driven by Hamiltonian

Time evolution by work (AoP 394, 155-161 (2018))

Time evolution

- Time evolution is typically “driven” by Hamiltonian.
- The theory is based on the fact that not all energy gravitates.
- \Rightarrow time evolution cannot be driven by Hamiltonian

The modified Heisenberg equation

$$\dot{A} = \frac{i}{\hbar} [H, A] - \frac{i}{\hbar} [U_p^\dagger H U_p, A] + \frac{\partial A}{\partial t}.$$

Time evolution by work (AoP 394, 155-161 (2018))

Time evolution

- Time evolution is typically “driven” by Hamiltonian.
- The theory is based on the fact that not all energy gravitates.
- \Rightarrow time evolution cannot be driven by Hamiltonian

The modified Heisenberg equation

$$\dot{A} = \frac{i}{\hbar}[H, A] - \frac{i}{\hbar}[U_p^\dagger H U_p, A] + \frac{\partial A}{\partial t}.$$

Heisenberg evolution

- $|\psi\rangle = \cos \theta |0\rangle + \sin \theta |1\rangle;$
- $\dot{\rho} = \frac{i}{\hbar}[\rho, A];$
- $p_{|0\rangle} = \cos^2 \theta$
 $p_{|1\rangle} = \sin^2 \theta.$

Time evolution by work (AoP 394, 155-161 (2018))

Time evolution

- Time evolution is typically “driven” by Hamiltonian.
- The theory is based on the fact that not all energy gravitates.
- \Rightarrow time evolution cannot be driven by Hamiltonian

The modified Heisenberg equation

$$\dot{A} = \frac{i}{\hbar}[H, A] - \frac{i}{\hbar}[U_p^\dagger H U_p, A] + \frac{\partial A}{\partial t}.$$

Heisenberg evolution

- $|\psi\rangle = \cos \theta |0\rangle + \sin \theta |1\rangle;$
- $\dot{\rho} = \frac{i}{\hbar}[\rho, A];$
- $p_{|0\rangle} = \cos^2 \theta$
 $p_{|1\rangle} = \sin^2 \theta.$

“Novel” time evolution

- $|\psi\rangle = \cos \theta |0\rangle + \sin \theta |1\rangle;$
- $\dot{\rho} = \frac{i}{\hbar}[H, \rho] - \frac{i}{\hbar}[U_p^\dagger H U_p, \rho];$
- $p_{|0\rangle} = \cos^2 \theta - \cos^2 \theta \cos^2(\sin \theta \frac{E_1}{\hbar} t)$
 $p_{|1\rangle} = \sin^2 \theta + \cos^2 \theta \cos^2(\sin \theta \frac{E_1}{\hbar} t).$

Testing in experiments (AoP 394, 155-161 (2018))

An experimental proposal

- Use highly entangled quantum state;
- Use Mach-Zender interferometer with arms at **different** heights;
- Use new time evolution equation.

Testing in experiments (AoP 394, 155-161 (2018))

An experimental proposal

- Use highly entangled quantum state;
- Use Mach-Zender interferometer with arms at **different** heights;
- Use new time evolution equation.

Scheme




Figure: CQG 29, 224010 (2012)

Testing in experiments (AoP 394, 155-161 (2018))

An experimental proposal

- Use highly entangled quantum state;
- Use Mach-Zender interferometer with arms at **different** heights;
- Use new time evolution equation.

Scheme




Figure: CQG 29, 224010 (2012)

Heisenberg evolution

- $|\psi\rangle = \frac{1}{\sqrt{2}} [|N0\rangle + |0N\rangle];$
- $\dot{\rho} = \frac{i}{\hbar} [H, \rho] - \frac{i}{\hbar} [U_p^\dagger H U_p, \rho];$
- $p_{|N0\rangle} = \frac{1}{2} [1 + \frac{r_s}{r_E} \frac{h}{r_E} \sin^2(\frac{N\omega_0 t}{2})];$
 $p_{|0N\rangle} = \frac{1}{2} [1 - \frac{r_s}{r_E} \frac{h}{r_E} \sin^2(\frac{N\omega_0 t}{2})];$
- Note: $\frac{r_s}{r_E} \frac{h}{r_E} \sim \frac{g h}{c^2};$
- Note: $\frac{r_s}{r_E} \frac{h}{r_E} \sim h \times 10^{-16}.$

Conclusions

We have:

- Studied role of quantum correlations/entanglement in gravitating quantum systems;
- Proposed that only extractible energy gravitates;
- Modified time evolution to be compatible with new proposal

Conclusions

We have:

- Studied role of quantum correlations/entanglement in gravitating quantum systems;
- Proposed that only extractible energy gravitates;
- Modified time evolution to be compatible with new proposal

Then:

- Applications include localized scenarios, as well as global ones;
- Predictions are (very) different from the standard known theories;
- More advanced mathematical tools can be used to propose better models;
- Experiments can be potentially done with current technology;
- More work to come...

Acknowledgments

Grazie

PLB 54, 182-186 (2016) — arXiv:1701.00699 — AoP 394, 155-161 (2018)

