

LIME background simulation

17/05/2021 CYGNO Simulation meeting

F. Di Giambattista

Background rates

PRELIMINARY

- **40cm water + 10cm copper**

External gammas ($0.56 \text{ cm}^{-2} \text{ s}^{-1}$)

(ER): $(2.05 \pm 0.10) \times 10^5$ events/yr (total)

(ER): $(4.0 \pm 0.4) \times 10^4$ events/yr (0-20 keV)

External neutrons ($2.7 \times 10^{-6} \text{ cm}^{-2} \text{ s}^{-1}$)

(ER+NR): (23.6 ± 0.6) events/yr (total)

(ER+NR): (9.9 ± 0.3) events/yr (0-20 keV)

(Only NR): (4.7 ± 0.1) events/yr (total)

(Only NR): (3.2 ± 0.1) events/yr (0-20 keV)

With Cu: 4.0×10^4 ER/yr (0-20keV)

With Pb: 4.5×10^4 ER/yr (0-20keV)

Internal background:

$0.5\text{-}1.0 \times 10^5$ ER/yr (0-20keV)

- **40cm water + 5cm lead + 5cm copper**

External gammas ($0.56 \text{ cm}^{-2} \text{ s}^{-1}$)

(ER): $(1.8 \pm 0.5) \times 10^5$ events/yr (total)

(ER): $(3.9 \pm 2.2) \times 10^4$ events/yr (0-20 keV)

External neutrons ($2.7 \times 10^{-6} \text{ cm}^{-2} \text{ s}^{-1}$)

(ER+NR): (30.1 ± 0.3) events/yr (total)

(ER+NR): (12.1 ± 0.2) events/yr (0-20 keV)

(Only NR): (4.38 ± 0.09) events/yr (total)

(Only NR): (2.66 ± 0.07) events/yr (0-20 keV)

Radioactivity from $^{210}\text{Bi}^*$:



(ER): $(2.47 \pm 0.05) \times 10^4$ events/yr (total)

(ER): $(6.2 \pm 0.3) \times 10^3$ events/yr (0-20 keV)

* $(80\text{Bq/kg} \times 5658.7\text{kg of lead} = 452693 \text{ Bq})$



Spectra – 40cm water + 10cm Cu

PRELIMINARY



Spectra – 40cm water + 5cm Cu + 5cm Pb

PRELIMINARY



Conclusions

- There might be problems of self-consistency in the simulation – to be checked
- Still need to simulate the radioactivity of **resistors of the field cage**
- *If these results are correct*, difference in background rate between copper and lead configurations is small – using lead might be a good choice
 - Not necessarily true for both LIME and CYGNO: the volume of lead in CYGNO is much higher for same thickness