



Contribution ID: 361

Type: Poster

LCDAs of heavy hadrons and their first inverse moments

Friday, 8 July 2022 20:10 (20 minutes)

Heavy-quark symmetry (HQS), despite being approximate, allows to relate dynamically many hadron systems. In the HQS-limit heavy mesons and doubly-heavy baryons are very similar as their dynamics is determined by a light quark moving in a color field of a static source. As in the meson case, matrix elements of non-local interpolation currents between the baryon state and vacuum are determined by light-cone distribution amplitudes (LCDAs). The first inverse moment of the leading twist B -meson distribution amplitude is a hadronic parameter needed for an accurate theoretical description of B -meson exclusive decays. It is quite natural that a similar moment of doubly-heavy baryon is of importance in exclusive doubly-heavy baryons' decays. We obtain HQET sum rules for the first inverse moment based on the correlation functions containing nonlocal heavy-light operator of the doubly-heavy baryon and its local interpolating current. Numerical estimates of this moment are presented.

In-person participation

No

Primary author: SHUKHTINA, Alisa**Co-author:** PARKHOMENKO, Alexander (P.G. Demidov Yaroslavl State University)**Presenter:** SHUKHTINA, Alisa**Session Classification:** Poster Session**Track Classification:** Strong interactions and Hadron Physics