



Contribution ID: 816

Type: Poster

The boosted $X \rightarrow b\bar{b}$ tagger calibration using $Z \rightarrow b\bar{b}$ events collected with the ATLAS detector

Friday, 8 July 2022 20:10 (20 minutes)

Many analyses in the ATLAS physics program are dependent on the identification of jets containing b-hadrons (b-tagging). The corresponding algorithms are referred to as b-taggers. The baseline b-taggers are optimized for jets containing one b-hadron. A new double b-tagging algorithm, the $X \rightarrow b\bar{b}$ tagger, provides better identification efficiency to reconstruct boosted resonant particles decaying into a pair of b-quarks. In the boosted regime, it is a challenging task because of high collimation of the two b-hadrons. This neural network based $X \rightarrow b\bar{b}$ tagger uses the kinematic information of the large radius ($R=1.0$) jet and the flavour information of associated track-jets. The performance of this tagger was evaluated using Monte Carlo simulation, therefore it could vary in collision data. Thus this poster presents the in situ tagging efficiency calibration using $Z \rightarrow b\bar{b}$ events with a recoiling photon or jet for this boosted $X \rightarrow b\bar{b}$ tagger. The efficiency data to simulation scale factor is derived using the Run 2 pp collision data collected by ATLAS experiment at $\sqrt{s} = 13$ TeV, with the integrated luminosity of 139 fb^{-1} .

In-person participation

No

Primary author: BATTULGA, Daariimaa (Humboldt University of Berlin (DE))**Presenter:** BATTULGA, Daariimaa (Humboldt University of Berlin (DE))**Session Classification:** Poster Session**Track Classification:** Operation, Performance and Upgrade (Incl. HL-LHC) of Present Detectors