



Contribution ID: 816

Type: Poster

## The boosted $X \rightarrow b\bar{b}$ tagger calibration using $Z \rightarrow b\bar{b}$ events collected with the ATLAS detector

*Friday, 8 July 2022 20:10 (20 minutes)*

Many analyses in the ATLAS physics program are dependent on the identification of jets containing b-hadrons (b-tagging). The corresponding algorithms are referred to as b-taggers. The baseline b-taggers are optimized for jets containing one b-hadron. A new double b-tagging algorithm, the  $X \rightarrow b\bar{b}$  tagger, provides better identification efficiency to reconstruct boosted resonant particles decaying into a pair of b-quarks. In the boosted regime, it is a challenging task because of high collimation of the two b-hadrons. This neural network based  $X \rightarrow b\bar{b}$  tagger uses the kinematic information of the large radius ( $R=1.0$ ) jet and the flavour information of associated track-jets. The performance of this tagger was evaluated using Monte Carlo simulation, therefore it could vary in collision data. Thus this poster presents the in situ tagging efficiency calibration using  $Z \rightarrow b\bar{b}$  events with a recoiling photon or jet for this boosted  $X \rightarrow b\bar{b}$  tagger. The efficiency data to simulation scale factor is derived using the Run 2 pp collision data collected by ATLAS experiment at  $\sqrt{s} = 13$  TeV, with the integrated luminosity of  $139 \text{ fb}^{-1}$ .

### In-person participation

No

**Primary author:** BATTULGA, Daariimaa (Humboldt University of Berlin (DE))**Presenter:** BATTULGA, Daariimaa (Humboldt University of Berlin (DE))**Session Classification:** Poster Session**Track Classification:** Operation, Performance and Upgrade (Incl. HL-LHC) of Present Detectors