



Contribution ID: 76

Type: **Parallel Talk**

$\psi(2S)$ production and nuclear modification factor in nucleus-nucleus collisions with ALICE

Saturday, 9 July 2022 09:35 (15 minutes)

Charmonium production is a probe sensitive to deconfinement in nucleus-nucleus collisions. The production of J/ψ via regeneration within the QGP or at the phase boundary has been identified as an important ingredient for the description of the observed centrality and p_T dependence at the LHC. $\psi(2S)$ production relative to J/ψ is one possible discriminator between the two different regeneration scenarios. At RHIC and at the LHC, there is so far no significant observation of the $\psi(2S)$ in nucleus-nucleus collisions in central events at low transverse momentum, where regeneration is the dominating process. The combined Run 2 data set of ALICE allows to extract a significant $\psi(2S)$ signal in such a kinematic region at forward rapidity in the dimuon decay channel. In this contribution, we present for the first time results on the $\psi(2S)$ -to- J/ψ double ratio and the $\psi(2S)$ nuclear modification factor in Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, calculated with respect to a new pp reference with improved precision. Results are compared with model calculations.

In-person participation

Yes

Primary author: CC CHAIRS, ALICE**Presenter:** PAUL, Biswarup (Istituto Nazionale di Fisica Nucleare)**Session Classification:** Heavy Ions**Track Classification:** Heavy Ions