

Searching for jet quenching effect using high multiplicity inclusive and semi-inclusive jets in pp collisions with ALICE




Kotliarov Artem

for the ALICE Collaboration

Nuclear Physics Institute of the Czech Academy of Sciences



Signatures of QGP-like formation in small-collision systems



How do QGP signatures evolve with the system size?

1. Enhancement of strange hadron production in high-multiplicity pp collisions
2. Pronounced ridge structures in high-multiplicity pp collisions
3. Non-zero v_2 coefficient for low and high- p_T particles in p-Pb collisions

Do we observe signs of jet quenching in small systems?

Measurement of nuclear modification factor in p-Pb collisions




Yield suppression of high- p_T inclusive jets/hadrons relative to minimum bias pp

$$R_{AA} = \frac{d^2 N_{AA}/dydp_T}{\langle T_{AA} \rangle d^2 \sigma_{pp}^{\text{INEL}}/dydp_T}$$

- Limited precision of $\langle T_{AA} \rangle$ for centrality biased events
- Undefined Glauber scaling for high-multiplicity pp collisions

No conclusive results on jet quenching in small systems

→ more sensitive approaches are needed




Semi-inclusive measurements of hadron-jet acoplanarity

$$\frac{1}{N_{\text{trig}}^{\text{AA}}} \frac{d^2 N_{\text{jet}}^{\text{AA}}}{dp_{\text{T,jet}}^{\text{ch}} d|\Delta\varphi|} \bigg|_{p_{\text{T,trig}} \in \text{TT}} = \left(\frac{1}{\sigma^{\text{pp} \rightarrow \text{h} + \text{X}}} \frac{d^2 \sigma^{\text{pp} \rightarrow \text{h} + \text{jet} + \text{X}}}{dp_{\text{T,jet}}^{\text{ch}} d|\Delta\varphi|} \right) \bigg|_{p_{\text{T,trig}} \in \text{TT}} \times \frac{\langle T_{\text{AA}} \rangle}{\langle T_{\text{AA}} \rangle}$$

Cross section for trigger hadron production

Differential cross section for coincidence production of trigger hadron and recoil jet



- Applicable in pp collisions
- Equality in case of no nuclear effects
- **Self-normalized observable, reference spectrum has no dependence on $\langle T_{\text{AA}} \rangle$**

Hadron-jet acoplanarity measurements in Pb-Pb collisions → talk by [Yongzhen Hou](#) on 8 July at 3:05 pm, “Heavy Ions” session

Event activity selection in pp collisions at $\sqrt{s} = 13$ TeV

Online data triggers based on V0 detectors:



- Minimum-bias (MB) trigger $\rightarrow L_{\text{Int}} \approx 32 \text{ nb}^{-1}$
- High-multiplicity (HM) trigger $\rightarrow L_{\text{Int}} \approx 10^4 \text{ nb}^{-1}$

Offline event activity (EA) selection:

$$V0M = V0A + V0C \rightarrow \text{sum of signals}$$

Characterization of EA in terms of $V0M/\langle V0M \rangle$




$\langle V0M \rangle$ - mean of MB distribution



Inclusive jet yield in different multiplicity classes

NEW

VOM multiplicity classes




- Jet p_T corrected for underlying events and instrumental effects
- **Event activity bias has a mild effect on the shape** for $p_{T,jet} > 20$ GeV/c
- Jet yield increases with event activity bias

Jet production ratio using different jet R


NEW

Ratio of jet p_T spectra




- **Small R** \rightarrow weak dependence on event activity bias
- **Large R** \rightarrow hint of jet production ordering

Ratio of p_T differential cross sections



- **No dependence** on \sqrt{s} observed
- p-Pb ratio compatible with pp one


Self-normalized jet production vs self-normalized multiplicity



- **Non-linear rising with multiplicity** observed for jet production in midrapidity, similar trend as for J/ψ ¹
- **Electrons from W decay follow linear trend**, talk by [Mingrui Zhao](#) on 7 July at 5.30 pm, “Heavy Ions” session
- PYTHIA 8 **overestimates** data at high particle multiplicities

¹ ALICE, Phys. Lett. B 810 (2020)

Semi-inclusive p_T distribution of recoil jets



- Jet p_T corrected for underlying event density

$$p_{T,jet}^{ch} = p_{T,jet}^{raw,ch} - \rho A_{jet}$$


- Negative and low p_T region

→ **contribution of combinatorial background jets**


- Yield of combinatorial jets **has no dependence on p_T of TT**
- Data-driven approach for removal of uncorrelated jet yield

$$\Delta_{recoil} = \frac{1}{N_{trig}} \frac{d^2 N_{jets}}{dp_{T,jet}^{ch} d|\Delta\phi|} \Big|_{TT\{20,30\}} - \frac{1}{N_{trig}} \frac{d^2 N_{jets}}{dp_{T,jet}^{ch} d|\Delta\phi|} \Big|_{TT\{6,7\}}$$

TT{x,y} → trigger-track with $p_T \in (x, y)$ GeV/c




Semi-inclusive azimuthal distribution of recoil jets




Construction of Δ_{recoil} observable as a function of TT-jet opening angle for a given $p_{T,\text{jet}}$

$$\Delta_{\text{recoil}} = \frac{1}{N_{\text{trig}}} \frac{d^2 N_{\text{jets}}}{dp_{T,\text{jet}}^{\text{ch}} d|\Delta\phi|} \Big|_{\text{TT}\{20,30\}} - \frac{1}{N_{\text{trig}}} \frac{d^2 N_{\text{jets}}}{dp_{T,\text{jet}}^{\text{ch}} d|\Delta\phi|} \Big|_{\text{TT}\{6,7\}} \quad \text{TT}\{x,y\} \rightarrow \text{trigger-track with } p_T \in (x,y) \text{ GeV}/c$$



ALI-PREL-339864

Projection for specified
jet p_T region



ALI-PREL-339825


Measurements of hadron-jet acoplanarity with $\Delta_{\text{recoil}} (\Delta\phi)$



NEW

Fully corrected data:

- Substantial suppression of jets back-to-back w.r.t. T^*T in HM collisions
- Broadening of HM acoplanarity distribution with respect to MB
- Resembles jet quenching effects



Measurements of hadron-jet acoplanarity with $\Delta_{\text{recoil}} (\Delta\phi)$



NEW


Fully corrected data:

- Substantial suppression of jets back-to-back w.r.t. $T\bar{T}$ in HM collisions
- Broadening of HM acoplanarity distribution with respect to MB
- Resembles jet quenching effects

PYTHIA 8 Monash simulation:


- $V0M$ = # of charged, final state particles within V0A & V0C acceptances
- Does not account for jet quenching effects
- Exhibits qualitatively similar features as experimental data

What can we learn from PYTHIA simulation
about recoil jets in MB and HM events?




ALI-PREL-502404

PYTHIA 8 simulation: recoil jet pseudorapidity distribution






ALI-SIMUL-347689






* Grey boxes represent acceptances of V0 detectors

PYTHIA 8 simulation: recoil jet pseudorapidity distribution





- Most likely HM trigger is induced by high- p_T recoil jets
 \leftrightarrow HM trigger imposes substantial bias on recoil jets
 - Lower enhancement in V0A is caused by asymmetric coverage of V0 detectors
- * Grey boxes represent acceptances of V0 detectors

PYTHIA 8 simulation: recoil jet pseudorapidity distribution




- Most likely HM trigger is induced by high- p_T recoil jets
↔ HM trigger imposes substantial bias on recoil jets
 - Lower enhancement in V0A is caused by asymmetric coverage of V0 detectors
- * Grey boxes represent acceptances of V0 detectors

PYTHIA 8 simulation: recoil jet pseudorapidity distribution







ALI-SIMUL-347693






- Most likely HM trigger is induced by high- p_T recoil jets
 \leftrightarrow HM trigger imposes substantial bias on recoil jets
 - Lower enhancement in V0A is caused by asymmetric coverage of V0 detectors
- * Grey boxes represent acceptances of V0 detectors

PYTHIA 8 simulation: recoil jet pseudorapidity distribution



- Most likely HM trigger is induced by high- p_T recoil jets
↔ HM trigger imposes substantial bias on recoil jets
 - Lower enhancement in V0A is caused by asymmetric coverage of V0 detectors
- * Grey boxes represent acceptances of V0 detectors


PYTHIA 8 simulation: recoil jet pseudorapidity distribution



ALI-SIMUL-347689




ALI-SIMUL-347693

ALI-SIMUL-347697



- Most likely HM trigger is induced by high- p_T recoil jets
↔ HM trigger imposes substantial bias on recoil jets
 - Lower enhancement in V0A is caused by asymmetric coverage of V0 detectors
- * Grey boxes represent acceptances of V0 detectors


PYTHIA 8 simulation: recoil jet pseudorapidity distribution



ALI-SIMUL-347689


ALI-SIMUL-347693

ALI-SIMUL-347697




- Most likely HM trigger is induced by high- p_T recoil jets
↔ HM trigger imposes substantial bias on recoil jets
 - Lower enhancement in V0A is caused by asymmetric coverage of V0 detectors
- * Grey boxes represent acceptances of V0 detectors


PYTHIA 8 simulation: number of high- p_T recoil jets vs event activity



PYTHIA 8 simulation: number of high- p_T recoil jets vs event activity




ALI-PREL-502416




- Lower probability to observe 1 hard recoil jet in HM events
 → result in suppression of acoplanarity

PYTHIA 8 simulation: number of high- p_T recoil jets vs event activity



ALI-PREL-502416



- Lower probability to observe 1 hard recoil jet in HM events
→ **results in suppression of acoplanarity**
- HM trigger → **bias toward multi-jet final state**

No jet quenching effects observed in high-multiplicity pp collisions. **Potentially, signal is too small**

Inclusive jet measurements

- Jet production **rises** with event activity
- Event activity bias has **weak impact on the spectrum slope for high- p_T jets**

Semi-inclusive jet measurements

- Broadening and suppression of back-to-back hadron-jet correlation in HM events relative to MB
- PYTHIA quantitatively reproduces the shape → **jet quenching signal is not genuine**
 - HM trigger enhances probability to measure high- p_T recoil jets in V0 acceptance
 - Bias towards multi-jet final state induced by HM trigger → **obscures possible jet quenching signal**
 - Multi-jet final state → **generic bias for HM measurements in small systems**

Thank you for your attention!