



Contribution ID: 126

Type: Parallel Talk

Measurement of lepton-jet correlation in deep-inelastic scattering with the H1 detector using machine learning for unfolding

Saturday, 9 July 2022 10:00 (15 minutes)

The lepton-jet momentum imbalance in deep inelastic scattering events offers a useful set of observables for unifying collinear and transverse-momentum-dependent frameworks for describing high energy Quantum Chromodynamics interactions. A recent first measurement was made [1] of this imbalance in the laboratory frame using positron-proton collision data recorded with the H1 experiment at HERA in the years 2006-2007. Using a new machine learning method, the measurement was performed simultaneously and unbinned in eight dimensions. The first results were presented as a set of four one-dimensional projections onto key observables. This work extends over those results by making use of the multi-differential nature of the unfolded result. In particular, distributions of lepton-jet correlation observables are studied as a function of the kinematic properties of the scattering process, i.e. as a function of the momentum transfer $Q^2 > 150 \text{ GeV}^2$ and the inelasticity $0.2 < y < 0.7$.

H1prelim-22-031

[1] arxiv:2108.12376, accepted by PRL

In-person participation

No

Primary authors: SCHMITT, Stefan (DESY); XU, YAOYUAN (Lawrence Berkeley National Laboratory)**Presenter:** XU, YAOYUAN (Lawrence Berkeley National Laboratory)**Session Classification:** Strong interactions and Hadron Physics**Track Classification:** Strong interactions and Hadron Physics