



Contribution ID: 1160

Type: Parallel Talk

## Linear power corrections to $e^+e^-$ shape variables in the three-jet region

Thursday, July 7, 2022 6:35 PM (15 minutes)

A proper understanding of non-perturbative effects, which manifest themselves as linear power corrections, is needed to describe many observables measured at colliders. We report on recent progress in the calculation of linear power corrections to shape variables such as the  $C$ -parameter and thrust in the three-jet region arising from infrared renormalons. Previously, only the results at the two Sudakov shoulders, namely the two-jet and symmetric three-jet limits, have been known in the literature. We develop a formalism that allows us to compute power corrections in the entire three-jet region, and discuss its implications for the determination of the strong coupling constant  $\alpha_s$ . We derive a factorisation formula for the power corrections in which the so-called Milan factor naturally arises, and present analytic results for the power corrections for the  $C$ -parameter and the thrust in the generic  $N$ -jet region.

### In-person participation

Yes

**Primary authors:** CAOLA, Fabrizio (University of Oxford, Wadham College); LIMATOLA, Giovanni (Università degli Studi di Milano Bicocca and INFN); MELNIKOV, Kirill (KIT); OZCELIK, Melih Arslan (TTP-KIT); NASON, Paolo (MIB); FERRARIO RAVASIO, Silvia (University of Oxford)

**Presenter:** OZCELIK, Melih Arslan (TTP-KIT)

**Session Classification:** Strong interactions and Hadron Physics

**Track Classification:** Strong interactions and Hadron Physics