



Contribution ID: 1012

Type: Parallel Talk

New physics behind the new muon $g-2$ puzzle?

Friday, July 8, 2022 11:49 AM (17 minutes)

The recent measurement of the muon $g-2$ at Fermilab confirms the previous Brookhaven result. The leading hadronic vacuum polarization (HVP) contribution to the muon $g-2$ represents a crucial ingredient to establish if the Standard Model prediction differs from the experimental value. A recent lattice QCD result by the BMW collaboration shows a tension with the low-energy $e+e-\rightarrow$ hadrons data which are currently used to determine the HVP contribution. We refer to this tension as the new muon $g-2$ puzzle. In this Letter we consider the possibility that new physics contributes to the $e+e-\rightarrow$ hadrons cross-section. This scenario could, in principle, solve the new muon $g-2$ puzzle. However, we show that this solution is excluded by a number of experimental constraints.

In-person participation

Yes

Primary authors: MASIERO, Antonio (U. Padova); DI LUZIO, Luca (U. Padova); PARADISI, Paride (Istituto Nazionale di Fisica Nucleare); PASSERA, Massimo (INFN Padova)

Presenters: MASIERO, Antonio (U. Padova); DI LUZIO, Luca (U. Padova); PARADISI, Paride (Istituto Nazionale di Fisica Nucleare); PASSERA, Massimo (INFN Padova)

Session Classification: Quark and Lepton Flavour Physics

Track Classification: Quark and Lepton Flavour Physics