



Contribution ID: 393

Type: **Parallel Talk**

## Stau searches and measurements with the ILD concept at the International Linear Collider

*Friday, 8 July 2022 11:45 (15 minutes)*

The direct pair-production of the tau-lepton superpartner, stau, is one of the most interesting channels to search for SUSY. First of all the stau is with high probability the lightest of the scalar leptons. Secondly the signature of stau pair production signal events is one of the most difficult ones, yielding to the 'worst' and so most global scenario for the searches. The current model-independent stau limits comes from analysis performed at LEP but they suffer from the low energy of this facility. The LHC exclusion reach extends to higher masses for large mass differences, but under strong model assumptions.

The ILC, a future electron-positron collider with energy up to 1 TeV, is ideally suited for SUSY searches. The capability of the ILC for determining exclusion/discovery limits for the stau in a model-independent way is shown in this contribution, together with an overview of the current state-of-the-art. A detailed study of the 'worst' scenario for stau exclusion/discovery, taking into account the effect of the stau mixing on stau production cross-section and efficiency is presented. For selected benchmarks, the prospect for measuring masses and polarised cross-sections will be shown. The studies were done studying events passed through the full detector simulation and reconstruction procedures of the International Large Detector concept (ILD) at the ILC. The simulation included all SM backgrounds, as well as the machine induced ones.

### In-person participation

No

**Primary authors:** LIST, Jenny (DESY); KAWAGOE, Kiyotomo (Kyushu University); PARDO DE VERA, Maria Teresa (DESY); BERGGREN, Mikael (DESY)

**Presenter:** PARDO DE VERA, Maria Teresa (DESY)

**Session Classification:** Beyond the Standard Model

**Track Classification:** Beyond the Standard Model