



Contribution ID: 445

Type: **Parallel Talk**

Higgs boson decay to J/ψ via c -quark fragmentation

Friday, 8 July 2022 11:45 (15 minutes)

With the discovery of the Higgs boson at the CERN Large Hadron Collider (LHC), the particle spectrum of the Standard Model (SM) is complete. The next target at the energy frontier will be to study the Higgs properties and to search for the next scale beyond the SM. Experimentally, the $H \rightarrow c\bar{c}$ channel would be extremely difficult to dig out because of both the weak Yukawa coupling and the daunting SM di-jet background. We propose to test the charm-quark Yukawa coupling at the LHC and future hadron colliders with the Higgs boson decay to J/ψ via the charm-quark fragmentation. Using the non-relativistic quantum chromodynamics (NRQCD), we study the Higgs decay channel $H \rightarrow c\bar{c} + J/\psi$ (or η_c), where both the color-singlet and color-octet contributions are considered. Our result opens another door to improve determinations at the LHC of the Higgs Yukawa couplings: the final state from this decay mode is quite distinctive with $J/\psi \rightarrow e^+e^-$, $\mu^+\mu^-$ and the branching fraction is logarithmically enhanced by the charm-quark fragmentation mechanism.

In-person participation

No

Primary authors: Prof. LEIBOVICH, Adam (University of Pittsburgh); MA, Yang (University of Pittsburgh); Prof. HAN, Tao (University of Pittsburgh); Mr TAN, Xiaoze (Harbin Institute of Technology)

Presenter: MA, Yang (University of Pittsburgh)

Session Classification: Higgs Physics

Track Classification: Higgs Physics