

Sterile Neutrino and Dipole Portal Explanations of the MiniBooNE Excess



Nicholas Kamp, A. Schneider, M. Hostert, S. Vergani, A. Diaz, C.A. Argüelles, J.M. Conrad, M.H. Shaevitz, M.A. Uchida & the MiniBooNE Collaboration


2201.01724

2206.07100



The MiniBooNE Excess and Sterile Neutrinos

- The simplest explanation of the LSND $\bar{\nu}_e$ -like excess and MiniBooNE electron-like excess invokes oscillations involving an eV-scale sterile neutrino
- Recent MicroBooNE results [1] have disfavored a generic excess of electron neutrinos in the Booster Neutrino Beam, but do not rule out MiniBooNE's allowed region in oscillation parameter space [2, 3]
- The MiniBooNE collaboration has recently performed a combined 3+1 fit using MiniBooNE data and the MicroBooNE CCQE result [2]





- Even so, an eV-scale sterile neutrino is not able to explain the lowest energy and most forward parts of the MiniBooNE excess.
- Additionally, removing MiniBooNE reduces tension in global 3+1 oscillation fits by $\sim 2\sigma$ [4]

$$p_{PG} = \begin{cases} 8 \times 10^{-7} (4.8\sigma) & \text{w/ MiniBooNE} \\ 7 \times 10^{-3} (2.5\sigma) & \text{w/o MiniBooNE} \end{cases}$$

- This motivates the study of more exotic BSM models in addition to standard 3+1 oscillations




The Dipole Portal Model

- To accommodate the remaining MiniBooNE excess, we consider the addition of a dipole-coupled heavy neutral lepton (HNL)
- MINERvA elastic scattering measurements [5] are also sensitive to HNL decays to photons
- NuMI energies are too large to be sensitive to oscillations
- Use LeptonInjector [6] to simulate HNL production and decay



Dipole Fit Results

- Energy and angular distributions of the MiniBooNE excess prefer different regions of dipole parameter space
- Allowed regions for each distribution overlap at the 2σ level
- MINERvA elastic scattering data places stringent limits on the MiniBooNE preferred regions [7]



Conclusions

- The recent MicroBooNE results do not rule out the MiniBooNE preferred region in 3+1 parameter space
- The addition of a dipole-coupled HNL can alleviate tension in 3+1 global fits while retaining a reasonable explanation of the MiniBooNE excess
- MINERvA elastic scattering measurements place constraints on MiniBooNE preferred region in dipole parameter space; however, allowed regions still remain at the 2σ - 3σ level
- Most stringent constraints come from the unpublished antineutrino analysis

References and Acknowledgements

- [1] MicroBooNE Collaboration. *Phys. Rev. Lett.* 128, 241801 (2021)
- [2] MiniBooNE Collaboration. 2201.01724 (2022)
- [3] Argüelles et al. *Phys. Rev. Lett.* 128, 241802 (2021)
- [4] Vergani et al. *Phys. Rev. D* 104, 095005 (2021)
- [5] MINERvA Collaboration. *Phys. Rev. D* 100, 092001 (2019)
- [6] IceCube Collaboration. *Computer Physics Communications* 266, 108018 (2019)
- [7] Kamp et al. 2206.07100 (2022)

This work is supported by the NSF Graduate Research Fellowship under Grant No. 1745302.