



Contribution ID: 684

Type: Parallel Talk

Constraining 3-3-1 Models at the LHC and Future Hadron Colliders

Friday, 8 July 2022 18:30 (15 minutes)

In this work, we derive lower mass bounds on the Z' gauge boson based on the dilepton data from LHC with 13 TeV of center-of-mass energy, and forecast the sensitivity of the High-Luminosity-LHC with $L = 3000 fb^{-1}$, the High-Energy LHC with $\sqrt{s} = 27\text{-TeV}$, and also at the Future Circular Collider with $\sqrt{s} = 100\text{-TeV}$. We take into account the presence of exotic and invisible decays of the Z' gauge boson to find a more conservative and robust limit, different from previous studies. We investigate the impact of these new decay channels for several benchmark models in the scope of two different 3-3-1 models. We found that in the most constraining cases, LHC with $139 fb^{-1}$ can impose $m_{Z'} > 4\text{-TeV}$. Moreover, we forecast HL-LHC, HE-LHC, and FCC-hh collider reach, and derived the projected bounds $m_{Z'} > 5.8\text{-TeV}$, $m_{Z'} > 9.9\text{-TeV}$, and $m_{Z'} > 27\text{-TeV}$, respectively. Lastly, we put our findings into perspective with dark matter searches to show the region of parameter space where a dark matter candidate with the right relic density is possible.

In-person participation

No

Primary authors: Ms SÁNCHEZ VILLAMIZAR, Yoxara (IIP & DFTE - Federal University of Rio Grande do Norte (UFRN)); QUEIROZ, Farinaldo (Max Planck Institute for Nuclear Physics - Heidelberg); KLASSEN, Michael

Presenter: Ms SÁNCHEZ VILLAMIZAR, Yoxara (IIP & DFTE - Federal University of Rio Grande do Norte (UFRN))

Session Classification: Beyond the Standard Model

Track Classification: Beyond the Standard Model