



Contribution ID: 506

Type: **Parallel Talk**

Integrability as a new method for exact results on quasinormal modes of black holes

Friday, 8 July 2022 12:15 (15 minutes)

In this talk, I will show a new connection between quantum integrable models and black holes perturbation theory. After an introduction to quasinormal modes and their role in gravitational waves observations, I will connect their mathematically precise definition with the integrability structures derived from the ordinary differential equation associated to the black hole perturbation. More precisely, I will derive the full system of functional and non linear integral equations (Thermodynamic Bethe Ansatz) typical of quantum integrability and prove that the quasinormal modes verify different equivalent exact quantization conditions. As a consequence, it follows a new simple and effective method to numerically compute quasinormal modes which I will compare with other methods. I will also give a mathematical explanation of the recently found connection between quasinormal modes and $N=2$ supersymmetric gauge theories, through the further connection we previously found of these to quantum integrable models. All this I will show for a generalization of extremal Reissner-Nordström (charged) black holes, but in the end I will explain how it should be possible to generalize it to many other black holes, branes, fuzzballs, etc. and thus provide a new effective tool for the study of quantum gravity and gravitational waves.

In-person participation

Yes

Primary author: GREGORI, Daniele (Istituto Nazionale di Fisica Nucleare)**Presenter:** GREGORI, Daniele (Istituto Nazionale di Fisica Nucleare)**Session Classification:** Formal Theory**Track Classification:** Formal Theory