

A Monte Carlo Event Generator For New Physics in $B \rightarrow D^* \ell \nu$ decays

Lopamudra Mukherjee

University of Mississippi, USA


Talk Based on : **2203.07189, 2206.11283 [hep-ph]**

In Collaboration with *B.Bhattacharya, T.Browder, Q.Campagna, A.Datta,
S.Dubey & A.Sibidanov*




Flavour Changing Charged Current B-decay

- 1 Semileptonic decays are theoretically clean : Leptonic current is decoupled from the hadronic current.
- 2 Here, we focus on $B \rightarrow D^* \ell \nu$ because :
 - Useful in the extraction of $|V_{cb}|$ (See talks by *Taichiro Koga*).
 - Testing CKM unitarity.
 - Sensitive probes of New Physics.
 - Test Lepton Flavour Universality of the SM.
 - Persistent hints of NP in τ modes ($R_{D^{(*)}}$ etc.) and now in μ -modes.



See talks by *Resmi Puthumanaim, Manuel Naviglio, Yunxuan Li*

Angular Distribution in $B \rightarrow D^* \ell \nu$




- Belle provided the first time lepton-flavour specific single-differential distribution data for each of the 4 kinematic variables

Phys. Rev. D 100 (2019), 052007

- They used it to extract V_{cb} and test lepton universality ratio (μ/e) .
- The electron and muon data are in good agreement with SM.


Angular Observables



2104.02094

- Study of angular observables using the binned CP-averaged measurements of the four single-differential distributions provided by Belle done by Bobeth et. al
Phys. Rev. D 100 (2019), 052007
arxiv : 2104.02094
- Observables integrated over the entire q^2 range.
- Reports $> 2\sigma$ anomaly in A_{FB}^{μ} and $\sim 4\sigma$ anomaly in $\Delta A_{FB} = A_{FB}^{\mu} - A_{FB}^e$.

Angular Observables




2104.02094

- Study of angular observables using the binned CP-averaged measurements of the four single-differential distributions provided by Belle done by Bobeth et. al
Phys. Rev. D 100 (2019), 052007
arxiv : 2104.02094
- Observables integrated over the entire q^2 range.
- Reports $> 2\sigma$ anomaly in A_{FB}^{μ} and $\sim 4\sigma$ anomaly in $\Delta A_{FB} = A_{FB}^{\mu} - A_{FB}^e$.

Are these angular observables really clean?

Angular Observables



2104.02094

- Study of angular observables using the binned CP-averaged measurements of the four single-differential distributions provided by Belle done by Bobeth et. al
Phys. Rev. D 100 (2019), 052007
arxiv : 2104.02094
- Observables integrated over the entire q^2 range.
- Reports $> 2\sigma$ anomaly in A_{FB}^{μ} and $\sim 4\sigma$ anomaly in $\Delta A_{FB} = A_{FB}^{\mu} - A_{FB}^e$.

Are these angular observables really clean?

Is it possible to study the distribution of angular observables as function of q^2 ? -
Future physics goals at Belle II/LHCb upgrade

Forward-backward Asymmetry


$$\frac{d^2\Gamma}{dq^2 d\cos\theta_\ell} = \frac{d\Gamma}{dq^2} \left(\frac{1}{2} + A_{FB} \cos\theta_\ell + \frac{1 - 3\tilde{F}_L^\ell}{4} \frac{3\cos^2\theta_\ell - 1}{2} \right)$$

Measure of the no. of fermions produced in the forward region of the detector
vs. that produced in the backward region.

Forward-backward Asymmetry

$$\frac{d^2\Gamma}{dq^2 d\cos\theta_\ell} = \frac{d\Gamma}{dq^2} \left(\frac{1}{2} + A_{FB} \cos\theta_\ell + \frac{1 - 3\tilde{F}_L^\ell}{4} \frac{3\cos^2\theta_\ell - 1}{2} \right)$$


Measure of the no. of fermions produced in the forward region of the detector vs. that produced in the backward region.



- Difficult to disentangle NP from SM due to heavy dependence on form factors.
- We instead consider Δ -observables with potential sensitivity to NP.

Δ Angular Observables


Future of Measurements : $\Delta X = X^\mu - X^e$



- In case of SM, there is an almost exact cancellation of the hadronic uncertainties.
- $\Delta A_{FB}^{SM} \approx 0$ except at threshold where $A_{FB}^\ell \rightarrow -1$.
- Deviation from SM due to potential NP can be reliably extracted.

Δ Angular Observables

Future of Measurements : $\Delta X = X^\mu - X^e$



- In case of SM, there is an almost exact cancellation of the hadronic uncertainties.
- $\Delta A_{FB}^{SM} \approx 0$ except at threshold where $A_{FB}^\ell \rightarrow -1$.
- Deviation from SM due to potential NP can be reliably extracted.

What kinds of NP would provide potential signals in experiments?

MC for NP in $b \rightarrow c\ell\bar{\nu}$ decays

To answer this question we now have a new Monte-Carlo based on Evtgen:
https://github.com/qdcampagna/BTODSTARLNUNP_EVTGEN_Model

MC for NP in $b \rightarrow c\ell\bar{\nu}$ decays

To answer this question we now have a new Monte-Carlo based on Evtgen:

https://github.com/qdcampagna/BTODSTARLNUNP_EVTGEN_Model

$$\mathcal{H}_{\text{eff}} = \frac{G_F V_{cb}}{\sqrt{2}} \left\{ \begin{aligned} &(1 + g_L) [\bar{c}\gamma_\mu(1 - \gamma_5)b] [\bar{\ell}\gamma^\mu(1 - \gamma_5)\nu_\ell] \\ &+ g_R [\bar{c}\gamma_\mu(1 + \gamma_5)b] [\bar{\ell}\gamma^\mu(1 - \gamma_5)\nu_\ell] \\ &+ g_S [\bar{c}b] [\bar{\ell}(1 - \gamma_5)\nu_\ell] \\ &+ g_P [\bar{c}\gamma_5b] [\bar{\ell}(1 - \gamma_5)\nu_\ell] \\ &+ g_T [\bar{c}\sigma^{\mu\nu}(1 - \gamma_5)b] [\bar{\ell}\sigma_{\mu\nu}(1 - \gamma_5)\nu_\ell] \end{aligned} \right\} + h.c.$$

MC for NP in $b \rightarrow c \ell \bar{\nu}$ decays

To answer this question we now have a new Monte-Carlo based on Evtgen:

https://github.com/qdcampagna/BTODSTARLNUNP_EVTGEN_Model

$$\mathcal{H}_{\text{eff}} = \frac{G_F V_{cb}}{\sqrt{2}} \left\{ (1 + g_L) [\bar{c} \gamma_\mu (1 - \gamma_5) b] [\bar{\ell} \gamma^\mu (1 - \gamma_5) \nu_\ell] \right. \\ + g_R [\bar{c} \gamma_\mu (1 + \gamma_5) b] [\bar{\ell} \gamma^\mu (1 - \gamma_5) \nu_\ell] \\ + g_S [\bar{c} b] [\bar{\ell} (1 - \gamma_5) \nu_\ell] \\ + g_P [\bar{c} \gamma_5 b] [\bar{\ell} (1 - \gamma_5) \nu_\ell] \\ \left. + g_T [\bar{c} \sigma^{\mu\nu} (1 - \gamma_5) b] [\bar{\ell} \sigma_{\mu\nu} (1 - \gamma_5) \nu_\ell] \right\} + h.c.$$


Caveats :

- ❶ Neutrinos are always left-handed.
- ❷ The scalar matrix element $\langle D^* | \bar{c} b | \bar{B} \rangle = 0$
- ❸ SM case : $g_L = g_R = g_P = g_T = 0$
- ❹ Hadronic matrix elements are expressed in terms of form factors which are non-perturbative objects (cannot be calculated from first principles of QCD).


New Physics Analysis

- We pick out a few NP scenarios as listed below.
- The choice is motivated such that :
 - the ratio of semi-leptonic branching fractions is constrained to be within 3% of unity.
 - they are able to explain the “experimental” $\langle \Delta A_{FB} \rangle : 0.0349 \pm 0.0089$.
 - they also satisfy constraints on other angular observables such as $\langle \Delta F_L \rangle^{exp} = -0.0065 \pm 0.0059$ and $\langle \Delta \tilde{F}_L \rangle^{exp} = -0.0107 \pm 0.0142$.

	g_L	g_R	g_P
Scenario 1:	0.06	0.075	0.2 i
Scenario 2:	0.08	0.090	0.6 i
Scenario 3:	0.07	0.075	0



Distributions of Angular Asymmetries



MC shown for 50 ab⁻¹ data set in q^2 bins of 0.4 GeV²

- True CP violating observable S_7 in presence of complex new physics.


Theoretical Predictions & Belle II Sensitivities

	$\langle \Delta A_{FB} \rangle$ %	$\langle \Delta S_3 \rangle$ %	$\langle \Delta S_5 \rangle$ %	$\langle S_7 \rangle$ $\times 10^{-3}$
SM:	-0.252 ± 0.004	0.0441 ± 0.0007	0.0286 ± 0.0013	0
NP 1:	2.89 ± 0.05	1.08 ± 0.04	$2.44^{+0.02}_{-0.03}$	0.7 ± 0.01
NP 2:	$2.89^{+0.05}_{-0.06}$	$1.49^{+0.05}_{-0.04}$	$2.43^{+0.02}_{-0.03}$	2.0 ± 0.1
NP 3:	$2.94^{+0.04}_{-0.05}$	1.04 ± 0.04	$2.47^{+0.03}_{-0.02}$	0


- Here we use Belle **fiducial cuts** :

- $p_T^{\mu,e} > 0.8 \text{ GeV}$
- $p_T^\pi > 0.1 \text{ GeV}$
- Angular acceptance of all final state particles :
 $-0.866 < \cos \theta < 0.956$

- Note that we use the same p_T cut for electron and muon since we did not include detector efficiencies for the leptons separately.



Correlated Angular Asymmetries



- If there is NP, then one will observe signals in other angular asymmetries, not just in ΔA_{FB} .
- If experiments measure ΔA_{FB} in future without observing a ΔS_5 (say), then the signature does not indicate new physics.
- The right-handed coupling mainly drives correlation in absence of tensor NP.

Summary & Outlook

- **Distributions of angular asymmetries** in $B \rightarrow D^* \ell \nu$ are interesting and important.
- We expect angular asymmetries to provide tighter constraints on NP LFU couplings.
- We propose **Δ -observables** to be the future of experimental measurements for this mode.
- We now have a **MC generator** for NP studies in such decays.
- We have pointed out possible NP scenarios that can generate $\Delta A_{FB} \sim 3\%$ and can be extracted by experiments.
- A lot can be achieved at and beyond the 50 ab^{-1} of Belle II and other hadron colliders.

Summary & Outlook

- **Distributions of angular asymmetries** in $B \rightarrow D^* \ell \nu$ are interesting and important.
- We expect angular asymmetries to provide tighter constraints on NP LFU couplings.
- We propose **Δ -observables** to be the future of experimental measurements for this mode.
- We now have a **MC generator** for NP studies in such decays.
- We have pointed out possible NP scenarios that can generate $\Delta A_{FB} \sim 3\%$ and can be extracted by experiments.
- A lot can be achieved at and beyond the 50 ab^{-1} of Belle II and other hadron colliders.

THANK YOU!