IFAE2011 Incontri di Fisica delle Alte Energie



Contribution ID: 120

Type: not specified

The Kaon identification detector for the NA62 rare Kaon decay experiment at CERN

Wednesday, 27 April 2011 19:39 (1 minute)

The NA62 experiment at the CERN SPS aims to collect about 100 events of the ultra rare Kaon decay $K+ \rightarrow \pi + \nu vbar$, with a Signal to Background (S/B) ratio of 10:1. The main purpose is the measurement of the Branching Ratio of the decay with a 10% accuracy. NA62 will use an unseparated charged beam with Kaon decays in flight technique. Pions and protons cannot be separated efficiently from Kaons at the beam level. To make a positive identification of the ~ 6% of Kaons in the high rate environment (~800 MHz) before they decay a Hydrogen gas--filled differential Cherenkov counter (CEDAR) is placed in the incoming beam. The counter will be insensitive to pions and protons and it will give precise timing to reconstruct the K+ $\rightarrow \pi$ + $\nu vbar$ decay. The CEDAR detector is required to achieve a Kaon tagging efficiency of at least 95% with a time resolution of at least 100 ps.

Presenter: ROMANO, Angela (University of Birmingham) **Session Classification:** Sessione Poster

Track Classification: Dottorandi e Posters