



# Mapping the parameter space of low-scale leptogenesis

---


Juraj Klarić

based on works in collaboration with M. Drewes, A. Granelli, Y. Georis, S. Petcov, M.E. Shaposhnikov, I. Timiryasov

Neutrino Oscillation Workshop, September 5<sup>th</sup> 2022

# Some puzzles for physics beyond the Standard Model

Neutrino masses



The Baryon Asymmetry of the Universe

$$n_B/n_\gamma = 6.05(7) \times 10^{-10}$$





Image credits: Kamioka Observatory, ICRR, U. Tokyo; ESA and the Planck Collaboration


# Some puzzles for physics beyond the Standard Model

## Neutrino masses




[Minkowski 1977...]

## The Baryon Asymmetry of the Universe




[Fukugita/Yanagida '86...]

Image credits: Kamioka Observatory, ICRR, U. Tokyo; ESA and the Planck Collaboration




# Leptogenesis mechanisms




## Sakharov conditions

1. Baryon number violation  
sphaleron processes
2.  $C$  and  $CP$  violation  
RHN decays and oscillations
3. Deviation from thermal equilibrium  
freeze-in and freeze-out of RHN




- for hierarchical RHN  $M_1 \gtrsim 10^9 \text{ GeV}$

# Leptogenesis mechanisms




## Sakharov conditions

1. Baryon number violation  
sphaleron processes
2.  $C$  and  $CP$  violation  
RHN decays and oscillations
3. Deviation from thermal equilibrium  
freeze-in and freeze-out of RHN




- for hierarchical RHN  $M_1 \gtrsim 10^9$  GeV
- leptogenesis works in a wide range of RHN masses

# Leptogenesis mechanisms




## Sakharov conditions

1. Baryon number violation  
sphaleron processes
2.  $C$  and  $CP$  violation  
RHN decays and oscillations
3. Deviation from thermal equilibrium  
freeze-in and freeze-out of RHN




- for hierarchical RHN  $M_1 \gtrsim 10^9$  GeV
- leptogenesis works in a wide range of RHN masses

# Leptogenesis mechanisms




## Sakharov conditions

1. Baryon number violation  
sphaleron processes
2.  $C$  and  $CP$  violation  
RHN decays and oscillations
3. Deviation from thermal equilibrium  
freeze-in and freeze-out of RHN




- for hierarchical RHN  $M_1 \gtrsim 10^9$  GeV
- leptogenesis works in a wide range of RHN masses

# Leptogenesis mechanisms



## Sakharov conditions

1. Baryon number violation  
sphaleron processes
2.  $C$  and  $CP$  violation  
RHN decays and oscillations
3. Deviation from thermal equilibrium  
freeze-in and freeze-out of RHN




- for hierarchical RHN  $M_1 \gtrsim 10^9$  GeV
- leptogenesis works in a wide range of RHN masses
- how are the low-scale mechanisms connected?

# Resonant leptogenesis

- the BAU is mainly produced in RHN decays
- The lepton asymmetries follow the equation

$$\frac{dY_{\ell_a}}{dz} = -\epsilon_a \frac{\Gamma_N}{Hz} (Y_N - Y_N^{\text{eq}}) - W_{ab} Y_{\ell_b}$$



The key quantity determining the BAU is the decay asymmetry

$$\epsilon_a \equiv \frac{\Gamma_{N \rightarrow \ell_a} - \Gamma_{N \rightarrow \bar{\ell}_a}}{\Gamma_{N \rightarrow \ell_a} + \Gamma_{N \rightarrow \bar{\ell}_a}} = \frac{1}{8\pi} \frac{\text{Im}(F^\dagger F)_{12}^2}{(F^\dagger F)_{11}} \frac{M_1 M_2}{M_1^2 - M_2^2}$$

Becomes **enhanced** if  $M_2 \rightarrow M_1$  [(baryogenesis) Kuzmin '70] [(leptogenesis):

Liu/Segrè/Flanz/Paschos/Sarkar/Weiss/Covi/Roulet/Vissani/Pilaftsis/Underwood/Buchmüller/Plumacher...]


This enhancement is known as **resonant leptogenesis**.

- divergent when  $M_2 = M_1$ ?

# Resonant leptogenesis

- the BAU is mainly produced in RHN decays
- The lepton asymmetries follow the equation

$$\frac{dY_{\ell_a}}{dz} = -\epsilon_a \frac{\Gamma_N}{Hz} (Y_N - Y_N^{\text{eq}}) - W_{ab} Y_{\ell_b}$$



The key quantity determining the BAU is the decay asymmetry

$$\epsilon_a \equiv \frac{\Gamma_{N \rightarrow \ell_a} - \Gamma_{N \rightarrow \bar{\ell}_a}}{\Gamma_{N \rightarrow \ell_a} + \Gamma_{N \rightarrow \bar{\ell}_a}} = \frac{1}{8\pi} \frac{\text{Im}(F^\dagger F)_{12}^2}{(F^\dagger F)_{11}} \frac{M_1 M_2}{M_1^2 - M_2^2 + A^2}$$

Becomes **enhanced** if  $M_2 \rightarrow M_1$  [(baryogenesis) Kuzmin '70] [(leptogenesis):

Liu/Segrè/Flanz/Paschos/Sarkar/Weiss/Covi/Roulet/Vissani/Pilaftsis/Underwood/Buchmüller/Plumacher...]

This enhancement is known as **resonant leptogenesis**.

- divergent when  $M_2 = M_1$ ?
- divergence is unphysical – it needs to be regulated!
- this process can instead be described with **density matrix equations**

# Leptogenesis via oscillations





figure from [Drewes/Garbrecht/Gueter/JK 1606.06690]

# Quantum Kinetic Equations (QKEs)

## System of QKEs

$$i \frac{dn_{\Delta\alpha}}{dt} = -2i \frac{\mu_\alpha}{T} \int \frac{d^3k}{(2\pi)^3} \text{Tr} [\Gamma_\alpha] f_N (1 - f_N) \\ + i \int \frac{d^3k}{(2\pi)^3} \text{Tr} [\tilde{\Gamma}_\alpha (\bar{\rho}_N - \rho_N)],$$

$$i \frac{d\rho_N}{dt} = [H_N, \rho_N] - \frac{i}{2} \left\{ \Gamma, \rho_N - \rho_N^{eq} \right\} \\ - \frac{i}{2} \sum_\alpha \tilde{\Gamma}_\alpha \left[ 2 \frac{\mu_\alpha}{T} f_N (1 - f_N) \right],$$

$$i \frac{d\bar{\rho}_N}{dt} = -[H_N, \bar{\rho}_N] - \frac{i}{2} \left\{ \Gamma, \bar{\rho}_N - \rho_N^{eq} \right\} \\ + \frac{i}{2} \sum_\alpha \tilde{\Gamma}_\alpha \left[ 2 \frac{\mu_\alpha}{T} f_N (1 - f_N) \right],$$

- coupled system of integro-differential equations for the lepton flavor asymmetries  $n_{\Delta\alpha}$ , and the helicity-dependent HNL density matrices  $\rho_N$  and  $\bar{\rho}_N$
- HNL oscillations described by the effective hamiltonian  $H_N$
- equilibration described by helicity and flavor-dependent matrices  $\Gamma$

# Quantum Kinetic Equations (QKEs)

## System of QKEs

$$i \frac{dn_{\Delta\alpha}}{dt} = -2i \frac{\mu_\alpha}{T} \int \frac{d^3k}{(2\pi)^3} \text{Tr} [\Gamma_\alpha] f_N (1 - f_N) \\ + i \int \frac{d^3k}{(2\pi)^3} \text{Tr} [\tilde{\Gamma}_\alpha (\bar{\rho}_N - \rho_N)],$$

$$i \frac{d\rho_N}{dt} = [H_N, \rho_N] - \frac{i}{2} \left\{ \Gamma, \rho_N - \rho_N^{eq} \right\} \\ - \frac{i}{2} \sum_\alpha \tilde{\Gamma}_\alpha \left[ 2 \frac{\mu_\alpha}{T} f_N (1 - f_N) \right],$$

$$i \frac{d\bar{\rho}_N}{dt} = -[H_N, \bar{\rho}_N] - \frac{i}{2} \left\{ \Gamma, \bar{\rho}_N - \rho_N^{eq} \right\} \\ + \frac{i}{2} \sum_\alpha \tilde{\Gamma}_\alpha \left[ 2 \frac{\mu_\alpha}{T} f_N (1 - f_N) \right],$$

- coupled system of integro-differential equations for the lepton flavor asymmetries  $n_{\Delta\alpha}$ , and the helicity-dependent HNL density matrices  $\rho_N$  and  $\bar{\rho}_N$
- HNL oscillations described by the effective hamiltonian  $H_N$
- equilibration described by helicity and flavor-dependent matrices  $\Gamma$

# Quantum Kinetic Equations (QKEs)

## System of QKEs

$$i \frac{dn_{\Delta\alpha}}{dt} = -2i \frac{\mu_\alpha}{T} \int \frac{d^3k}{(2\pi)^3} \text{Tr} [\Gamma_\alpha] f_N (1 - f_N) \\ + i \int \frac{d^3k}{(2\pi)^3} \text{Tr} [\tilde{\Gamma}_\alpha (\bar{\rho}_N - \rho_N)],$$

$$i \frac{d\rho_N}{dt} = [H_N, \rho_N] - \frac{i}{2} \left\{ \Gamma, \rho_N - \rho_N^{eq} \right\} \\ - \frac{i}{2} \sum_\alpha \tilde{\Gamma}_\alpha \left[ 2 \frac{\mu_\alpha}{T} f_N (1 - f_N) \right],$$

$$i \frac{d\bar{\rho}_N}{dt} = -[H_N, \bar{\rho}_N] - \frac{i}{2} \left\{ \Gamma, \bar{\rho}_N - \rho_N^{eq} \right\} \\ + \frac{i}{2} \sum_\alpha \tilde{\Gamma}_\alpha \left[ 2 \frac{\mu_\alpha}{T} f_N (1 - f_N) \right],$$

- coupled system of integro-differential equations for the lepton flavor asymmetries  $n_{\Delta\alpha}$ , and the helicity-dependent HNL density matrices  $\rho_N$  and  $\bar{\rho}_N$
- HNL oscillations described by the effective hamiltonian  $H_N$
- equilibration described by helicity and flavor-dependent matrices  $\Gamma$

# Quantum Kinetic Equations (QKEs)

## System of QKEs

$$i \frac{dn_{\Delta\alpha}}{dt} = -2i \frac{\mu_\alpha}{T} \int \frac{d^3k}{(2\pi)^3} \text{Tr} [\Gamma_\alpha] f_N (1 - f_N) + i \int \frac{d^3k}{(2\pi)^3} \text{Tr} [\tilde{\Gamma}_\alpha (\bar{\rho}_N - \rho_N)],$$

$$i \frac{d\rho_N}{dt} = [H_N, \rho_N] - \frac{i}{2} \left\{ \Gamma, \rho_N - \rho_N^{eq} \right\} - \frac{i}{2} \sum_\alpha \tilde{\Gamma}_\alpha \left[ 2 \frac{\mu_\alpha}{T} f_N (1 - f_N) \right],$$

$$i \frac{d\bar{\rho}_N}{dt} = -[H_N, \bar{\rho}_N] - \frac{i}{2} \left\{ \Gamma, \bar{\rho}_N - \rho_N^{eq} \right\} + \frac{i}{2} \sum_\alpha \tilde{\Gamma}_\alpha \left[ 2 \frac{\mu_\alpha}{T} f_N (1 - f_N) \right],$$

- coupled system of integro-differential equations for the lepton flavor asymmetries  $n_{\Delta\alpha}$ , and the helicity-dependent HNL density matrices  $\rho_N$  and  $\bar{\rho}_N$
- HNL oscillations described by the effective hamiltonian  $H_N$
- equilibration described by helicity and flavor-dependent matrices  $\Gamma$

# Quantum Kinetic Equations (QKEs)

## System of QKEs

$$i \frac{dn_{\Delta\alpha}}{dt} = -2i \frac{\mu_\alpha}{T} \int \frac{d^3k}{(2\pi)^3} \text{Tr} [\Gamma_\alpha] f_N (1 - f_N) \\ + i \int \frac{d^3k}{(2\pi)^3} \text{Tr} [\tilde{\Gamma}_\alpha (\bar{\rho}_N - \rho_N)],$$

$$i \frac{d\rho_N}{dt} = [H_N, \rho_N] - \frac{i}{2} \left\{ \Gamma, \rho_N - \rho_N^{eq} \right\} \\ - \frac{i}{2} \sum_\alpha \tilde{\Gamma}_\alpha \left[ 2 \frac{\mu_\alpha}{T} f_N (1 - f_N) \right],$$

$$i \frac{d\bar{\rho}_N}{dt} = -[H_N, \bar{\rho}_N] - \frac{i}{2} \left\{ \Gamma, \bar{\rho}_N - \rho_N^{eq} \right\} \\ + \frac{i}{2} \sum_\alpha \tilde{\Gamma}_\alpha \left[ 2 \frac{\mu_\alpha}{T} f_N (1 - f_N) \right],$$

- coupled system of integro-differential equations for the lepton flavor asymmetries  $n_{\Delta\alpha}$ , and the helicity-dependent HNL density matrices  $\rho_N$  and  $\bar{\rho}_N$
- HNL oscillations described by the effective hamiltonian  $H_N$
- equilibration described by helicity and flavor-dependent matrices  $\Gamma$

# Quantum Kinetic Equations (QKEs)

## System of QKEs

$$i \frac{dn_{\Delta\alpha}}{dt} = -2i \frac{\mu_\alpha}{T} \int \frac{d^3k}{(2\pi)^3} \text{Tr} [\Gamma_\alpha] f_N (1 - f_N) \\ + i \int \frac{d^3k}{(2\pi)^3} \text{Tr} [\tilde{\Gamma}_\alpha (\bar{\rho}_N - \rho_N)],$$

$$i \frac{d\rho_N}{dt} = [H_N, \rho_N] - \frac{i}{2} \left\{ \Gamma, \rho_N - \rho_N^{eq} \right\} \\ - \frac{i}{2} \sum_\alpha \tilde{\Gamma}_\alpha \left[ 2 \frac{\mu_\alpha}{T} f_N (1 - f_N) \right],$$

$$i \frac{d\bar{\rho}_N}{dt} = - [H_N, \bar{\rho}_N] - \frac{i}{2} \left\{ \Gamma, \bar{\rho}_N - \rho_N^{eq} \right\} \\ + \frac{i}{2} \sum_\alpha \tilde{\Gamma}_\alpha \left[ 2 \frac{\mu_\alpha}{T} f_N (1 - f_N) \right],$$

- similar sets of equations derived using different strategies for both regimes
- for resonant leptogenesis relativistic corrections were typically negligible helicity effects could be neglected  $\rho_N \approx \bar{\rho}_N^*$
- leptogenesis via oscillations assumed ultra-relativistic HNLs non-relativistic corrections found to be important in recent years  
[Hambye/Teresi '16; Laine/Ghiglieri '17; Eijima/Shaposhnikov '17]
- gradual convergence towards the same set of equations

# The low-scale leptogenesis mechanisms

## Resonant leptogenesis

- often sufficient to use **decay asymmetries**  $\epsilon_a$
- conceptual issues arise when  $M_2 \rightarrow M_1$
- **relativistic effects** can typically be neglected
- heavy neutrino decays require  $M \gtrsim T$ , not clear what happens for  $M \lesssim 130 \text{ GeV}$
  
- both can be described by the **same density-matrix equations**


## Leptogenesis via oscillations

- initial conditions are crucial, all BAU is generated during RHN **equilibration (freeze-in)**
- important to distinguish the **helicities** of the RHN
- the decay of the RHN equilibrium distribution can typically be neglected  $Y_N^{\text{eq}} \approx 0$

# The parameter space of low-scale leptogenesis


## Resonant leptogenesis

- early estimates lead to successful leptogenesis for  $\mathcal{O}(200)$  GeV [Pilaftsis/Underwood '05]
- Higgs decay leptogenesis mechanism proposed in [Hambye/Teresi '16; '17]



## Leptogenesis via oscillations

- for  $M_M > M_W$  new channels open up
- large equilibration rates for both FNV and FNC processes
- generically we have  $\Gamma_N/H \gtrsim 30$  for  $T \sim 150$  GeV,  $M \sim 80$  GeV
- early estimate [Blondel/Graverini/Serra/Shaposhnikov 2014]




- Baryogenesis window closes at  $M_M \sim 80$  GeV?

A quantitative study is necessary!

# How to navigate the parameter space

- we use a single set of equations for both leptogenesis
  - for  $M \gg T$  we recover resonant leptogenesis
  - for  $M \ll T$  we recover leptogenesis via oscillations
- we separate the **freeze-in** and **freeze-out** regimes
  - for thermal initial conditions **freeze-out** is the only source of BAU: “resonant” leptogenesis dominates
  - for vanishing initial conditions with  $Y_N^{eq} \rightarrow 0$  **freeze-in** is the only source of BAU: LG via oscillations dominates
- biggest challenge: **rates!**
  - so far estimates of the rates only exist for  $M \ll T$  and  $M \gg T$
  - we combine the two by *extrapolating* the relativistic rate and adding it to the non-relativistic decays
- we perform a comprehensive numerical scan over the parameters between  $100 \text{ MeV} < M_M < 10 \text{ TeV}$

# Results: The minimal model with 2 RHNs




[JK/Timiryasov/Shaposhnikov 2103.16545]

- the baryogenesis window remains open!
- two main contributions to the BAU, from freeze-in and freeze-out
- there is significant overlap of the two regimes

- in resonant leptogenesis freeze-out (HNL decays) dominates, we can start with thermal initial conditions  $Y_N(0) = Y_N^{\text{eq}}$
- leptogenesis via oscillations is freeze-in dominated,  $Y_N(0) = 0$ , we set the “source” term to  $dY_N^{\text{eq}}/dz \rightarrow 0$  by hand
- results depend on low-energy CP phases:
  - optimal phases  $\delta = 0$  and  $\eta = \pi/2$
  - less overlap for e.g.  $\delta = \pi$  and  $\eta = 0$
  - maximal  $\Delta M/M \lesssim 10^{-1} \rightarrow 10^{-3}$

# Results: The minimal model with 2 RHNs




[JK/Timiryasov/Shaposhnikov 2103.16545]

- the baryogenesis window remains open!
- two main contributions to the BAU, from freeze-in and freeze-out
- there is significant overlap of the two regimes

- in resonant leptogenesis freeze-out (HNL decays) dominates, we can start with thermal initial conditions  $Y_N(0) = Y_N^{\text{eq}}$
- leptogenesis via oscillations is freeze-in dominated,  $Y_N(0) = 0$ , we set the “source” term to  $dY_N^{\text{eq}}/dz \rightarrow 0$  by hand
- results depend on low-energy CP phases:
  - optimal phases  $\delta = 0$  and  $\eta = \pi/2$
  - less overlap for e.g.  $\delta = \pi$  and  $\eta = 0$
  - maximal  $\Delta M/M \lesssim 10^{-1} \rightarrow 10^{-3}$

# Results: The minimal model with 2 RHNs




[JK/Timiryasov/Shaposhnikov 2103.16545]

- the baryogenesis window remains open!
- two main contributions to the BAU, from freeze-in and freeze-out
- there is significant overlap of the two regimes

- in resonant leptogenesis freeze-out (HNL decays) dominates, we can start with thermal initial conditions  $Y_N(0) = Y_N^{\text{eq}}$
- leptogenesis via oscillations is freeze-in dominated,  $Y_N(0) = 0$ , we set the “source” term to  $dY_N^{\text{eq}}/dz \rightarrow 0$  by hand
- results depend on low-energy CP phases:
  - optimal phases  $\delta = 0$  and  $\eta = \pi/2$
  - less overlap for e.g.  $\delta = \pi$  and  $\eta = 0$
  - maximal  $\Delta M/M \lesssim 10^{-1} \rightarrow 10^{-3}$

# Results: The minimal model with 2 RHNs




[JK/Timiryasov/Shaposhnikov 2103.16545]

- the baryogenesis window remains open!
- two main contributions to the BAU, from freeze-in and freeze-out
- there is significant overlap of the two regimes

- in resonant leptogenesis freeze-out (HNL decays) dominates, we can start with thermal initial conditions  $Y_N(0) = Y_N^{\text{eq}}$
- leptogenesis via oscillations is freeze-in dominated,  $Y_N(0) = 0$ , we set the “source” term to  $dY_N^{\text{eq}}/dz \rightarrow 0$  by hand
- results depend on low-energy CP phases:
  - optimal phases  $\delta = 0$  and  $\eta = \pi/2$
  - less overlap for e.g.  $\delta = \pi$  and  $\eta = 0$
  - maximal  $\Delta M/M \lesssim 10^{-1} \rightarrow 10^{-3}$

# Results: The minimal model with 2 RHNs




[JK/Timiryasov/Shaposhnikov 2103.16545]

- the baryogenesis window remains open!
- two main contributions to the BAU, from freeze-in and freeze-out
- there is significant overlap of the two regimes

- in resonant leptogenesis freeze-out (HNL decays) dominates, we can start with thermal initial conditions  $Y_N(0) = Y_N^{\text{eq}}$
- leptogenesis via oscillations is freeze-in dominated,  $Y_N(0) = 0$ , we set the “source” term to  $dY_N^{\text{eq}}/dz \rightarrow 0$  by hand
- results depend on low-energy CP phases:
  - optimal phases  $\delta = 0$  and  $\eta = \pi/2$
  - less overlap for e.g.  $\delta = \pi$  and  $\eta = 0$
  - maximal  $\Delta M/M \lesssim 10^{-1} \rightarrow 10^{-3}$

# Results: The minimal model with 2 RHNs




[JK/Timiryasov/Shaposhnikov 2103.16545]

- the baryogenesis window remains open!
- two main contributions to the BAU, from freeze-in and freeze-out
- there is significant overlap of the two regimes

- in resonant leptogenesis freeze-out (HNL decays) dominates, we can start with thermal initial conditions  $Y_N(0) = Y_N^{\text{eq}}$
- leptogenesis via oscillations is freeze-in dominated,  $Y_N(0) = 0$ , we set the “source” term to  $dY_N^{\text{eq}}/dz \rightarrow 0$  by hand
- results depend on low-energy CP phases:
  - optimal phases  $\delta = 0$  and  $\eta = \pi/2$
  - less overlap for e.g.  $\delta = \pi$  and  $\eta = 0$
  - maximal  $\Delta M/M \lesssim 10^{-1} \rightarrow 10^{-3}$

# Results: Leptogenesis with 3 RHNs




[ Snowmass White Paper 2203.08039]

leptogenesis bounds from [ Drewes/Georis/JK 2106.16226]

- for experimentally accessible heavy neutrino masses, all  $U^2$  are allowed
- both **freeze-in** and **freeze-out** leptogeneses already testable at existing experiments
- the maximal value of  $U^2$  depends on  $m_1$

# Results: Leptogenesis with 3 RHNs




[ Snowmass White Paper 2203.08039]

leptogenesis bounds from [ Drewes/Georis/JK 2106.16226]

- for experimentally accessible heavy neutrino masses, all  $U^2$  are allowed
- both **freeze-in** and **freeze-out** leptogeneses already testable at existing experiments
- the maximal value of  $U^2$  depends on  $m_1$

# Results: Leptogenesis with 3 RHNs




[ Snowmass White Paper 2203.08039]

leptogenesis bounds from [ Drewes/Georis/JK 2106.16226]

- for experimentally accessible heavy neutrino masses, all  $U^2$  are allowed
- both freeze-in and freeze-out leptogeneses already testable at existing experiments
- the maximal value of  $U^2$  depends on  $m_1$


# Indirect probes: Charged LFV



[Graneli/JK/Petcov 2206.04342]

- parameters space in the TeV region already severely constrained by cLFV observables
- future  $\mu \rightarrow e$  conversion experiments can probe a large part of the  $N = 3$  parameter space


# Indirect probes: Charged LFV



[Granelli/JK/Petcov 2206.04342]


- parameters space in the TeV region already severely constrained by cLFV observables
- future  $\mu \rightarrow e$  conversion experiments can probe a large part of the  $N = 3$  parameter space

# Allowed ranges of mixing angles $U^2$ and mass splittings $\Delta M$




- benchmark with fixed  $U_{\alpha I}^2/U^2$
- upper bound on  $U^2$  arises through a combination of baryogenesis + fine tuning constraints
- leptogenesis consistent with both LNV and LNC RHN decays
- nontrivial LNV/LNC ratios can further constrain the RHN parameters

# Allowed ranges of mixing angles $U^2$ and mass splittings $\Delta M$



- benchmark with fixed  $U_{\alpha I}^2/U^2$
- upper bound on  $U^2$  arises through a combination of baryogenesis + fine tuning constraints
- leptogenesis consistent with both LNV and LNC RHN decays
- nontrivial LNV/LNC ratios can further constrain the RHN parameters


# Allowed ranges of mixing angles $U^2$ and mass splittings $\Delta M$



- benchmark with fixed  $U^2_{\alpha I}/U^2$
- upper bound on  $U^2$  arises through a combination of baryogenesis + fine tuning constraints
- leptogenesis consistent with both LNV and LNC RHN decays
- nontrivial LNV/LNC ratios can further constrain the RHN parameters

[ Drewes/Georis/JK 220x.xxxx ]

# Allowed ranges of mixing angles $U^2$ and mass splittings $\Delta M$



- benchmark with fixed  $U^2_{\alpha I}/U^2$
- upper bound on  $U^2$  arises through a combination of baryogenesis + fine tuning constraints
- leptogenesis consistent with both LNV and LNC RHN decays
- nontrivial LNV/LNC ratios can further constrain the RHN parameters

[ Drewes/Georis/JK 220x.xxxx ]

# Conclusions

- resonant leptogenesis and leptogenesis through neutrino oscillations are really **two regimes of the same mechanism**
- freeze-out is already **possible for GeV-scale** RHNs
- freeze-in remains **important at the TeV-scale** and beyond
- leptogenesis is a viable baryogenesis mechanism for **all heavy neutrino masses** above the  $\mathcal{O}(100)$  MeV scale
- leptogenesis is testable at planned future experiments
  - synergy between **high-energy** and **high-intensity** frontiers!
  - together they can cover a large portion of the low-scale leptogenesis parameter space

Thank you!

# Large mixing angles and approximate B-L symmetry

- large  $U^2$  require cancellations between different entries of the Yukawa matrices  $F$
- this cancellation can be associated with an approximate lepton number symmetry

[Shaposhnikov hep-ph/0605047, Kersten Smirnov

0705.3221, Moffat Pascoli Weiland 1712.07611]

- symmetry broken by small parameters  $\epsilon, \epsilon', \mu, \mu'$

## Pseudo-Dirac pairs

$$N_s = \frac{N_1 + iN_2}{\sqrt{2}}, N_w = \frac{N_1 - iN_2}{\sqrt{2}}$$

## B-L parametrisation

$$M_M = \bar{M} \begin{pmatrix} 1 - \mu & 0 & 0 \\ 0 & 1 + \mu & 0 \\ 0 & 0 & \mu' \end{pmatrix}$$

$$F = \frac{1}{\sqrt{2}} \begin{pmatrix} F_e(1 + \epsilon_e) & iF_e(1 - \epsilon_e) & F_e\epsilon'_e \\ F_\mu(1 + \epsilon_\mu) & iF_\mu(1 - \epsilon_\mu) & F_\mu\epsilon'_\mu \\ F_\tau(1 + \epsilon_\tau) & iF_\tau(1 - \epsilon_\tau) & F_\tau\epsilon'_\tau \end{pmatrix}$$


# Fine tuning

- if present, symmetries are manifest to all orders in p.t.
- in the case of a large B-L breaking, radiative corrections can cause large neutrino masses
- we can use the size of radiative corrections to the light neutrino masses to quantify tuning

## Fine Tuning


$$f.t.(m_\nu) = \sqrt{\sum_{i=1}^3 \left( \frac{m_i^{\text{loop}} - m_i^{\text{tree}}}{m_i^{\text{loop}}} \right)^2}$$

# Slices of the parameter space




- two characteristic mass splittings
- mass splitting induced by the Higgs  $\Delta M_{\theta\theta}$
- mass splitting induced by RG running  $\delta M_{RG}$

# Slices of the parameter space




- two characteristic mass splittings
- mass splitting induced by the Higgs  $\Delta M_{\theta\theta}$
- mass splitting induced by RG running  $\delta M_{RG}$

# Slices of the parameter space



- two characteristic mass splittings
- mass splitting induced by the Higgs  $\Delta M_{\theta\theta}$
- mass splitting induced by RG running  $\delta M_{RG}$

# Results: Leptogenesis with 3 RHN (Normal Ordering)




# Hierarchy in the washout

- lepton asymmetry can survive washout if hidden in a particular flavor
- washout suppression

$$f \equiv \frac{\Gamma_a}{\Gamma} \sim \frac{U_a^2}{U^2}$$

- for 2 RHN  $f > 5 \times 10^{-3}$
- for 3 RHN  $f \ll 1$  possible

2 RHNs:



[Snowmass White Paper 2203.08039]

[Drewes/Garbrecht/Gueter/JK 1609.09069]

[Caputo/Hernandez/Lopez-Pavon/Salvado 1704.08721]


# Hierarchy in the washout

- lepton asymmetry can survive washout if hidden in a particular flavor
- washout suppression

$$f \equiv \frac{\Gamma_a}{\Gamma} \sim \frac{U_a^2}{U^2}$$

- for 2 RHN  $f > 5 \times 10^{-3}$
- for 3 RHN  $f \ll 1$  possible

3 RHNs:



[ Drewes/Georis/JK 220x.xxxx ]

[ Chrzaszcz/Drewes/Gonzalo/Harz/Krishnamurthy/Weniger 1908.02302 ]


# Hierarchy in the washout

- lepton asymmetry can survive washout if hidden in a particular flavor
- washout suppression

$$f \equiv \frac{\Gamma_a}{\Gamma} \sim \frac{U_a^2}{U^2}$$

- for 2 RHN  $f > 5 \times 10^{-3}$
- for 3 RHN  $f \ll 1$  possible

3 RHNs:





[ Drewes/Georis/JK 220x.xxxx ]

[ Chrzaszcz/Drewes/Gonzalo/Harz/Krishnamurthy/Weniger 1908.02302 ]


# Enhancement due to level crossing

- in the  $B - L$  symmetric limit two heavy neutrinos form a pseudo-Dirac pair
- the “3rd” heavy neutrino can be heavier than the pseudo-Dirac pair
- for  $T \gg T_{EW}$ , the pseudo-Dirac pair also has a thermal mass

$T \gg T_{EW}$




$T \ll T_{EW}$




# Enhancement due to level crossing


## Heavy Neutrino Densities



## Lepton flavour asymmetries



## Heavy Neutrino correlations



## Lepton number asymmetry

