



On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

# On the origin of antimatter in cosmos

Zurab Berezhiani

University of L'Aquila and LNGS

NOW 2022, Ostuni, 4-11 Sept. 2022





# Contents

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary



# Anti-particles and anti-matter (antinulei)

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

From discovery of positron, 1930-32

*and all other antiparticles  
(antiproton, antineutron etc.)*



$$\left( i\hbar \frac{\partial}{\partial t} + \sum_{\alpha=1}^3 \alpha_{\alpha} p_{\alpha} \right) \psi(\mathbf{x}, t) = i\hbar \frac{\partial \psi(\mathbf{x}, t)}{\partial t}$$




... to a great vision 1967

Matter (Baryon asymmetry) in the early universe can be originated (from zero) by New Interactions which

- Violate  $B$  (now better  $B - L$ ) and also CP
- and go out-of-equilibrium at some early epoch

$$\sigma(bb \rightarrow \bar{b}\bar{b})/\sigma(\bar{b}\bar{b} \rightarrow bb) = 1 - \epsilon$$

$\epsilon \sim 10^{-9}$ : for every  $\sim 10^9$  processes *one unit of  $B$*   
is left in the universe after the process is frozen



*There should be no antimatter in the Universe!*

In any case, matter should dominate the entire visible Universe  
No antimatter domain can exist within the horizon!

– Cohen, De Rujula, Glashow 1997



# Antiparticles in cosmic rays ...

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

$$\Phi_{\bar{p}}/\Phi_p \sim 10^{-4} \quad \text{AMS-02}$$

can be produced as secondaries in collisions of cosmic rays with interstellar gas, or can be signature of Dark Matter annihilation?

Anti-deuteron test? [Donato, Fornengo, Maurin, 2008](#)

But 8 anti-helium candidates were observed by AMS-02:

$$\Phi(\overline{\text{He}})/\Phi(\text{He}) \sim 10^{-8} \quad \text{6 helium-3 and 2 helium-4}$$

The discovery of a single antihelium-4 nucleus is challenging to all known physics.

If true, this should be strong indication to non-trivial new physics

Some *specifically tuned* DM models could explain the flux of antihelium-3 – but they fail to explain antihelium-4 !

Ting promised that AMS-02 will publish the anti-nuclei data as soon as they see first sl anti-carbon




# My hypothesis ...

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

- we observed neutron stars (NS) and NS-NS gravitational mergers:  
merger rate  $R \sim (1 \div 10^3) \text{ Gpc}^{-3} \text{ yr}^{-1}$
- there can exist dark neutron stars (NS') built of mirror neutrons  $n'$
- the neutron-mirror neutron mixing induces  $n' \rightarrow \bar{n}$  transition  
– antimatter "eggs" grow inside NS'  $N_{\bar{b}} \sim 10^{57} \cdot t_{\text{NS}} / \tau_{nn'}$
- in NS'-NS' mergers the anti-nuclei are "liberated" with  $v \sim c$
- $n_{\bar{b}} \sim R N_{\bar{b}} \tau_{\text{surv}} \sim \left( \frac{R}{1 \text{ Gpc}^{-3} \text{ yr}^{-1}} \right) \left( \frac{N_{\bar{b}}}{10^{51}} \right) \left( \frac{\tau_{\text{surv}}}{10^{14} \text{ s}} \right) \times 10^{-7} \text{ cm}^{-3}$   
 $\tau_{\text{surv}} = (n_p \langle \sigma_{\text{ann}} v \rangle)^{-1} \simeq 10^{14} \times (1 \text{ cm}^{-3} / n_p)$





# Standard Model $SU(3) \times SU(2) \times U(1)$

## Matter and Antimatter

*fermions and anti-fermions :*

$$\begin{array}{llll}
 q_L = \begin{pmatrix} u_L \\ d_L \end{pmatrix}, & \ell_L = \begin{pmatrix} \nu_L \\ e_L \end{pmatrix}; & u_R, d_R, & e_R \\
 \text{B}=1/3 & \text{L}=1 & \text{B}=1/3 & \text{L}=1 \\
 \bar{q}_R = \begin{pmatrix} \bar{u}_R \\ \bar{d}_R \end{pmatrix}, & \bar{\ell}_R = \begin{pmatrix} \bar{\nu}_R \\ \bar{e}_R \end{pmatrix}; & \bar{u}_L, \bar{d}_L, & \bar{e}_L \\
 \text{B}=-1/3 & \text{L}=-1 & \text{B}=-1/3 & \text{L}=-1
 \end{array}$$



$\updownarrow$  CP



C and P are maximally broken in weak interactions  
(not respected by gauge interactions)

but CP:  $F_L \rightarrow F_L^c \equiv \bar{F}_R = C \bar{F}_L^T = C \gamma_0 (F_L)^*$  is a nearly good symmetry  
transforming **Left-handed matter**  $\rightarrow$  **Right-handed antimatter**

– broken *only* by complex phases of Yukawa couplings to Higgs doublet  $\phi$

$$\mathcal{L}_{\text{Yuk}} = Y_{ij} \bar{F}_{Ri} F_{Lj} \phi = Y_{ij} \bar{F}_{Li} F_{Lj} \phi + \text{h.c.} \quad + \theta\text{-term in QCD}$$

B and L are automatically conserved in (renormalizable) couplings:  
accidental global symmetries  $U(1)_B$  and  $U(1)_L$

**B–L is conserved also by non-perturbative effects**

B–L breaking needs New Physics



## B-L violation: Majorana masses of neutrinos

On the origin of  
antimatter in  
cosmos



Zurab Berezhiani

Summary

- $\frac{A}{M}(\ell\phi)(\ell\phi) \quad (\Delta L = 2)$   
induces Majorana masses of  
neutrinos:  $m_\nu \sim v^2/M$   
– seesaw mechanism



$M \simeq 10^{15}$  GeV is the scale of new physics beyond EW scale  $\langle\phi\rangle = v$   
 $\simeq$  Majorana masses of "new" singlet fermions (RH neutrinos)



Back to Sakharov: **baryon asymmetry of the Universe can be induced by L and CP-violation in decays:**  $\Gamma(N \rightarrow \ell\phi) \neq \Gamma(N \rightarrow \bar{\ell}\bar{\phi})$

"redistributed" to non-zero B via non-perturbative SM effects

– **Baryogenesis via Leptogenesis** – but the price is rather expensive



# Visible vs. Dark matter: $\Omega_D/\Omega_B \sim 1$ ?

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

Visible matter from Baryogenesis

$B$  ( $B - L$ ) & CP violation, Out-of-Equilibrium

$\rho_B = n_B m_B$ ,  $m_B \simeq 1 \text{ GeV}$ ,  $\eta = n_B/n_\gamma \sim 10^{-9}$

$\eta$  is model dependent on several factors:

coupling constants and CP-phases, particle degrees of freedom, mass scales and out-of-equilibrium conditions, etc.



• Sakharov 1967

Dark matter:  $\rho_D = n_X m_X$ , but  $m_X = ?$ ,  $n_X = ?$

$n_X$  is model dependent: DM particle mass and interaction strength (production and annihilation cross sections), freezing conditions, etc.

- Axion  $m_a \sim 10^{-5} \text{ eV}$   $n_a \sim 10^4 n_\gamma$  - CDM
- Neutrinos  $m_\nu \sim 10^{-1} \text{ eV}$   $n_\nu \sim n_\gamma$  - HDM (✗)
- Sterile  $\nu'$   $m_{\nu'} \sim 10 \text{ keV}$   $n_{\nu'} \sim 10^{-3} n_\nu$  - WDM
- Mirror baryons  $m_{B'} \sim 1 \text{ GeV}$   $n_{B'} \sim n_B$  - ???
- WIMP  $m_X \sim 1 \text{ TeV}$   $n_X \sim 10^{-3} n_B$  - CDM
- WimpZilla  $m_X \sim 10^{14} \text{ GeV}$   $n_X \sim 10^{-14} n_B$  - CDM





$$SU(3) \times SU(2) \times U(1) + SU(3)' \times SU(2)' \times U(1)'$$

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary



$$G \times G'$$

Regular world

Mirror world

Elementary Particles

Elementary Particles



- Two identical gauge factors, e.g.  $SU(5) \times SU(5)'$ , with identical field contents and Lagrangians:  $\mathcal{L}_{\text{tot}} = \mathcal{L} + \mathcal{L}' + \mathcal{L}_{\text{mix}}$
- Mirror sector ( $\mathcal{L}'$ ) is dark – or perhaps grey? ( $\mathcal{L}_{\text{mix}} \rightarrow$  portals)
- MM is similar to standard matter, (asymmetric/dissipative/atomic) but realized in somewhat different cosmological conditions ( $T'/T \ll 1$ )
- $G \leftrightarrow G'$  symmetry ( $Z_2$  or  $Z_2^{LR}$ ): no new parameters in  $\mathcal{L}'$
- Cross-interactions between O & M particles  
 $\mathcal{L}_{\text{mix}}$ : new operators – new parameters! limited only by experiment!



$$SU(3) \times SU(2) \times U(1) \quad \text{vs.} \quad SU(3)' \times SU(2)' \times U(1)'$$

Two possible parities: with and without chirality change

*fermions and anti-fermions :*

$$q_L = \begin{pmatrix} u_L \\ d_L \end{pmatrix}, \quad \ell_L = \begin{pmatrix} \nu_L \\ e_L \end{pmatrix}; \quad u_R, d_R, e_R$$

$B=1/3 \quad L=1 \quad B=1/3 \quad L=1$



$\updownarrow$  CP

$$\bar{q}_R = \begin{pmatrix} \bar{u}_R \\ \bar{d}_R \end{pmatrix}, \quad \bar{\ell}_R = \begin{pmatrix} \bar{\nu}_R \\ \bar{e}_R \end{pmatrix}; \quad \bar{u}_L, \bar{d}_L, \bar{e}_L$$

$B=-1/3 \quad L=-1 \quad B=-1/3 \quad L=-1$



*Mirror fermions and antifermions :*

$$q'_L = \begin{pmatrix} u'_L \\ d'_L \end{pmatrix}, \quad \ell'_L = \begin{pmatrix} \nu'_L \\ e'_L \end{pmatrix}; \quad u'_R, d'_R, e'_R$$

$B'=1/3 \quad L'=1 \quad B'=1/3 \quad L'=1$



$\updownarrow$  CP

$$\bar{q}'_R = \begin{pmatrix} \bar{u}'_R \\ \bar{d}'_R \end{pmatrix}, \quad \bar{\ell}'_R = \begin{pmatrix} \bar{\nu}'_R \\ \bar{e}'_R \end{pmatrix}; \quad \bar{u}'_L, \bar{d}'_L, \bar{e}'_L$$

$B'=-1/3 \quad L'=-1 \quad B'=-1/3 \quad L'=-1$



$$\mathcal{L}_{\text{Yuk}} = F_L Y \bar{F}_L \phi + \text{h.c.} \quad \mathcal{L}'_{\text{Yuk}} = F'_L Y' \bar{F}'_L \phi' + \text{h.c.}$$

$$Z_2: L(R) \leftrightarrow L'(R'): Y'_{u,d,e} = Y_{u,d,e} \quad B, L \leftrightarrow B', L'$$

$$Z_2^{LR}: L(R) \leftrightarrow R'(L'): Y'_{u,d,e} = Y_{u,d,e}^* \quad B, L \leftrightarrow -B', L' \quad Z_2^{LR} = Z_2 \times \text{CP}$$



## – Sign of mirror baryon asymmetry ?

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

Ordinary BA is positive:  $\mathcal{B} = \text{sign}(n_b - n_{\bar{b}}) = 1$   
– as produced by (unknown) baryogenesis a la Sakharov!

Sign of mirror BA,  $\mathcal{B}' = \text{sign}(n_{b'} - n_{\bar{b}'}),$  is a priori unknown!

Imagine a baryogenesis mechanism *separately* acting in O and M sectors!  
– without involving cross-interactions in  $\mathcal{L}_{\text{mix}}$

E.g. EW baryogenesis or leptogenesis  $N \rightarrow \ell \phi$  and  $N' \rightarrow \ell' \phi'$

$Z_2: \rightarrow Y'_{u,d,e} = Y_{u,d,e}$  i.e.  $\mathcal{B}' = 1$

– O and M sectors are CP-identical in same chiral basis! O=left, M=left

$Z_2^{LR}: \rightarrow Y'_{u,d,e} = Y_{u,d,e}^*$  i.e.  $\mathcal{B}' = -1$

– O sector in L-basis is identical to M sector in R-basis! O=left, M=right

In the absence of cross-interactions in  $\mathcal{L}_{\text{mix}}$  we cannot measure sign of BA  
(or chirality in weak interactions) in M sector – so all remains academic ...

But switching on cross-interactions, violating B and  $\mathcal{B}'$  – as neutron–mirror  
neutron mixing:  $\epsilon n' n + \text{h.c.}$  or  $\nu - \nu'$  mixing

$\mathcal{B}' = -1 \rightarrow \bar{n}' \rightarrow n$  M (anti)matter  $\rightarrow$  O matter but  $\bar{\nu}' \rightarrow \bar{\nu}$

$\mathcal{B}' = 1 \rightarrow n' \rightarrow \bar{n}$  M matter  $\rightarrow$  O antimatter but  $\nu' \rightarrow \nu$



# Quick overview of mirror dark matter ...

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

Parallel/mirror sector of particles as a duplicate of our SM:  $SM \times SM'$  (or  $SU(5) \times SU(5)'$  or  $E_8 \times E_8'$  or parallel branes ... or more sectors)  
– all our particles ( $e, p, n, \nu, \gamma \dots$ ) have dark M twins ( $e', p', n', \nu', \gamma' \dots$ ) of exactly (or almost) the same masses

M matter is viable DM (asymmetric/baryonic/atomic/self-interacting/dissipative etc. as ordinary (O) baryon matter) – but M sector must be colder than O sector:  $T'/T < 0.2$  or so (BBN, CMB, LSS etc.)

– asymmetric reheating between the two sectors after inflation



– O matter mainly hydrogen (H 75%,  $^4\text{He}$  25%)  
while M matter mostly helium (H' 25%,  $^4\text{He}'$  75%) – first M stars are formed earlier than O stars, are bigger, helium dominated and end up in heavy BH:  $M \sim (10 \div 10^2) M_\odot$  (inferring  $\sim 80\%$  of DM in galactic halo and for the rest of  $\sim 20\%$  – M gas clouds,  $\sim M_\odot$  stars etc.)

There can exist interactions between O and M particles, e.g. photon kinetic mixing  $\varepsilon F^{\mu\nu} F'_{\mu\nu}$ , some common gauge bosons, etc.  
Most interesting are the ones which violate baryon and lepton numbers between two sectors, and namely  $B - L$  and  $B' - L'$  which can co-generate baryon asymmetries in both sectors – and naturally explain why the DM and baryon fractions are comparable,  $\Omega_{B'}/\Omega_B \simeq 5$  or so




# B-L violation in O and M sectors: Active-sterile mixing

- $\frac{A}{M}(\ell\phi)(\ell\phi)$  ( $\Delta L = 2$ ) – neutrino (seesaw) masses  $m_\nu \sim v^2/M$   
 $M$  is the (seesaw) scale of new physics beyond EW scale.



- Neutrino -mirror neutrino mixing – (active - sterile mixing)  
 $L$  and  $L'$  violation:  $\frac{A}{M}(\ell\phi)(\ell\phi)$ ,  $\frac{A}{M}(\ell'\phi')(\ell'\phi')$  and  $\frac{B}{M}(\ell\phi)(l\ell'\phi')$



Mirror neutrinos as natural candidates for sterile neutrinos

ZB and Mohapatra 95, ZB, Dolgov and Mohapatra 96





## Co-leptogenesis: B-L violating interactions between O and M worlds

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

L and  $L'$  violating operators  $\frac{1}{M}(\ell\phi)(\ell\phi)$  and  $\frac{1}{M}(\ell\phi)(\ell'\phi')$  lead to processes  $\ell\phi \rightarrow \bar{\ell}\bar{\phi}$  ( $\Delta L = 2$ ) and  $\ell\phi \rightarrow \bar{\ell}'\bar{\phi}'$  ( $\Delta L = 1, \Delta L' = 1$ )



After inflation, our world is heated and mirror world is empty:  
but ordinary particle scatterings transform them into mirror particles,  
heating also mirror world.

- These processes should be **out-of-equilibrium**
- **Violate** baryon numbers in both worlds,  $B - L$  and  $B' - L'$
- **Violate** also CP, given complex couplings

Green light to celebrated conditions of Sakharov



# Co-leptogenesis:


Z.B. and Bento, PRL 87, 231304 (2001)

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

Operators  $\frac{1}{M}(l\bar{\phi})(l\bar{\phi})$  and  $\frac{1}{M}(l\bar{\phi})(l'\bar{\phi}')$  via seesaw mechanism –  
heavy RH neutrinos  $N_j$  with  
Majorana masses  $\frac{1}{2}Mg_{jk}N_jN_k + \text{h.c.}$



Complex Yukawa couplings  $Y_{ij}l_iN_j\bar{\phi} + Y'_{ij}l'_iN_j\bar{\phi}' + \text{h.c.}$

$Z_2$  (Xerox) symmetry  $\rightarrow Y' = Y$ ,


$Z_2^{LR}$  (Mirror) symmetry  $\rightarrow Y' = Y^*$



# Co-leptogenesis: Sign of Mirror BA

Z.B., arXiv:1602.08599

*Hot O World*  $\rightarrow$  *Cold M World*



$$\frac{dn_{\text{BL}}}{dt} + (3H + \Gamma)n_{\text{BL}} = \Delta\sigma n_{\text{eq}}^2$$

$$\frac{dn'_{\text{BL}}}{dt} + (3H + \Gamma')n'_{\text{BL}} = \Delta\sigma' n_{\text{eq}}^2$$

$$\sigma(l\phi \rightarrow \bar{l}\bar{\phi}) - \sigma(\bar{l}\bar{\phi} \rightarrow l\phi) = \Delta\sigma$$

$$\sigma(l\phi \rightarrow \bar{l}'\bar{\phi}') - \sigma(\bar{l}'\bar{\phi}' \rightarrow l'\phi') = -(\Delta\sigma + \Delta\sigma')/2 \rightarrow 0 \quad (\Delta\sigma = 0)$$

$$\sigma(l\phi \rightarrow l'\phi') - \sigma(\bar{l}'\bar{\phi}' \rightarrow \bar{l}\bar{\phi}) = -(\Delta\sigma - \Delta\sigma')/2 \rightarrow \Delta\sigma \quad (0)$$

$$\Delta\sigma = \text{Im Tr}[g^{-1}(Y^\dagger Y)^* g^{-1}(Y'^\dagger Y') g^{-2}(Y^\dagger Y)] \times T^2/M^4$$

$$\Delta\sigma' = \Delta\sigma(Y \rightarrow Y')$$

Mirror ( $Z_2^{LR}$ ):  $Y' = Y^* \rightarrow \Delta\sigma' = -\Delta\sigma \rightarrow B > 0, B' > 0$

Xerox ( $Z_2$ ):  $Y' = Y \rightarrow \Delta\sigma' = \Delta\sigma = 0 \rightarrow B, B' = 0$

If  $k = \left(\frac{\Gamma}{H}\right)_{T=T_R} \ll 1$ , neglecting  $\Gamma$  in eqs  $\rightarrow n_{\text{BL}} = n'_{\text{BL}}$

$$\Omega'_B = \Omega_B \simeq 10^3 \frac{J M_{\text{Pl}} T_R^3}{M^4} \simeq 10^3 J \left(\frac{T_R}{10^{11} \text{ GeV}}\right)^3 \left(\frac{10^{13} \text{ GeV}}{M}\right)^4$$






If  $k = \left(\frac{\Gamma_2}{H}\right)_{T=T_R} \sim 1$ , Boltzmann Eqs.

$$\frac{dn_{\text{BL}}}{dt} + (3H + \Gamma)n_{\text{BL}} = \Delta\sigma n_{\text{eq}}^2 \quad \frac{dn'_{\text{BL}}}{dt} + (3H + \Gamma')n'_{\text{BL}} = \Delta\sigma n_{\text{eq}}^2$$

should be solved with  $\Gamma$ :



$D(k) = \Omega_B/\Omega'_B$ ,  $x(k) = T'/T$  for different  $g_*(T_R)$  and  $\Gamma_1/\Gamma_2$ .

So we obtain  $\Omega'_B = 5\Omega_B$  when  $m'_B = m_B$  but  $n'_B = 5n_B$

– the reason: mirror world is colder



Now the neutrons: since 1932 they make 50% of mass in our bodies ...

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

Neutrons are closely mass degenerate with the proton  
(in the SM  $n = udd$ ,  $p = uud$ ) since  $B$  is conserved in the SM,  $n$  and  $p$  both are Dirac particles with  $B = 1$ )

Neutrons are stable in basic nuclei but decay in free state:  $n \rightarrow pe\bar{\nu}_e$   
... and decay also in some ( $\beta^-$  unstable) nuclei  
... and can be even born in other ( $\beta^+$  unstable) nuclei:  $p \rightarrow ne^+\nu_e$


*Fermi V-A Theory  $\rightarrow$  Standard Model*

$$\frac{G_V}{\sqrt{2}} \bar{u}(1 - \gamma^5)\gamma^\mu d \bar{\nu}_e(1 - \gamma^5)\gamma_\mu e + \text{h.c.}$$

$$G_V = G_F |V_{ud}|$$

$$\frac{G_V}{\sqrt{2}} \bar{p}(g_V - g_A\gamma^5)\gamma^\mu n \bar{\nu}_e(1 - \gamma^5)\gamma_\mu e + \text{h.c.}$$

$$g_V = 1 \text{ (CVC)} \quad \& \quad g_A \simeq 1.2 \text{ (PCAC)}$$



Yet, we do not know all its secrets in depth...



# Majorana mass of the neutron = $n - \bar{n}$ mixing

On the origin of  
antimatter in  
cosmos


Zurab Berezhiani

Summary

Neutron is a Dirac particle:  $m \bar{n} n$  ( $\Delta B = 0$ ) with  $m \simeq 1$  GeV

In principle, being neutral, it could have also a Majorana mass  $\epsilon(n^T C n + \bar{n}^T C \bar{n})$  ( $\Delta B = 2$ ) even with  $\epsilon$  larger than  $m$

But being composite, this Majorana mass can come only from six-fermions effective operator  $\frac{1}{M^5}(udd)(udd)$ ,  $M > 1$  TeV or so



$$\epsilon = \langle n | (udd)(udd) | \bar{n} \rangle \sim \frac{\Lambda_{\text{QCD}}^6}{M^5} \sim \left( \frac{10 \text{ TeV}}{M} \right)^5 \times 10^{-15} \text{ eV} \quad (\text{or } \sim 1 \text{ s}^{-1})$$

Induces transition  $n(udd) \rightarrow \bar{n}(\bar{u}\bar{d}\bar{d})$ , with oscillation time  $\tau_{n\bar{n}} = \epsilon^{-1}$

$$M > 10 \text{ TeV} \rightarrow \epsilon < 10^{-15} \text{ eV} \rightarrow \tau_{n\bar{n}} > 1 \text{ s}$$

$$M \sim 10^3 \text{ TeV} \rightarrow \epsilon \sim 10^{-25} \text{ eV} \rightarrow \tau_{n\bar{n}} \sim 10^{10} \text{ s}$$

$$\epsilon < 7.5 \times 10^{-24} \text{ eV} \rightarrow \tau_{n\bar{n}} > 0.86 \times 10^8 \text{ s} \quad \text{direct limit free } n$$



# $n - \bar{n}$ oscillation: Free (and bound)

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

Two states,  $n$  and  $\bar{n}$

$$H = \begin{pmatrix} m + \mu \vec{B} \vec{\sigma} - V_n & \epsilon \\ \epsilon & m - \mu \vec{B} \vec{\sigma} - V_{\bar{n}} \end{pmatrix}$$

Free oscillation probability  $P_{n\bar{n}}(t) = \frac{\epsilon^2}{\omega_B^2} \sin^2(\omega_B t)$ ,  $\epsilon \ll \omega_B = \mu B$

$$\omega_B t < 1 \rightarrow P_{n\bar{n}}(t) = (\epsilon t)^2 = (t/\tau_{n\bar{n}})^2$$

$$\omega_B t \gg 1 \rightarrow P_{n\bar{n}}(t) = \frac{1}{2}(\epsilon/\omega_B)^2 < \frac{(\epsilon t)^2}{(\omega_B t)^2}$$

for a given free flight time  $t$ , magn. field should be properly suppressed to achieve "quasi-free" regime:  $\omega_B t < 1$

Baldo-Ceolin et al, 1994 (ILL, Grenoble) :  $t \simeq 0.1$  s,  $B < 1$  mG

$$P_{n\bar{n}}(t) = (t/\tau_{n\bar{n}})^2 < 10^{-18} \rightarrow \epsilon < 7.7 \times 10^{-24} \text{ eV}$$

In nuclei:  $\Delta V = V_{\bar{n}} - V_n \sim 100 \text{ MeV}$   $\theta \simeq \epsilon/\Delta V < 10^{-23}$

$P_{n\bar{n}} \simeq \theta^2 \simeq (\epsilon/\Delta V)^2 < 10^{-46}$  – is unobservable? Not really ...



# $n - \bar{n}$ mixing and instability of matter (nuclei)

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

In principle,  $n - \bar{n}$  oscillation could be rather fast:

E.g. for  $M \sim 10$  TeV (otherwise safe scale for new physics)

one would have  $\tau_{n\bar{n}} \sim 1$  s  $\rightarrow P_{n\bar{n}}(t = 0.1 \text{ s}) \simeq (t/\tau_{n\bar{n}})^2 \simeq 10^{-4}$

However:  $n - \bar{n}$  oscillation destabilizes nuclei:

$(A, Z) \rightarrow (A - 1, \bar{n}, Z) \rightarrow (A - 2, Z/Z - 1) + \pi$ 's

Present bounds on  $\epsilon$  from nuclear stability

$\epsilon < 1.2 \times 10^{-24}$  eV  $\rightarrow \tau > 1.3 \times 10^8$  s Fe, Soudan 2002

$\epsilon < 2.5 \times 10^{-24}$  eV  $\rightarrow \tau > 2.7 \times 10^8$  s O, SK 2015

$\epsilon < 7.5 \times 10^{-24}$  eV  $\rightarrow \tau > 0.86 \times 10^8$  s direct limit free  $n$



# $B$ violating operators between $O$ and $M$ particles in $\mathcal{L}_{\text{mix}}$

On the origin of  
antimatter in  
cosmos



Zurab Berezhiani

Summary

Ordinary quarks  $u, d$  ( antiquarks  $\bar{u}, \bar{d}$ )  
Mirror quarks  $u', d'$  ( antiquarks  $\bar{u}', \bar{d}'$ )

- Neutron -mirror neutron mixing – (Active - sterile neutrons)

$$\frac{1}{M^5} (udd)(udd) \quad \& \quad \frac{1}{M^5} (udd)(u'd'd')$$



Oscillations  $n \rightarrow \bar{n}$  ( $\Delta B = 2$ )

Oscillations  $n \rightarrow \bar{n}'$  ( $\Delta B = 1, \Delta B' = 1$ )  $B - B'$  is conserved




# Neutron – mirror neutron mixing

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

Effective operator  $\frac{1}{M^5}(udd)(u'd'd')$   $\rightarrow$  mixing  $\epsilon n C n' + \text{h.c.}$   
violating  $B$  and  $B'$  – but conserving  $B - B'$



$$\epsilon = \langle n | (udd)(u'd'd') | \bar{n}' \rangle \sim \frac{\Lambda_{\text{QCD}}^6}{M^5} \sim \left( \frac{10 \text{ TeV}}{M} \right)^5 \times 10^{-15} \text{ eV}$$

Key observation:  *$n - \bar{n}'$  oscillation cannot destabilise nuclei:*  
 *$(A, Z) \rightarrow (A - 1, Z) + n' (p' e' \bar{\nu}')$  forbidden by energy conservation*  
(In principle, it can destabilise Neutron Stars)

For  $m_n = m_{n'}$ ,  $n - \bar{n}'$  oscillation can be as fast as  $\epsilon^{-1} = \tau_{nn'} \sim 1 \text{ s}$   
without contradicting experimental and astrophysical limits.  
(c.f.  $\tau > 10 \text{ yr}$  for neutron – antineutron oscillation)

Neutron disappearance  $n \rightarrow \bar{n}'$  and regeneration  $n \rightarrow \bar{n}' \rightarrow n$   
can be searched at small scale 'Table Top' experiments



# Free Neutrons: Where to find Them ?

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

Neutrons are making  $1/7$  fraction of baryon mass in the Universe.

But most of neutrons bound in nuclei ....

$n \rightarrow \bar{n}'$  conversions can be seen only with free neutrons ... and, under some parameters, it can explain the neutron lifetime puzzle !

Free neutrons are present only in

- Reactors and Spallation Facilities (experiments are looking for)
- In Cosmic Rays ( $n - n'$  can reconcile TA and Auger experiments)
- During BBN epoch (fast  $n' \rightarrow \bar{n}$  can solve Lithium problem)

– Transition  $n \rightarrow \bar{n}'$  can take place for (gravitationally bound)  
Neutron Stars – conversion of NS into mixed ordinary/mirror NS

We do not observe the strong effects since  $n \rightarrow \bar{n}'$  is suppressed by some environmental factors (matter, magnetic field) or simply by some mass splitting between  $n - n'$





# Neutron Stars: $n - n'$ conversion

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

Two states,  $n$  and  $n'$

$$H = \begin{pmatrix} m_n + V_n + \mu_n \vec{B} \vec{\sigma} & \epsilon \\ \epsilon & m'_n + V'_n - \mu_n \vec{B}' \vec{\sigma} \end{pmatrix}$$

$$n_1 = \cos \theta n + \sin \theta n', \quad n_2 = \sin \theta n - \cos \theta n', \quad \theta \simeq \frac{\epsilon}{V_n - V'_n}$$

Fermi degenerate neutron liquid  $p_F \simeq (n_b/0.3 \text{ fm}^{-3})^{2/3} \times 400 \text{ MeV}$

$nn \rightarrow nn'$  with rate  $\Gamma = 2\theta^2 \eta \langle \sigma v \rangle n_b$

$$\frac{dN}{dt} = -\Gamma N \quad \frac{dN'}{dt} = \Gamma N \quad N + N' = N_0 \text{ remains Const.}$$

$$\tau_\epsilon = \Gamma^{-1} = \epsilon_{15}^{-2} \left( \frac{M}{1.5 M_\odot} \right)^{2/3} \times 10^{15} \text{ yr} \quad N'/N_0 = t/\tau_\epsilon$$

for  $t = 10 \text{ Gyr}$ ,  $\tau_\epsilon = 10^{15} \text{ yr}$  gives  $M$  fraction  $10^{-5}$  – few Earth mass

$$\dot{\mathcal{E}} = \frac{E_F N}{\tau_\epsilon} = \left( \frac{10^{15} \text{ yr}}{\tau_\epsilon} \right) \left( \frac{M}{1.5 M_\odot} \right) \times 10^{31} \text{ erg/s} \quad \text{NS heating – surface T}$$



# Neutron Star transformation


On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

$$\frac{dN}{dt} = -\Gamma N \quad \frac{dN'}{dt} = \Gamma \quad N + N' = N_0 \quad \text{remains Const.}$$

Initial state  $N = N_0$ ,  $N' = 0$       final state  $N = N' = \frac{1}{2} N_0$



Quark stars: in strange quark matter (color-superconducting phase) transition is not energetically favored. So Quark stars (which perhaps are heavy pulsars with  $M \simeq 2 M_{\odot}$  or so) are insensitive to  $n \rightarrow n'$ .



# Mixed Neutron Stars: TOV and $M - R$ relations

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

$$g_{\mu\nu} = \text{diag}(-g_{tt}, g_{rr}, r^2, r^2 \sin^2 \theta) \quad g_{tt} = e^{2\phi}, \quad g_{rr} = \frac{1}{1-2m/r}$$

$$T_{\mu\nu} = T_{\mu\nu}^1 + T_{\mu\nu}^2 = \text{diag}(\rho g_{tt}, p g_{rr}, p r^2, p r^2 \sin^2 \theta)$$



$$\rho = \rho_1 + \rho_2 \quad \& \quad p = p_1 + p_2, \quad p_\alpha = F(\rho_\alpha)$$

$$\frac{dm}{dr} = 4\pi r^2 \rho \rightarrow \frac{dm_{1,2}}{dr} = 4\pi r^2 \rho_{1,2} \quad m = m_1 + m_2$$

$$\frac{d\phi}{dr} = -\frac{1}{\rho+p} \frac{dp}{dr} \rightarrow \frac{dp_1/dr}{\rho_1+p_1} = \frac{dp_2/dr}{\rho_2+p_2}$$

$$\frac{dp}{dr} = (\rho + p) \frac{m+4\pi p r^3}{2mr-r^2}$$

$$(m_1 \neq 0, m_2 = 0)_{\text{in}} \rightarrow (m_1 = m_2)_{\text{fin}} \quad r \rightarrow \frac{r}{\sqrt{2}}, \quad m_\alpha \rightarrow \frac{m_\alpha}{2\sqrt{2}}$$



$$\sqrt{2} \text{ rule: } M_{\text{mix}}^{\text{max}} = \frac{1}{\sqrt{2}} M_{\text{NS}}^{\text{max}} \quad R_{\text{mix}}(M) = \frac{1}{\sqrt{2}} R_{\text{NS}}(M)$$



# Transforming Dark Matter into Antimatter: $n$ or $\bar{n}$ ?

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

Cross-interactions can induce mixing of neutral particles between two sectors, e.g.  $\nu - \nu'$  oscillations (M neutrinos = sterile neutrinos)

Oscillation  $n \rightarrow n'$  can be very effective process, **faster than the neutron decay**. For certain parameters it can explain the neutron lifetime problem,  $4.5\sigma$  discrepancy between the decay times measured by different experimental methods (bottle and beam), or anomalous neutron losses observed in some experiments and paradoxes in the UHECR detections

$n \rightarrow n'$  transition can have observable effects on neutron stars. It creates dark cores of M matter in the NS interiors, or eventually can transform them into maximally mixed stars with equal amounts of O and M neutrons

Such transitions in mirror NS create O matter cores. If baryon asymmetry in M sector has opposite sign, transitions  $\bar{n}' \rightarrow \bar{n}$  create antimatter cores which can be seen by LAT **by accreting ordinary gas** and explain the origin of anti-helium nuclei in cosmic rays **supposedly seen by AMS2**



# Mergers of neutron stars .. and mirror neutron stars

On the origin of  
antimatter in  
cosmos


Zurab Berezhiani

Summary

NS-NS merger and kilonova (GW170817 ?)  
r-processes can give heavy \*trans-Iron\* elements

Mirror NS-NS merger is invisible (GW190425 ?  $M_{\text{tot}} = 3.4M_{\odot}$  )

But not completely ... if during the evolution they developed small  
core of our **antimatter** (depends on the mirror BA sign)  
– their mergers can be origin of antinuclei for AMS-2





# Looking for antimatter stars/planets

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

DUPOURQUÉ, TIBALDO, and VON BALLMOOS

PHYS. REV. D **103**, 083016 (2021)




FIG. 1. Positions and energy flux in the 100 MeV–100 GeV range of antistar candidates selected in 4FGL-DR2. Galactic coordinates. The background image shows the *Fermi* 5-year all-sky photon counts above 1 GeV (image credit: NASA/DOE/Fermi LAT Collaboration).

$$\text{Antimatter production rate: } \dot{N}_{\bar{b}} = \frac{N_0}{\tau_{\epsilon}} \simeq \epsilon_{15}^2 \left( \frac{M}{M_{\odot}} \right)^{2/3} \times 3 \cdot 10^{34} \text{ s}^{-1}$$

$$\text{ISM accretion rate: } \dot{N}_b \simeq \frac{(2GM)^2 n_{\text{is}}}{v^3} \simeq \frac{10^{32}}{v_{100}^3} \times \left( \frac{n_{\text{is}}}{1/\text{cm}^3} \right) \left( \frac{M}{M_{\odot}} \right)^2 \text{ s}^{-1}$$

Annihilation  $\gamma$ -flux from the mirror NS as seen at the Earth:

$$J \simeq \frac{10^{-12}}{v_{100}^3} \left( \frac{n_{\text{is}}}{1/\text{cm}^3} \right) \left( \frac{M}{1.5 M_{\odot}} \right)^2 \left( \frac{50 \text{ pc}}{d} \right)^2 \frac{\text{erg}}{\text{cm}^2 \text{ s}} \quad d - \text{distance to source}$$



# Getting Energy from Dark Parallel World

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary


I argued that in O and M worlds baryon asymmetries can have same signs:  $B > 0$  and  $B' > 0$ . Since  $B - B'$  is conserved, our neutrons have transition  $n \rightarrow \bar{n}'$  (which is the antiparticle for M observer)

while  $n'$  (of M matter) oscillates  $n' \rightarrow \bar{n}$  into our antineutron

Neutrons can be transformed into antineutrons, but (happily) with low efficiency:  $\tau_{n\bar{n}} > 10^8$  s

dark neutrons, before they decay, can be effectively transformed into our antineutrons in controllable way, by tuning vacuum and magnetic fields, if  $\tau_{n\bar{n}'} < 10^3$  s

$E = 2m_n c^2 = 3 \times 10^{-3}$  erg  
per every  $\bar{n}$  annihilation



Two civilisations can agree to built scientific reactors and exchange neutrons ... we could get plenty of energy out of dark matter !

E.g. mirror source with  $3 \times 10^{17}$  n/s (PSI)  $\longrightarrow$  power = 100 MW




# Asimov Machine: the "Pump"

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary



**First Part:** Against Stupidity ...

**Second Part:** ...The Gods Themselves ...

**Third Part:** ... Contend in Vain?

*"Mit der Dummheit kämpfen Götter  
selbst vergebens!"* – Schiller

Radiochemist Hallam constructs the "Pump": a cheap, clean, and apparently endless source of energy functioning by the matter exchange between our universe and a parallel universe .... His "discovery" was inspired by beings of parallel (mirror) world where stars were very old and so too cold – they had no more energy resources and were facing full extinction ...





# Backup

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

## Some auxiliary slides



# Neutron – mirror neutron oscillation probability

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary


$$H = \begin{pmatrix} m + \mu \vec{B} \vec{\sigma} + V & \epsilon \\ \epsilon & m + \mu \vec{B}' \vec{\sigma} + V' \end{pmatrix}$$

The probability of n-n' transition depends on the relative orientation of magnetic and mirror-magnetic fields. The latter can exist if mirror matter is captured by the Earth

$$P_B(t) = p_B(t) + d_B(t) \cdot \cos \beta$$

$$p(t) = \frac{\sin^2 [(\omega - \omega')t]}{2\tau^2(\omega - \omega')^2} + \frac{\sin^2 [(\omega + \omega')t]}{2\tau^2(\omega + \omega')^2}$$

$$d(t) = \frac{\sin^2 [(\omega - \omega')t]}{2\tau^2(\omega - \omega')^2} - \frac{\sin^2 [(\omega + \omega')t]}{2\tau^2(\omega + \omega')^2}$$



where  $\omega = \frac{1}{2}|\mu B|$  and  $\omega' = \frac{1}{2}|\mu B'|$  ;  $\tau$  - oscillation time

$$A_B^{\text{det}}(t) = \frac{N_{-B}(t) - N_B(t)}{N_{-B}(t) + N_B(t)} = N_{\text{collis}} d_B(t) \cdot \cos \beta \leftarrow \text{assymetry}$$



# Earth mirror magnetic field via the electron drag mechanism

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary



Earth can accumulate some, even tiny amount of mirror matter due to Rutherford-like scattering of mirror matter due to photon-mirror photon kinetic mixing.

High temperature of the Earth core  $\rightarrow$  mirror gas is partially ionized.

Rotation of the Earth drags mirror electrons but cannot move as well mirror ions which are much heavier. So circular electric currents can emerge which seed the mirror magnetic field. Rather tiny amount of captured mirror matter (say  $\sim 10^{45}$  particles) would suffice

These seeds can be strongly enhanced by the dynamo mechanism: mirror plasma captured in the Earth must differentially rotate and also have convective motions




# Experiments

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

By now 8 experiments were done at ILL/PSI



Several new experiments are underway at PSI, ILL and ORNL  
and are projected at ESS




# 2009 – magnetic field vertical

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

Experiment sequence:  $\{B_-, B_+, B_+, B_-, B_+, B_-, B_-, B_+\}$  ,  
 $B \simeq 0.2G$



Careful analysis has shown the non-zero effect: **Z.B. and Nesti, 2012**

$$A(B) = (7.0 \pm 1.3) \times 10^{-4} \quad \chi^2_{dof} = 0.9 \longrightarrow 5.2\sigma$$

Modulation with the period  $T = 24$  hour  $\longrightarrow 5.5\sigma$




# 2009 – magnetic field Horizontal

large field  $B_{\pm} = 0.2$  G and small field  $b_{\pm} < 10^{-2}$  G

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary









# 2009 – magnetic field Horizontal

large field  $B = 0.2$  G and small field  $b < 10^{-2}$  G

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary



small field:  $A_b \simeq 0$ , but large field measurements show non-zero  $A_B$   
and  $E_B$ , both with the period  $T \simeq 300$  hours  
(Unpublished and not included in Fig. of exp. limits)



# Can neutron be transformed into antineutron ... effectively?

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

Small Majorana mass of neutron  $\frac{\epsilon}{2} (n^T C n + \bar{n} C \bar{n}^T) = \frac{\epsilon}{2} (\bar{n}_c n + \bar{n} n_c)$   
 $\equiv n - \bar{n}$  oscillation ( $\Delta B = 2$ )

Oscillation probability for free flight time  $t$

$$P_{n\bar{n}}(t) = (\epsilon t)^2 = (t/\tau_{n\bar{n}})^2 \quad \text{in quasi-free regime} \quad \omega_B t < 1$$

Present bounds on oscillation time  $\tau_{n\bar{n}} = \epsilon^{-1}$  are severe:

$$\tau_{n\bar{n}} > 0.86 \times 10^8 \text{ s} \quad \text{direct limit (free } n) \quad \text{ILL, 1994}$$

$$\tau_{n\bar{n}} > 2.7 \times 10^8 \text{ s} \quad \text{nuclear stability (bound } n) \quad \text{SK, 2020 (this conf.)}$$

$$P_{n\bar{n}}(t) = \frac{t^2}{\tau_{n\bar{n}}^2} = \left( \frac{10^8 \text{ s}}{\tau_{n\bar{n}}} \right)^2 \left( \frac{t}{0.1 \text{ s}} \right)^2 \times 10^{-18}$$

Shortcut through mirror world:  $n \rightarrow n' \rightarrow \bar{n}$ :

Experimental search to be tuned against (dark) environmental conditions

$$P_{n\bar{n}}(t) = P_{nn'}(t) P_{n\bar{n}'}(t) = \frac{t^4}{\tau_{nn'}^2 \tau_{n\bar{n}'}^2} = \left( \frac{1 \text{ s}^2}{\tau_{nn'} \tau_{n\bar{n}'}} \right)^2 \left( \frac{t}{0.1 \text{ s}} \right)^4 \times 10^{-4}$$

No danger for nuclear stability !

Nor for Neutron Stars





$$2 \times 2 = 4 !$$

Z.B., Eur.Phys.J C81:33 (2021), arXiv:2002.05609

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

4 states:  $n, \bar{n} : n', \bar{n}'$  and mixing combinations:

$$n \longleftrightarrow \bar{n} \quad (\Delta B = 2) \quad \& \quad n' \longleftrightarrow \bar{n}' \quad (\Delta B' = 2)$$

$$n \longleftrightarrow n' \quad + \quad \bar{n}' \longleftrightarrow \bar{n} \quad \Delta(B - B') = 0$$

$$n \longleftrightarrow \bar{n}' \quad + \quad n' \longleftrightarrow \bar{n} \quad \Delta(B + B') = 0$$

Full Hamiltonian is  $8 \times 8$ :

$$\begin{pmatrix} m_n + \mu \vec{B} \vec{\sigma} & \epsilon_{n\bar{n}} & \epsilon_{nn'} & \epsilon_{n\bar{n}'} \\ \epsilon_{n\bar{n}} & m_n - \mu \vec{B} \vec{\sigma} & \epsilon_{n\bar{n}'} & \epsilon_{nn'} \\ \epsilon_{nn'} & \epsilon_{n\bar{n}'} & m'_n + V'_n + \mu' \vec{B}' \vec{\sigma} & \epsilon_{n\bar{n}} \\ \epsilon_{n\bar{n}'} & \epsilon_{nn'} & \epsilon_{n\bar{n}} & m'_n + V'_n - \mu' \vec{B}' \vec{\sigma} \end{pmatrix}$$

Present bounds on oscillation time  $\tau_{n\bar{n}} = \epsilon^{-1}$ :

$$\tau_{n\bar{n}} > 0.86 \times 10^8 \text{ s} \quad (\text{free } n), \quad \tau_{n\bar{n}} > 4.7 \times 10^8 \text{ s} \quad (\text{bound } n)$$

$$P_{n\bar{n}}(t) = \frac{t^2}{\tau_{n\bar{n}}^2} = \left( \frac{10^8 \text{ s}}{\tau_{n\bar{n}}} \right)^2 \left( \frac{t}{0.1 \text{ s}} \right)^2 \times 10^{-18}$$



# Shortcut for $n \rightarrow \bar{n}$ via $n \rightarrow n' \rightarrow \bar{n}$

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

Consider case when direct  $n - \bar{n}$  mixing simply absent:  $\epsilon_{n\bar{n}} = 0$

Anyway,  $n \rightarrow \bar{n}$  emerges as second order effect via  $n \rightarrow n' \bar{n}' \rightarrow \bar{n}$

$$\bar{P}_{n\bar{n}} = \bar{P}_{nn'} \bar{P}_{n\bar{n}'}$$

$$\bar{P}_{nn'} = \frac{2\epsilon_{nn'}^2 \cos^2(\beta/2)}{(\Omega - \Omega')^2} + \frac{2\epsilon_{nn'}^2 \sin^2(\beta/2)}{(\Omega + \Omega')^2}, \quad \bar{P}_{n\bar{n}'} = \frac{2\epsilon_{n\bar{n}'}^2 \sin^2(\beta/2)}{(\Omega - \Omega')^2} + \frac{2\epsilon_{n\bar{n}'}^2 \cos^2(\beta/2)}{(\Omega + \Omega')^2}$$

where  $\beta$  is the (unknown) angle between the vectors  $\vec{B}$  and  $\vec{B}'$

Disappearance experiments measure the sum  $P_{nn'} + P_{n\bar{n}'} \propto \epsilon_{nn'}^2 + \epsilon_{n\bar{n}'}^2$

$n - \bar{n}$  transition measures the product  $P_{n\bar{n}} = P_{nn'} P_{n\bar{n}'} \propto \epsilon_{nn'}^2 \epsilon_{n\bar{n}'}^2$

From the ILL'94 limit  $P_{n\bar{n}} < 10^{-18}$  (measured at  $B = 0$ ) we get

$$\tau_{nn'} \tau_{n\bar{n}'} > \frac{2 \times 10^9}{\Omega'^2} \approx \left( \frac{0.5 \text{ G}}{B'} \right)^2 \times 100 \text{ s}^2$$

E.g.  $\tau_{nn'} \tau_{n\bar{n}'} \sim 1$  second is possible if  $B' \sim 5 \text{ G}$

Limits become even weaker if  $\Delta m > 0.1 \text{ neV}$



# How good the shortcut can be?

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

Assuming e.g.  $\tau_{nn'} \tau_{n\bar{n}'} = 100$  s and  $B' = 0.5$  G, we see that

ILL94-like measurement at  $B = 0.45$  G (or  $B = 0.49$  G) would give

$$P_{n\bar{n}} \simeq \sin^2 \beta \times 10^{-15} \quad (\text{or } P_{n\bar{n}} \simeq \sin^2 \beta \times 10^{-12})$$

To maximize  $n - \bar{n}$  probability, one has to match resonance with about 1 mG precision: we get

$$P_{nn'}(t) = \left(\frac{t}{\tau_{nn'}}\right)^2 \cos^2 \frac{\beta}{2}, \quad P_{n\bar{n}'}(t) = \left(\frac{t}{\tau_{n\bar{n}'}}\right)^2 \sin^2 \frac{\beta}{2}$$

and

$$P_{n\bar{n}}(t) = P_{nn'}(t)P_{n\bar{n}'}(t) = \frac{\sin^2 \beta}{4} \left(\frac{t}{0.1 \text{ s}}\right)^4 \left(\frac{100 \text{ s}^2}{\tau_{nn'} \tau_{n\bar{n}'}}\right)^2 \times 10^{-8}$$

**Practically no limit from nuclear stability**

E.g.  $^{16}\text{O}$  decay time predicted  $\sim 10^{60}$  yr vs. present limit  $\sim 10^{32}$  yr !




# How effective $n \rightarrow \bar{n}$ can be?

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

simulations for  $n - \bar{n}$  experiment with  
 $t = 0.1$  s ( $\ell = 100$  m as ILL) and  $t = 0.02$  s ( $\ell = 20$  m)



– and perhaps a chance for free energy ?



# Majorana Machine

On the origin of  
antimatter in  
cosmos


Zurab Berezhiani

Summary



Che cretini! Hanno scoperto il protone neutro e non se ne accorgono!

La fisica è su una strada sbagliata. Siamo tutti su una strada sbagliata...



La fantomatica macchina forse teorizzata da Ettore Majorana! Nella sua formulazione attuale violerebbe un'infinità di principi scientifici, producendo enormi quantità di energia a costo zero. Non può affatto esistere ...



# References

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

Z.B., “Neutron lifetime puzzle and neutron–mirror neutron oscillation,” Eur.Phys.J. C **79**, 484 (2019), arXiv:1807.07906

Z.B., Biondi, Mannarelli and Tonelli, “Neutron-mirror neutron mixing and neutron stars,” Eur.Phys.J. C **81**, 1036 (2021), arXiv:2012.15233

Z.B., “Antistars or antimatter cores in mirror neutron stars?,” Universe **8** (2022) 6, 313, arXiv:2106.11203

Z.B., “A possible shortcut for neutron–antineutron oscillation through mirror world,” Eur.Phys.J. C **81**, 33 (2021), arXiv:2002.05609

Oldies ... but Goldies – Psychedelia or Serendipities ?

Z.B. and Bento, “Neutron - mirror neutron oscillations: How fast might they be?,” Phys.Rev.Lett. **96**, 081801 (2006), hep-ph/0507031

Z.B. “More about neutron - mirror neutron oscillations,” Eur. Phys. J. C **64**, 421 (2009), arXiv:0804.2088

Z.B., “Mirror world and its cosmological consequences,” Int.J.Mod.Phys. A **19**, 3775-3806 (2004), hep-ph/0312335

Z.B., Comelli and Villante, “Early mirror universe: Inflation, baryogenesis, nucleosynthesis and dark matter,” Phys.Lett. B **503**, 362 (2001), hep-ph/0008105



# The Universe: Anthropic or Intelligent Design?

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

Is the Universe Anthropic?      multiverse...

or Anthropomorphic?      with basic instincts to survive

or Anthrophilic?      has sapience and purposes ...

Conspiracy in the fine selection of the SM parameters:

$M_W$ ,  $\Lambda_{\text{QCD}}$ , Yukawa constants       $\theta_{\text{QCD}}?$

and Cosmological term? (Weinberg's anthropic argument ...)

E.g. Neutron-proton-electron mass conspiracy:  $m_e < m_n - m_p$   
– neutron decays if free but is stable in nuclei with  $E_b \sim \text{few MeV}$

Taken Standard Model with all coupling constants fixed in UV,  
sort of "explanation" why  $M_W \sim 10^2 \text{ GeV}$

$M_W < 10 \text{ GeV} \longrightarrow m_e > m_n - m_p$       hydrogen atom decays  $pe \rightarrow n\nu$

$M_W > 10^3 \text{ GeV} \longrightarrow m_n > m_p + m_e + E_b$       only hydrogen, no nuclei

– in either case, no life!      And noone can ask stupid questions  
"Why?"



# Anthropic limit on $n - \bar{n}$ mixing

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

Nuclear instability against

$(A, Z) \rightarrow (A - 1, \bar{n}, Z) \rightarrow (A - 2, Z/Z - 1) + \pi$ 's scales as

Scale of new physics unknown – but  $\tau_{\text{nucl}} \propto \epsilon^2 \propto 1/M^{10}$  ( $\epsilon \propto 1/M^5$ )

Present limit  $\tau_{\text{nucl}} > 10^{32}$  yr implies

$\epsilon < 2.5 \times 10^{-24}$  eV  $\rightarrow M > 500$  TeV or so

$M \rightarrow M/3$  (just 3 times less) would give  $\tau_{\text{nucl}} \rightarrow \tau_{\text{nucl}}/3^{10} \approx 10^{27}$  yr

$\bar{n}n$  ( $\bar{n}p$ ) annihilation releases energy  $E_{\text{ann}} = 2m_n c^2 \approx 3 \times 10^{-10}$  J

Then the Earth power =  $E_{\text{ann}} N_{\oplus} / \tau_{\text{nucl}} \simeq 10$  TW

.. the Earth radioactivity turns dangerous for the Life!

And (happily) the neutron is not elementary particle

– in which case it could have unsuppressed Majorana mass  $\epsilon n^T C n$

It is composite  $n = (udd)$  of three quarks – Majorana mass

can be induced only by D=9 operator  $\frac{1}{M^5} (udd)^2$

Life can exist because of the (intelligent) structure of the SM





# Digression: Anthropic $\theta$ -term in QCD

Z.B., EPJ C 76, 705 (2016), arXiv:1507.05478



On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

QCD forms quark condensate  $\langle \bar{q}q \rangle \sim \Lambda_{\text{QCD}}^3$  breaking chiral symmetry (and probably 4-quark condensates  $\langle \bar{q}q\bar{q}q \rangle$  not reducible to  $\langle \bar{q}q \rangle^2$ )

Can six-quark condensates  $\langle qqqqqq \rangle$  be formed? (i.e. 3 diquarks)  $\langle (udd)^2 \rangle$  or  $\langle (uds)^2 \rangle$  inducing  $n - \bar{n}$ ,  $\Lambda - \bar{\Lambda}$  mixings ( $\Delta B = 2$ )



Vafa-Witten theorem: QCD cannot break vector symmetries ...

.. the prove relies on the absence of  $\theta$ -term (valid strictly for  $\theta = 0$ )

Imagine then world with  $\theta \sim 1$  where  $\langle qqqqqq \rangle \sim \Lambda_{\text{QCD}}^9$

– bad for Life: enormous  $n - \bar{n}$  or  $\Lambda - \bar{\Lambda}$  ...

Let us assume  $\langle qqqqqq \rangle_\theta \sim F(\theta) \Lambda_{\text{QCD}}^9$  with  $F(\theta)$  being a smooth periodic and even function:  $F(\theta) \simeq \theta^2 + \dots$

Then for  $\theta \sim 10^{-10}$ ,  $\langle qqqqqq \rangle_\theta = \theta^2 \Lambda_{\text{QCD}}^9 \sim (1 \text{ MeV})^9$

Then  $\epsilon \sim \theta^2 \Lambda_{\text{QCD}}^9 / m^8 \rightarrow \theta < 10^{-11}$  or so



# Neutron Stars Evolution to mixed star



On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

$$\tau_{\epsilon} = (10^{-15} \text{ eV}/\epsilon)^2 \times 10^{15} \text{ yr} \quad \text{Two regimes are allowed :}$$

1. slow transformation ( $\tau_{\epsilon} \gg 14 \text{ Gyr}$  age of universe)  
then limit from pulsar heating tells  $\tau_{\epsilon} > 10^{15} \text{ yr} \rightarrow \epsilon < 10^{-15} \text{ eV}$  or so  
matches exp. limits for exactly degenerate  $n - n'$
2. fast transformation  $\tau_{\epsilon} < 10^5 \text{ yr}$  or so  $\rightarrow \epsilon > 10^{-10} \text{ eV}$  or so  
– then old pulsars all should be transformed into maximally mixed stars  
matches explanation of neutron lifetime anomaly, non-degenerate  $n - n'$







# Neutron Stars: mass distribution

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary






# Neutron Stars: observational $M - R$

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary




**Figure 4**

The combined constraints at the 68% confidence level over the neutron star mass and radius obtained from (Left) all neutron stars in low-mass X-ray binaries during quiescence (Right) all neutron stars with thermonuclear bursts. The light grey lines show mass-relations corresponding to a few representative equations of state (see Section 4.1 and Fig. 7 for detailed descriptions.)



# Back to trap-beam problem: $\tau_n$ vs. $\beta$ -asymmetry

Updated Fig.7 from Belfatto, Beradze and Z.B, EPJ C 80, 149 (2020)



$$g_A = 1.27625(50)$$

$$\tau_{\text{beam}} = 888.0 \pm 2.0 \text{ s}$$

$$\tau_{\text{trap}} = 878.5 \pm 0.5 \text{ s}$$

Free neutron decay:

$$G_V^2 = \frac{K / \ln 2}{\mathcal{F}_n \tau_n (1 + 3g_A^2)(1 + \Delta_R)}$$

$0^+ - 0^+$  decays:

$$G_V^2 = \frac{K}{2\mathcal{F}t(1 + \Delta_R)}$$

$$\tau_n = \frac{2\mathcal{F}t}{\mathcal{F}_n(1 + 3g_A^2)} = \frac{5172.1(1.1 \rightarrow 2.8)}{1 + 3g_A^2} \text{ s} \quad \text{Czarnecki et al. 2018}$$

$G_V$  and  $\Delta_R$  cancel out even in BSM  $G_V \neq G_F|V_{ud}|$  :  $g_A = -G_A/G_V$



$$g_A = 1.27625(50) \quad \longrightarrow \quad \tau_n^{\text{theor}} = 878.7 \pm (0.6 \rightarrow 1.5) \text{ s} \quad \approx \tau_{\text{trap}}$$



On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary



$$\tau_n^{\text{theor}} = 878.7 \pm 1.5 \text{ s}$$

$$\tau_{\text{trap}} = 878.5 \pm 0.5 \text{ s} \quad (\text{compatible})$$

$$\tau_{\text{beam}} = 888.0 \pm 2.0 \text{ s} \quad (4.5\sigma)$$

$$\tau_{\text{mat}} = 880.1 \pm 0.7 \text{ s} \quad \tau_{\text{magn}} = 877.8 \pm 0.3 \text{ s} \quad (3.3\sigma \text{ discrepancy})$$

So experimentally we have  $\tau_{\text{magn}} < \tau_{n \rightarrow p}^{\text{theor}} < \tau_{\text{mat}} < \tau_{\text{beam}}$

which is possible in  $n - n'$  oscillation scenario So far so Good!



# Dark matter Factory ?


On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary

If my hypothesis is correct, a simple solenoid (magn. field  $\sim$  Tesla) can be an effective machine transforming neutrons into DM neutrons

With good adiabatic conditions 50 % transformation can be achieved



$$P_{nn'}^{\text{tr}} \approx \frac{\pi}{4} \xi \simeq 10^{-2} \left( \frac{2 \text{ km/s}}{v} \right) \left( \frac{P_{nn'}^0}{10^{-6}} \right) \left( \frac{B_{\text{res}}}{1 \text{ T}} \right) \left( \frac{R_{\text{res}}}{10 \text{ cm}} \right)$$

ORNL experiment via  $n \rightarrow n' \rightarrow n$  in strong magn. fields




# Cabibbo Angle Anomaly

On the origin of  
antimatter in  
cosmos

Zurab Berezhiani

Summary



If CKM unitarity is assumed – strong discrepancy between

A:  $|V_{us}| = \sin \theta_C$

B:  $|V_{us}/V_{ud}| = \tan \theta_C$

C:  $|V_{ud}| = \cos \theta_C$

Unitarity excluded at  $> 3\sigma$