



Contribution ID: 1

Type: not specified

Report on the AGATA@GANIL experiment E793S

The region around the magic numbers $N = 28$ and $Z = 20$ is of great interest in nuclear structure physics. Moving away from the doubly-magic isotope ^{48}Ca , in the neutron-rich direction there is evidence of an emergent shell gap at $N = 34$ [1], and in the proton-deficient direction, the onset of shape deformation suggests a weakening of the $N = 28$ magic number [2]. The $^{47}\text{K}(d,p)^{48}\text{K}$ reaction is uniquely suited to investigating this region, as the ground state configuration of ^{47}K has an exotic proton structure, with an odd proton in the $\pi(1s_{1/2})$ orbital, below a fully occupied $\pi(0d_{3/2})$ orbital [3]. As such, the selective neutron transfer reaction (d,p) will preferentially populate states in ^{48}K arising from $\pi(1s_{1/2}) \otimes \nu(fp)$ cross-shell interactions. The implications of this extend both down the proton-deficient $N = 28$ isotonic chain, where these interactions are expected to dominate the structure of the exotic, short-lived ^{44}P nucleus [4], and across the neutron-rich region, where the relative energies of the $\nu(fp)$ orbitals is the driving force behind shell evolution.

The first experimental study of states arising from the interaction between $\pi(1s_{1/2})$ and the orbitals $\nu(1p_{3/2})$, $\nu(1p_{1/2})$ and $\nu(0f_{5/2})$ has been conducted, by way of the $^{47}\text{K}(d,p)$ reaction in inverse kinematics. A beam of radioactive ^{47}K ions was delivered by the GANIL-SPIRAL1+ facility, with a beam energy of 7.7 MeV/nucleon. This beam was estimated to be $> 99.99\%$ pure, with a typical intensity of 5×10^5 pps, and was impinged upon a 0.13 mg/cm^2 CD_2 target. The MUGAST+AGATA+VAMOS detection setup [5] allowed for triple coincidence gating, providing a great amount of selectivity. An analysis based both on excitation and gamma-ray energy measurements has revealed a number of previously unobserved states, and preliminary differential cross sections for the most strongly populated of these states will be presented.

[1] D. Steppenbeck *et al.*, Nature **502**, 207 (2013).

[2] O. Sorlin and M.-G. Porquet, Prog. Part. Nucl. Phys. **61**, 602 (2008).

[3] J. Papuga *et al.*, Phys. Rev. C, **90**, 034321 (2014).

[4] L. Gaudefroy, Phys. Rev. C **81**, 064329 (2010).

[5] M. Assié *et al.*, Nucl. Instrum. Methods A **1014**, 165743 (2021).

Primary authors: PAXMAN, Charlie (University of Surrey); MATTA, Adrien (LPC Caen CNRS/IN2P3); CATFORD, Wilton (University of Surrey)

Presenter: PAXMAN, Charlie (University of Surrey)