



Contribution ID: 47

Type: not specified

## Octupole Collectivity: Coulomb Excitation of $^{224}\text{Ra}$ at ISOLDE-CERN

*Tuesday, June 28, 2011 5:55 PM (10 minutes)*

There is considerable theoretical and experimental evidence that atomic nuclei can assume reflection asymmetric shapes that arise from the octupole degree of freedom. From a microscopic point of view, the wave functions of low-lying 3- octupole excitations must contain components which include the intruding unique parity state  $(l, j)$ . Because of the nature of the octupole-octupole interaction in nuclei, strong octupole correlations arise when the Fermi level lies between this intruder subshell and a subshell with  $\Delta j, \Delta l = 3$ , giving rise to  $[l, j; l-3, j-3]$  particle-hole configurations at relatively low excitation energies. The strongest correlations occur near the proton numbers  $Z = 34, 56$  and  $88$  and the neutron numbers  $N = 34, 56, 88$  and  $134$  where, for the heaviest nuclei, an octupole deformation can occur in the ground state. Indeed, at these values of  $Z$  and  $N$ , nuclei exhibit phenomena associated with reflection asymmetry such as odd-even staggering of the positive- and negative-parity yrast bands in even-even nuclei, parity doublets in odd mass nuclei, and enhanced  $E 1$  moments due to a division of the centre of charge and centre of mass. The only observable that provides unambiguous and direct evidence for enhanced octupole correlations in nuclei is the  $E 3$  matrix element, and the measure of octupole correlations in the ground state is the  $B(E 3; 0^+ \rightarrow 3^-)$ . In the mass region where octupole correlations are expected to be largest, i.e. at  $Z = 88$  and  $N = 134$ , there is a lack of spectroscopic data on  $E 3$  moments. So far, only for  $^{226}\text{Ra}$ , with its comparatively long half life of 1600 years, has it been possible to measure the  $B(E 3)$  strength using Coulomb excitation. This talk will present the current status and the first results from the recent Coulomb excitation the post-accelerated  $^{224}\text{Ra}$  beam at REX-ISOLDE facility, CERN, using the MINIBALL setup.

**Primary author:** GAFFNEY, Liam P. (University of Liverpool)

**Co-authors:** PETER A., Butler (University of Liverpool); MARCUS, Scheck (University of Liverpool)

**Presenter:** GAFFNEY, Liam P. (University of Liverpool)

**Session Classification:** Contributions