



Contribution ID: 14

Type: **not specified**

## Gamma-ray spectroscopy experiments at RIKEN

*Tuesday, June 28, 2011 3:25 PM (25 minutes)*

In the Radioactive Isotope Beam Factory (RIBF) stable primary beams of energies up to 345 MeV/nucleon are used to produce radioactive isotope beams via in-flight separation with the BigRIPS fragment separator.

For gamma-ray spectroscopy experiments these radioactive beams are either incident on a secondary target for in-beam experiments or stopped to measure isomeric or beta-delayed gamma-emission.

In my presentation I will report on the results obtained from the first gamma-ray spectroscopy experiments performed at the RIBF. For stopped-beam experiments neutron-rich nuclei in the mass region  $A=110$  were investigated after fission of a  $^{238}\text{U}$  primary beam. In this mass region, shape transitions around the Zr isotopes were studied. The first in-beam gamma-ray experiments targeted the "Island of Inversion", a region in which the standard ordering of shells is disturbed by neutron intruder configuration across the  $N=20$  shell gap.

Besides showing the results from these first experiments an outlook on future gamma-ray spectroscopy campaigns at the RIBF will be given.

**Primary author:** DOORNENBAL, Pieter (RIKEN)

**Presenter:** DOORNENBAL, Pieter (RIKEN)

**Session Classification:** New Gamma-arrays