

Study of $B^+ \rightarrow K^+ \tau^+ \tau^-$ using hadronic tagging


Debjit Ghosh

August 5, 2022

Study of $B^+ \rightarrow K^+ \tau^+ \tau^-$

Motivation:

1. FCNC: highly suppressed in SM, $\mathcal{O}(10^{-7})$
2. 3rd generation strongly couples to NP




Earlier searches:

1. Attempt in Belle (by Simon Wehle, 2016): Belle Note- 1394
upper limit at 90 % confidence level, $\mathcal{B}(B^+ \rightarrow K^+ \tau^+ \tau^-) < 3.17 \times 10^{-4}$
2. BaBar (2017): [arXiv:1605.09637](https://arxiv.org/abs/1605.09637)
upper limit at 90 % confidence level, $\mathcal{B}(B^+ \rightarrow K^+ \tau^+ \tau^-) < 2.25 \times 10^{-3}$

Initial step: Perform similar to Simon's study in Belle II
modify Vidya's reconstruction script for Belle II

SignalMC generator



#simulated sample size: 50 million


generator model: BTOSLLBALL

release-06-00-10

globalTag: mc_production_MC15ri_a

bkg: early phase III (release-06-00-05), BGx1

Reconstruction



B_{sig}^+ is composed of K^+ , h^+ , and h'^- :

1. $K^+ e^+ e^-$
2. $K^+ e^+ \mu^-$
3. $K^+ e^- \mu^+$
4. $K^+ e^+ \pi^-$
5. $K^+ e^- \pi^+$
6. $K^+ \mu^+ \mu^-$
7. $K^+ \mu^+ \pi^-$
8. $K^+ \mu^- \pi^+$
9. $K^+ \pi^+ \pi^-$

Selection

Charged tracks (e, μ, K, π) cuts:

- transverse distance from IP, $dr < 0.5$
- distance in beam direction from IP, $|dz| < 2$
- polar angle is within CDC acceptance (`thetaInCDCAcceptance`)

PID cuts:

- e : $\mathcal{L}(\text{electronID}) > 0.9$
- μ : $\mathcal{L}(\text{muonID}) > 0.9$
- K : $\mathcal{L}(\text{kaonID}) > 0.6$
- π : $\mathcal{L}(\text{pionID}) > 0.6$

photon: use all photons (`gamma:all`)

Selection

Reconstruct FEI hadronic B_{tag} :

- weight file prefix - 'FEIv4_2021_MC14_release_05_01_12'
- two most probable B_{tag} candidates are accepted
- $M_{bc} > 5.27$
- $|\Delta E| < 0.1$
- FEI signal probability > 0.001
- ROE of B_{tag} has 3 charged tracks

Continuum suppression:

- event sphericity > 0.2
- $\cos TBTO < 0.9$

ROE mask:

- $dr < 0.5$, $|dz| < 2$, thetaInCDCAcceptance
- $E > 0.06$ and $|\text{cluster time}| < 20$

Analysis globalTag: 'analysis_tools_light-2203-zeus'

Reconstruction

- Reconstruct B_{sig}^+ with $K^+h^+h'^-$ combinations
- Reconstruct $\Upsilon(4S) \rightarrow B_{sig}^+ B_{tag}^-$
- Build ROE with **mask** and reconstruct ROE π^0

ROE mask:

- $dr < 0.5$, $|dz| < 2$, $\theta \text{InCDCAcceptance}$
- $E > 0.06$ and $|\text{cluster time}| < 20$


ROE π^0 :

- daughter photons are in ROE and passes ROE mask
- select π^0 with least difference between M and $\text{Inv}M$


..let's look at few distributions

Signal side

M_{bc}




ΔE




"signal" is selected by "isSignalAcceptMissingNeutrino"

Tag side

M_{bc}




ΔE




Extra ECL energy (E_{ECL})

Energy left in ECL cluster after removing B_{tag} and B_{sig} related deposition




Ideally it should peak at zero



subtract energy of ROE π^0 from E_{ECL}


q_K^2 and $M(\tau^+\tau^-)$



Constraint invariant mass of τ pair:

$$q_K^2 \equiv (p_{\Upsilon(4S)} - p_{B_{tag}} - p_K)^2$$

where $p_{\Upsilon(4S)}$ is the four momentum of beam



reconstructed invariant mass of τ pair