

Longitudinal polarization asymmetry of Ξ_b decays with the effect of non-universal Z' boson

giovedì 16 febbraio 2023 19:00 (1O 30m)

Rare baryonic decays induced by flavour changing neutral current (FCNC) have been of immense interest in recent years because of their sensitivities towards new physics (NP) beyond the standard model (SM). The exploration had been triggered with the observation of $\Lambda_b \rightarrow \Lambda \mu^+ \mu^-$ transition at the Fermilab [1] and the LHCb [2]. Theoretically these decays are also studied at different NP models [3-5]. Inspired by these results obtained for baryonic decays [3-7], we are interested to study the polarization asymmetry for Ξ_b baryon with the effect of NP. Various theoretical studies of branching fractions for $\Xi_b \rightarrow \Xi l^+ l^-$ decays in the standard model (SM) [8] proclaim the possibility of observation of these decays at the LHC. In this work, we will mainly concentrate on longitudinal polarization asymmetries for muonic, electronic and taunic channels in family non-universal Z' model [9, 10]. Asymmetry parameters characterize the angular dependence of differential decay width with polarized and unpolarized heavy baryons. We will investigate the observables with the contribution of Z' boson. The phenomenology of Z' is one of the important sectors to the accelerators. Due to its heavy mass, it may be used to calibrate the upcoming runs of the experiments. Here, we will introduce the NP couplings and use their constrained values. We will show the variation of the observables throughout the whole allowed kinematic region. These results for Ξ_b decays will help the experimental community to observe the decays in colliders and will unlock a new horizon to the theoretical community to probe NP with heavy baryons.

References:

- T. Aaltonen et al. [CDF Collaboration], Phys. Rev. Lett. 107, 201802 (2011) [arXiv: 1107.3753 [hep-ph]].
- R. Aaij et al. [LHCb Collaboration], Phys. Lett. B 725, 25 (2013).
- S. Sahoo, C. K. Das and L. Maharana, Int. J. Mod. Phys. A 24, 6223 (2009).
- S. Sahoo and R. Mohanta, New J. Phys. 18, 093051 (2016) [arXiv: 1607.04449 [hep-ph]].
- D. Banerjee and S. Sahoo, Chin. Phys. C 41, 083101 (2017).
- A. Nasrullah, M. J. Aslam and S. Shafaq, Prog. Theor. Exp. Phys. 2018, 043B08 (2018).
- R. L. Workman et al. [Particle data group], Prog. Theor. Exp. Phys. 2022, 083C01 (2022).
- K. Azizi, M. Bayar, A. Ozpineci, Y. Sarac and H. Sundu, Phys. Rev. D 85, 016002 (2012) [arXiv: 1112.5147 [hep-ph]].
- P. Langacker, Rev. Mod. Phys. 81, 1199 (2009).
- P. Maji, S. Mahata, S. Biswas and S. Sahoo, Int. J. Theor. Phys. 61, 162 (2022).

Autori principali: CHAKRABARTY, RAJESH (NATIONAL INSTITUTE OF TECHNOLOGY DURGAPUR); BISWAS, SWAGATA (NATIONAL INSTITUTE OF TECHNOLOGY DURGAPUR)

Relatore: BISWAS, SWAGATA (NATIONAL INSTITUTE OF TECHNOLOGY DURGAPUR)

Classifica Sessioni: Poster Session and Discussion Session