

SIGRAV Winter School 2023 - QSimFP Tutorial

Sam Patrick, Sebastian Erne, Silke Weinfurtner

January 2023

1. The equations of motion for a shallow fluid flow are,

$$\partial_t \phi + \frac{1}{2} \mathbf{v}^2 + gh = 0, \quad \partial_t h + \nabla \cdot (h \mathbf{v}) = 0.$$

where $\mathbf{v} = \nabla \phi$ is a d -dimensional velocity field, $g \simeq 9.81 \text{ms}^{-2}$ is the gravitational acceleration and h is the height of the fluid. By perturbing $\phi \rightarrow \phi + \delta \phi$ and $h \rightarrow h + \delta h$, show that velocity potential fluctuations obey,

$$(\partial_t + \nabla \cdot \mathbf{v})(\partial_t + \mathbf{v} \cdot \nabla) \delta \phi - g \nabla \cdot (h \nabla \delta \phi) = 0. \quad (1)$$

2. Using (1), find a dispersion relation for $\delta \phi$ in a homogeneous condensate, i.e. when \mathbf{v} and n are constant. Give an expression for the wave speed c in a frame co-moving with the fluid.
3. Show that (1) can be written in the form,

$$\partial_\mu (f^{\mu\nu} \partial_\nu \delta \phi) = 0, \quad \partial_\mu = \begin{pmatrix} \partial_t \\ \partial_i \end{pmatrix}, \quad (2)$$

where $i = 1 \dots d$, and find the components of $f^{\mu\nu}$. Hint: it will help to write,

$$f^{\mu\nu} = \begin{pmatrix} f^{tt} & f^{tj} \\ f^{it} & f^{ij} \end{pmatrix}.$$

4. Write the form of the metric tensor $g^{\mu\nu}$ such that $f^{\mu\nu} = \sqrt{-g} g^{\mu\nu}$, where g is the determinant of the metric. Find a condition for which the identification above is valid.

You may find the following facts helpful. A matrix M of the form,

$$M = C \begin{pmatrix} a^2 - b^i b^i & -b^i \\ b^j & \delta^{ij} \end{pmatrix},$$

has inverse and determinant,

$$M^{-1} = \frac{1}{a^2 C} \begin{pmatrix} -1 & -b^j \\ b^i & a^2 \delta^{ij} - b^i b^j \end{pmatrix}, \quad \det(M) = -a^2 C^{d+1},$$

where C is a conformation factor, a is a scalar and b is a $1 \times d$ vector.