



# Initial Physics Performance and Status of the MPD at NICA

Ivonne Maldonado\* for the MPD Collaboration

\*ivonne.alicia.maldonado@gmail.com

The 20th International Conference on Hadron Spectroscopy and Structure (HADRON 2023)

Genova, Italy. June 5th-9th, 2023



### NICA: Unique and complementary







T. Galatyuk, Nucl.Phys. A982(2019); https://github.com/tgalatyuk/interaction\_rate\_facilities

Neutron stars mergers → similar density and temperature achieved by HIC at NICA.



Lukas R. Weih & L. Rezzolla (GUF)/CERN

- At  $\mu_B \sim 0$ , smooth crossover (lattice QCD + data)
- At large  $\mu_B$ , we expect 1st order phase transition  $\rightarrow$  QCD critical point
- Thermal model indicates that highest baryon density is achieved at NICA energy
- Energy range  $\sqrt{s_{NN}} < 6 \text{ GeV} \rightarrow \text{most}$  appropriate to search CEP

BM@N and MPD will study QCD Medium at extreme net baryon densities.

Many ongoing (NA61/Shine, STAR-BES) and future experiments (CBM at FAIR) in the same energy range



# MPD at NICA Complex in Dubna

- 👉 Two injection chains
  - Ion sources  $(A/Z \le 3) \rightarrow \text{LINAC LU-20}$ (5 MeV/u)  $\rightarrow$  Nuclotron
  - ESIS KRION sources  $(A/Z \le 6) \rightarrow$  HILAc (3.24 MeV/u)  $\rightarrow$  Booster
- ✤ SC Booster synchrotron

injection up to  $2\cdot 10^9$  accelerated up to  $\sim 600~{\rm MeV/u}$  ions of  $^{197}Au^{31+}$ 

### ✤ Nuclotron synchrotron

injection up to  $1\cdot 10^9$  ions accelerated up to  $1-4.5~{\rm GeV/n}$ 

→ BM@N

- Two Collider superconducting storage rings
  - MPD
  - SPD



#### Collider. Parameters for 45 T·m, 11 GeV/u for $Au^{79+}$

Ring circumference (m)	503.4	Luminosity (cm $^{-2}$ s $^{-1}$ )	$10^{27}$
Number of bunches	22	RMS bunch length (m)	0.6
$\beta$ (m)	0.35	Energy in CM (GeV)	4 - 11
RMS $\Delta  ho /  ho$ ( $10^{-3}$ )	1.6	IBS growth time (s)	1800



### **Progress of civil construction**







Initial Physics Performance and MPD status

I. Maldonado

### **NICA** status



#### Accelerator chain - Injection System – is constructed & commissioned!





Initial Physics Performance and MPD status

I. Maldonado

### Status of the collider construction



- In June 2022, all the collider dipole magnets were installed and mechanically adjusted in the collider tunnel, connected by pairs to each other.
- The assembly of the collider is postponed until completion of the engineering infrastructure mounting that is expected during this year.





### Multi-Purpose Detector (MPD) Collaboration



11 Countries > 500 participants, 35 institutes and JINR

#### Organization

Acting Spokesperson: Deputy Spokesperson: Institutional Board Chair: Project Manager: Victor Riabov Zebo Tang, Arkadiy Taranenko Alejandro Ayala Slava Golovatyuk

#### Joint Institute for Nuclear Research

A.Alikhanyan National Lab of Armenia, Yerevan, Armenia; University of Ploydiv Bulgaria: Tsinahua University Beilina China: University of Science and Technology of China, Hefei, China: Huzhou University, Huzhou, China; Institute of Nuclear and Applied Physics, CAS, Shanahai, China; Central China Normal University. China: Shandona University, Shandong, China; University of Chinese Academy of Sciences, Beijing, China; University of South China, China: Three Goraes University. China: Institute of Modern Physics of CAS Lanzhou China: Tbilisi State University, Tbilisi, Georgia; Institute of Physics and Technology, Almaty, Kazakhstan Benemerita Universidad Autónoma de Puebla Mexico: Centro de Investigación y de Estudios Avanzados Mexico:

MPD International Collaboration was established in 2018 to construct, commission and operate the detector



05062023



Initial Physics Performance and MPD status

I. Maldonado

7/23

### **MPD** apparatus

#### Stage I: TPC, TOF, ECAL, FFD, FHCal



Beam configuration

- Not-optimal beam optics with wide z-vertex distribution,  $\sigma_z\sim 50{\rm cm}$
- Reduced luminosity(  $\sim 10^{25}$  is the goal for 2023)  $\rightarrow$  collision rate  $\sim 50~{\rm Hz}$
- Collision system available with the current sources: C(A = 12), N(A = 14), Ar(A = 40), Fe(A = 56), Kr(A = 78 86), Xe(A = 124 134), Bi(A = 209)  $\rightarrow$  start with Bi+Bi @ 9.2 GeV





Initial Physics Performance and MPD status I. Maldonado

05.06.2023

Acceptance

 $< 2\pi$  < 1.4

 $< 2\pi$  2.9 - 3.3

 $< 2\pi$  2 - 5

TPC:

FFD:

FHCal:

TOF. EMC:

### MPD subsystem in production









100% of MRPCs(modules) are ready,

Cool down and power the magnet + magnetic field Cosmic tests ongoing.

#### Support structure



made of carbon fiber sagite  $\sim 5$  mm, 0.13  $X_0$ 

#### TPC central tracking detector



#### ECAL (projective geometry)



FHCal



Forward detectors - are in advanced state of production(electronic and integration)

38400 towers 66-83% of the whole detector will be produced for Stage-I

FFD



NICA

Initial Physics Performance and MPD status

I. Maldonado

### **MPD Status and plan**





#### 2023:

preparation for Vacuum test of Solenoid with Cryostat. Solenoid cooling down.

#### 0 2024:

Supplying the current to the solenoid and Correction coils. MF measurements. Support frame and detectors installation. Cabling and Installation of beam pipe.

#### 2025:

Move the MPD on Collider beam line and Commisioning.

#### beyond:

MPD commissioning first run with Bi+Bi @ 9.2 GeV, ~ 50-100 M events for alignment calibration and physics. Au+Au @ 11 GeV, design luminosity system size and collision energy scans

- Preparation of the MPD detector and experimental program is ongoing, all activities are continued
- All components of the MPD 1-st stage detector are in advanced state of production (subsystems, support frame, electronics platforms, LV/HV, control systems, cryogenics, cabling, etc.)

Schedule of the MPD-NICA is significantly affected by the current geopolitical situation (suspension of collaboration with CERN and Polish & Czech Republic member institutions, economic sanctions and problems with supplies of many components from companies).



### **Computing Resources for the MPD**





- Software framework MpdRoot object oriented set of tools to simulate, transport and reconstruct MC events within MPD experiment
  - Centralized Analysis Framework



- Main technological elements at VBLHEP and LIT
- LIT NICA part of MICC → connection with computing complexes of other organizations involved in NICA Complex
- DIRAC  $\rightarrow$  infrastructure enables integration of heterogeneous computing resources at multiple sites

# MPD physics program



#### G. Feofilov, A. Aparin Global Observables

- Total event multiplicity
- Total event energy
- Centrality determination
- Total cross-section measurement
- Event plane measurement at all rapidities
- Spectator measurement

#### V. Kolesnikov, Xianglei Zhu Spectra of light flavor and hypernuclei

- Light flavor spectra
- Hyperons and hypernuclei
- Total particle yields and yield ratios
- Kinematic and chemical properties of the event
- Mapping QCD Phase Diagram

#### K. Mikhailov, A. Taranenko Correlations and Eluctuations

- Collective flow for hadrons
- Vorticity, Λ polarization
- E-by-E fluctuation of multiplicity, momentum and conserved quantities
- Femtoscopy
- Forward-Backward correlations
- Jet-like correlations

#### D. Peresunko, C. Yang Electromagnetic probes

- Electromagnetic calorimeter measurement
- Photons in ECAL and central barrel
- Low mass dilepton spectra in-medium modification of resonances and intermediate mass region

#### Wangmei Zha, A. Zinchenko **Heavy flavor**

- Study of open charm production
- Charmoinium with ECAL and central barrel
- Charmed meson through secondary vertices in ITS and HF electrons
- Explore production at charm threshold



### **MPD** paper





 First collaboration paper recently published in EPJA
 Status and initial physics performance studies of the MPD experiment at NICA, Eur.Phys.J.A 58 (2022) 7, 140



Initial Physics Performance and MPD status

# 

# Centrality and reaction plane determination

#### **Centrality with TPC multiplicity**



- MC Glauber approach to extract b
- MC Glauber compatible with Bayesian inversion method (Γ-fit)
- Similar results with different event generators and energies



• Centrality estimation consistent with STAR  $\rightarrow$  good for cross checks between experiments

Acta Physica Polonica B 14 (2021) 3, 503-506, Particles 4 (2021) 2, 275-287, Phys.Part.Nucl.Lett. 19 (2022) 5,



#### **Centrality with FHCal**

I. Maldonado



Two dimensional linear fit of the energy deposition in FHCal modules, to extract the correlation between the maximum energy and the deposited energy in 10% centrality classes



### Bulk Properties: hadron spectra, yields and ratios

- Particle spectra, yields and ratios are sensitive to bulk fireball properties and phase transformations in the medium
- Uniform acceptance and large phase coverage are crucial for precise mapping of the QCD phase diagram



- MPD provides large phase-space coverage for identified pions and kaons (> 70% of the full phasespace at 9 GeV)
- Hadron spectra can be measured from  $p_T=(0.1)0.2$  to  $2.5~{
  m GeV/c}$  (for  $\pi$ )
- Extrapolation to full  $p_T$ -range and to the full phase space can be performed exploiting the spectra shapes (see BW fits for  $p_T$ -spectra and Gaussian for rapidity distributions)



# **Hyperon Reconstruction**

- \*\* MPD
- Strangeness enhancement is considered as a signature of the QGP formation Rafelski, Phys. Rep. 88(1982)331, Rafelski, Múller, P.R.Lett.
   48(1982)1066
- Experimentally observed in HIC at AGS, SPS, RHIC and LHC energies.
- Differential measurements (vs. pT, multiplicity, event shape, energy balance) of strange baryons are needed in different collision systems (pp, pA, AA) at NICA energies

No consensus on the dominant strangeness enhancement mechanisms:

- strangeness enhancement in QGP contradicts with the observed collision energy dependence
- strangeness suppression in pp within canonical suppression models reproduces most of results except for φ(1020)



Acta Phys. Pol. B Proc. Suppl. 14, 529 (2021)

Strange baryons can be reconstructed with good S/B ratios using charged hadron identification in the TPC and TOF
and different decay topology selections. And different techniques like TMVA. https://doi.org/10.3390/particles6020027



Initial Physics Performance and MPD status

I. Maldonado

# **Global Polarization**





 $\Lambda$  and  $\bar{\Lambda}$  polarization can be measured by its self analyzing charged decay  $\rightarrow$  preferential emission of p is along spin direction.

Nature 548,62-65(2017)





- Global polarization decrease with  $\sqrt{s_{NN}}$
- Transport models used to reproduce global polarization are: AMPT, 3FD, UrQMD+vHLLE



Kosuke Okubo, QM022 - STAR, Phys.Rev.C, 104(6):L061901, 2021

MPD will cover  $\sqrt{s_{NN}} = 4 - 11$  GeV as a function of centrality,  $p_T$  and Y not only for  $\Lambda$  but other hyperons( $\overline{\Lambda}, \Xi, \overline{\Xi}, \Sigma, \overline{\Sigma}$ ).

PHSD simulation of 15M events for Bi+Bi at  $\sqrt{s_{NN}} = 9.2 \text{ GeV}$ Full event/detector simulation and reconstruction First global measurements for  $\Lambda/\bar{\Lambda}$  will be possible with  $\sim 10$ M data sampled events





#### I. Maldonado

### **Reconstruction of resonances**

- MPD
- Resonances are best suited to probe density and lifetime of the late hadronic phase of HI collisions
- Suppression of short-lived resonances was observed in central HI collisions at SPS, RHIC and LHC  $\rightarrow$  dominance of rescattering over regeneration  $\rightarrow$  consistent with existence of a long enough hadronic phase  $\rightarrow$  hadronic phase lifetime  $\sim 10 \text{ fm/c}$
- Hadronic phase affects most of observables measured in the final state (flow, correlations, yields, etc.)
- Measurements for resonances are vital to cross check the hadronic phase models



- MPD is capable of reconstruction the resonance peaks in the invariant mass distributions using combined charged hadron identification with TPC and TOF
- Decays with weakly decaying daughters require additional second vertex and topology cuts for reconstruction
- First measurements for resonances will be possible with accumulation of  $\sim 10~{\rm M}$  of Bi+Bi events

NICA

### **Reconstruction of hypertritons**



19/23

- Hyper nuclei measurement provides information about production mechanism, Y-N potential, strange sector of nuclear EoS
- It has strong implications for astronuclear physics, since are expected in the inner core of neutron stars
- Production mechanism can be described by different phenomenological models: statistical hadronization (SHM) and coalescence
- Models predict enhanced hypernuclear production at NICA, double hypernuclei are reachable



### First measurements for hypertriton will be possible with 50M Bi+Bi @ 9.2 events

Phys.Part.Nucl.Lett. 19 (2022) 1, 46-53, https://doi.org/10.3390/physics5020028



Hypertriton reconstruction allows extraction of its lifetime  $\tau$ 



$$\begin{aligned} (\tau) &= & N(0) \exp(-\frac{\tau}{\tau_0}) \\ &= & N(0) \exp(-\frac{ML}{cp\tau_0}) \end{aligned}$$

 $^{\rm Proper time:\, m}$  where  $\tau$  is the proper time, L is the decay distance, p is the particle momentum, M is the hypertriton rest mass and c is the speed of light



Initial Physics Performance and MPD status

#### I. Maldonado

### Anisotropic flow

Flow has high sensitivity to the transport properties of the QCD matter: EoS, speed of sound ( $c_s$ ), specific viscosity ( $\eta$ /s), etc.

$$\frac{dN}{d\phi} = \frac{N}{2\pi} (1 + \sum 2v_n \cos(n(\phi - \Psi_n)))$$

with  $v_n = \langle cos(n(\phi - \Psi_n)) \rangle$ 

 $v_1$  and  $v_2$  for identified hadrons

UrQMD events for Bi+Bi @ 9.2 GeV measured at mid-centrality 10-40%



MPD detector is able to provide detailed differential measurements of directed and elliptic flows with high accuracy.

# \*\* MPD

#### Higher harmonics $v_3$





Models show that higher harmonic ripples are more sensitive to the existence of a QGP phase

#### Collective flow for VO ( $K_s^0$ and $\Lambda$ )

Results for Au+Au @ 11 GeV (UrQMD), 25M events  $\rightarrow$  full event/detector simulation and reconstruction



Differential flow signal extraction using invariant mass fit method  $% \left( {{{\rm{D}}_{{\rm{s}}}}_{{\rm{s}}}} \right)$ 

# **Dielectron studies**

- HBT measurements for identical particles
- Yield and flow of  $e^+e^-$  pairs:
  - probe deconfinement and chiral symmetry restoration
  - effective temperature



T. Galatyuk et al., Eur. Phys. J. A 52 (2016) 131; R. Rapp and H. v. Hess, PLB 753 (2016) 586 J.Cleymans et al. 2006 Phys. Rev. C73, 034905 NA60; H. Specht, AIP Conf. Proc. 1322 (2010) 160; HADES: Nature Physics 15 (2019) 1040

NA60: H. Specht, AIP Conf. Proc. 1322 (2010) 160; HADES: Nature Physics 15 (2019) 1040

BiBi@9.2 GeV (UrQMD+PHSD), 10 M events  $\rightarrow$  full event/detector simulation and reconstruction



- S/B (integrated in 0.2 -1.5 GeV/c) 5~10%
- Methods to improve S/B ratio with a minimal penalty for pair reconstruction are being developed
- Meaningful measurements for  $e^+\,e^-$  continuum and LVMs would require  $\sim 10^8$  events, first observations will be possible with  $\sim 50$  M events



Initial Physics Performance and MPD status I. Me

I. Maldonado





- MPD collaboration is steadily coming to final integration of the detector and first data taking on the beams from NICA
- Physics program for the first years of MPD data taking is formulated and the first physics paper was recently published
- MPD will provide a unique opportunity for investigating properties of nuclear matter at maximal densities to map the QCD phase diagram, to search for phase transition and the Critical End Point
- First operations of the MPD detector are expected with cosmic studies.
- Comissioning and start of data taking with Bi + Bi collisions at  $\sqrt{s_{NN}} = 9.2$  GeV at the NICA complex is expected at 2025











Initial Physics Performance and MPD status I. Maldonado

