

# The $D_s^+$ decay into $\pi^+ K_S^0 K_S^0$ reaction and the $I=1$ partner of the $f_0(1710)$ state

Wednesday, 7 June 2023 17:10 (20 minutes)

We have identified the decay modes of the  $D_s^+ \rightarrow \pi^+ K^{*+} K^{*-}$ ,  $\pi^+ K^{*0} \bar{K}^{*0}$  reactions producing a pion and two vector mesons. The posterior vector-vector interaction generates two resonances that we associate to the  $f_0(1710)$  and the  $a_0(1710)$  recently claimed, and they decay to the observed  $K^+ K^-$  or  $K_S^0 K_S^0$  pair, leading to the reactions  $D_s^+ \rightarrow \pi^+ K^+ K^-$ ,  $\pi^+ K_S^0 K_S^0$ . The results depend on two parameters related to external and internal emission. We determine a narrow region of the parameters consistent with the large  $N_c$  limit within uncertainties which gives rise to decay widths in agreement with experiment. With this scenario we make predictions for the branching ratio of the  $a_0(1710)$  contribution to the  $D_s^+ \rightarrow \pi^0 K^+ K_S^0$  reaction, finding values within the range of  $(1.3 \pm 0.4) \times 10^{-3}$ . Comparison of these predictions with coming experimental results on that latter reaction will be most useful to deepen our understanding on the nature of these two resonances.

**Primary authors:** DAI, Lianrong; OSET, Eulogio; GENG, Lisheng

**Presenter:** DAI, Lianrong

**Session Classification:** Light meson spectroscopy

**Track Classification:** Light meson spectroscopy