

## Sivers function of sea quarks in the light-cone model

We calculate the Sivers function of  $\bar{u}$  and  $\bar{d}$  quarks using the overlap representation within the light cone formalism. The light-cone wave functions of the proton are obtained in terms of the  $|q\bar{q}B\rangle$  Fock states motivated by the meson-baryon fluctuation model. We consider the final-state interaction at the level of one gluon exchange. In a simplified scenario, the Sivers function of  $\bar{u}$  and  $\bar{d}$  can be expressed as the convolution of the Sivers function of the pion inside the proton and the unpolarized distribution of  $\bar{q}$  inside the pion. The model parameters are fixed by fitting the unpolarized sea quark distributions to the known parameterizations. We present the numerical results for  $f_{1T}^{\bar{u}/P}(x, \boldsymbol{k}_T^2)$  and  $f_{1T}^{\bar{d}/P}(x, \boldsymbol{k}_T^2)$ . The first transverse moment of the sea quark Sivers functions in our model are found to be negative and the magnitude is about 0.004 at most.

**Primary author:** LUAN, Xiaoyan (Southeast University)

**Presenter:** LUAN, Xiaoyan (Southeast University)

**Session Classification:** Exotic hadrons and candidates

**Track Classification:** QCD and hadron structure