ID contributo: 82 Tipo: Contributed

A novel approach for determining spatial moments of the proton charge density

giovedì 8 giugno 2023 18:00 (20 minuti)

The internal structure of the nucleon is a field of intensive study. This structure can be experimentally probed through electron elastic scattering off a proton target. This allows the extraction of the proton electric and the magnetic form factors which characterize the charge and the magnetization densities. The spatial moments of the proton charge density are extracted using the electric form factor (EFF) data. Up to now, methods rely on the evaluation of the EFF derivative in the limit of zero four-momentum transfer Q2 enabling access only to positive even orders of spatial moments. A novel approach [1] based on integral forms of the Fourier transform of the density function allows the determination of spatial moments of densities to any real-valued order. Within this approach, we compute spatial moments of different orders from a reanalysis of EFF data obtained with Rosenbluth separation and from low Q2 experiments covering a range of Q2 from $2 \times 10^{\circ}(-4)$ up to 8.8 GeV2. We pay specific attention to the evaluation of systematic uncertainties. In this context, the evaluation of the proton charge radius corresponding to the second-order moment of the proton charge density will be discussed.

Autore principale: ATOUI, Mariam

Relatore: ATOUI, Mariam

Classifica Sessioni: QCD and hadron structure

Classificazione della track: QCD and hadron structure