## Latest results and precision measurements from the NA62 experiment

Tuesday, 6 June 2023 14:00 (25 minutes)

The NA62 experiment at CERN collected the world's largest dataset of charged kaon decays in 2016-2018, leading to the first measurement of the branching ratio of the ultra-rare K+  $\rightarrow \pi + v v$  decay, based on 20 candidates, and presented in 2021. In this talk the NA62 experiment reports new results from analyses of K+  $\rightarrow \pi$ + $\mu$ + $\mu$ - and K+  $\rightarrow \pi$ + $\gamma\gamma$  decays, using a data sample recorded in 2017–2018. The K+  $\rightarrow \pi$ + $\mu$ + $\mu$ - sample comprises about 27k signal events with negligible background contamination, and the presented analysis results include the most precise determination of the branching ratio and the form factor. The K+  $\rightarrow \pi$ + $\gamma\gamma$  sample contains about 4k signal events with 10% background contamination, and the analysis improves the precision of the branching ratio measurement by a factor of 3 with respect to the previous measurements. The NA62 experiment can also be run as a "beam-dump experiment" by removing the Kaon production target and moving the upstream collimators into a "closed" position. More than 1017 protons on target have been collected in this way during a week-long data-taking campaign by the NA62 experiment. We report on new results from analysis of this data, with a particular emphasis on Dark Photon and Axion-like particle Models.

**Primary authors:** ROMANO, Angela (University of Birmingham); PINZINO, Jacopo (Istituto Nazionale di Fisica Nucleare)

Presenter: PINZINO, Jacopo (Istituto Nazionale di Fisica Nucleare)

Session Classification: Hadrons and physics beyond the standard model

Track Classification: Hadrons and physics beyond the standard model