


The background of the slide features a faded, light blue musical score. It consists of multiple staves of music, likely for a piano or keyboard, with various notes, rests, and phrasing marks. The score is arranged in a grid-like pattern across the slide, with some staves being more prominent than others.

Kaon-nuclei interaction studies at low energies:
the AMADEUS project

Kristian Piscicchia
INFN, Laboratori Nazionali di Frascati
on behalf of the AMADEUS Collaboration

8th Int. Conf. On Nuclear Physics at Storage Rings STORI'11
Laboratori Nazionali di Frascati October 9-14, 2011




**ANTIKAONIC
MATTER
AT
DAΦNE: AN
EXPERIMENT
WITH UNRAVELING
SPECTROSCOPY**

**AMADEUS collaboration
116 scientists from 14 Countries and 34 Institutes**

**Inf.infn.it/esperimenti/siddharta
and**

LNF-07/24(IR) Report on Inf.infn.it web-page (Library)

**AMADEUS started in 2005 and
was presented and discussed in all the
LNF Scientific Committees**



**EU Fundings FP7 – I3HP2:
Network WP9 – LEANNIS;
WP24 (SiPM JRA);
WP28 (GEM JRA)**

CONTENTS

Scientific case

State of the
experimental
search

Experimental
programme

Setup performance
requirements

Experimental setup



KLOE data analysis

AMADEUS scientific case

An important hadron physics unresolved problem:

- how hadron masses and interactions change in nuclear medium

Approach by means of the predicted **kaonic nuclear clusters**



- from which to deduce the **hadron-nucleus potential** and the **in-medium hadron mass**

AMADEUS scientific case

Deeply bound Kaonic nuclear states requires the presence of a strong attractive $\bar{K}N$ interaction in the isospin $I=0$ channel.

The pillars of the existence of narrow \bar{K} -nuclear states are:

- The low energy $\bar{K}N$ scattering data
- The kaonic hydrogen shift and width of the ground state
- The binding energy and decay width of $\Lambda(1405)$ regarded as an isospin

$I = 0$ bound state of $\bar{K} + N$

AMADEUS scientific case

- In presence of strong $\bar{K}N$ attractive potential were firstly suggested by Wycech

(S. Wycech, Nucl. Phys. A450 (1986) 399c)

- Y. Akaishi and T. Yamazaki 'nuclear bound states in light nuclei'

(Phys. Rev. C65 (2002) 044005)



K^- cluster	M (MeV/ c^2)	$-E_B$ (MeV)	Γ_B (MeV)	$\rho(0)$ (fm $^{-3}$)	R_{rms} (fm)
pK^-	1407	27	40	0.59	0.45
ppK^-	2322	48	61	0.52	0.99
$pppK^-$	3211	97	13	1.56	0.81
$ppnK^-$	3192	118	21	1.50	0.72
ppK^-K^-	2747	117	35	-	-
$ppnK^-K^-$	3582	221	37	2.97	0.69

State of the experimental search and theoretical debate for DBKS

- E471, E549, E570 @ KEK
- FINUDA @ DAΦNE
- FOPI @ GSI
- OBELIX


future experiments

- FOPI @ GSI
- E15 @ J-PARC
- FAIR @ GSI

... and AMADEUS

International workshop on Hadronic Atoms and
Kaonic Nuclei, ECT* Trento,
26-Oct-09

present experimental/theoretical situation (J. Mares)



The background of the slide features a faint, light-colored musical score with various notes and clefs, providing a thematic context for the presentation.

Experimental programme

- The scientific case of the so-called “deeply bound kaonic nuclear states” is hotter than ever, both in the theoretical (intensive debate) and experimental sectors.
- What emerges is the strong need for a complete experimental study of the scientific case, i.e. a clear and clean experiment (so without the need to make hypothesis on involved physics processes), measuring kaonic clusters both in formation and in the decay processes.
- AMADEUS’s main aim is to perform a full acceptance, high precision measurement of DBKNS both in formation and in the decay processes, by implementing the KLOE detector with an inner AMADEUS-dedicated setup, containing a cryogenic target and a trigger system.

Experimental programme

- The scientific case of the so-called “deeply bound kaonic nuclear states” is hotter than ever, both in the theoretical (intensive debate) and experimental sectors.
- What emerges is the strong need for a complete experimental study of the scientific case, i.e. a clear and clean experiment (so without the need to model, or to involve processes), measuring kaonic clusters both in formation and in the decay processes.
Either situations: existence or not existence of DBKNS will have strong impact in kaon-nucleon/nuclei physics!
- AMADEUS’s main aim is to perform a full acceptance, high precision measurement of DBKNS both in formation and in the decay processes, by implementing the KLOE detector with an inner AMADEUS-dedicated setup, containing a cryogenic target and a trigger system.

The background of the slide features a faint, light-colored musical score with multiple staves and notes, typical of a piano or guitar arrangement. The score is partially obscured by the text and the title.

Experimental programme

study of the (most) fundamental antikaon deeply bound nuclear systems,

the **kaonic dibaryon states: $pp\bar{K}^-$ and $(pn\bar{K}^-)$**

produced in a ^3He gas target, in formation and decay processes

as next step, the **kaonic 3-baryon states: $ppn\bar{K}^-$ and $pnn\bar{K}^-$**

produced in a ^4He gas target, in formation and decay processes

Experimental programme

- Low-energy charged kaon cross sections and interactions on H, d, Helium(3 and 4), for K^- momentum lower than 100 MeV/c (missing today);
- The K^- nuclear interactions in Helium reactions (poorly known, based on one paper from 1970 ...)
- Resonance states as the $\Lambda(1405)$ or the $\Sigma(1385)$ could be better understood with high statistics; their behaviour in the nuclear medium can be studied too.

The search for antikaon-mediated deeply bound nuclear states with AMADEUS

AMADEUS aims to confirm or deny the existence of such exotic states performing a full acceptance, high precision measurement of DBKNS both in formation and in the decay process, implementing the KLOE detector with an inner AMADEUS dedicated setup.


;

This requires the detection of:

- charged and neutral particles
- up to about 800 MeV/c
- in a 4π geometry
- with high efficiency and resolution

The search for antikaon-mediated deeply bound nuclear states with AMADEUS

Reaction channels (simplified)



Measure 1 particle of a 2-body decay.
 Transform to cms of the decaying
 Object. Gives 2nd particle properties.
Missing mass spectroscopy

Measure all outgoing particles to obtain the
 total cms energy = *invariant mass of the object*

Setup performance requirements

Formation processes




Study of the exotic states by the energy distribution of the ejected protons and neutrons. The setup should be able to measure:

- position of K^- stop: primary vertex and K^+ tracking (trigger)
- outgoing neutrons and protons

Setup performance requirements

Decay processes



Invariant mass spectroscopy

this requires:



- identification of all decay products, including protons neutrons and pions from hyperons decay
- measurement of 4-momenta of charged and neutral particles
 - protons 200 – 800 Mev/c ; pions 50 – 200 Mev/c ; neutrons 200 – 800 Mev/c ; deuterons ...

requirements satisfied by..

double ring $e^+ e^-$ collider working
in C. M. energy of ϕ , producing
 $\approx 600 K^+ K^- /s$

- **low momentum** Kaons
 $\approx 127 \text{ Mev}/c$
- **back to back** $K^+ K^-$
topology

DAΦNE



KLOE

- 96% acceptance,
- optimized in the energy range of all charged particles involved
- good performance in detecting neutrons checked by kloNe group


M. Anelli et al., Nucl. Instr. Meth. A 581, 368 (2007)

The experimental setup of AMADEUS

•The AMADEUS setup will be implemented in the 50 cm. gap in KLOE DC around the beam pipe:

•**Target**(A gaseous He target for a first phase of study)

•**Trigger** (1 or 2 layers of ScFi surrounding the interaction point)



KLOE -
EMC

KLOE -
Drift Chamber


experimental setup: trigger system

- **Cilindrical layer of scintillating fibers** surrounding the beam pipe to **trigger $K^+ K^-$ in opposite directions**
- Single or double layer





In this case possibility of perform tracking as well:
X-Y measurement with high granularity layers

- Readout to be done by **SiPM (silicon photo-multipliers)**



experimental setup: trigger system





prototipe of the trigger system

layers of BCF-10 fibers double cladded
free to rotate
read at both sides by Hamamatsu
S10362-11-050-U SiPM

is now under test

experimental setup: target

AMADEUS Monte Carlo




Low-mass cryogenic
gas target cell:
T = 10 K
P = 1.0 bar
R_{in} = 5 cm
R_{out} = 15 cm
L = 20 cm

half-toroidal cryogenic target cell
inside a vacuum chamber,
and
two more layers of fibers

Analysis of K^- He interactions in the KLOE drift chamber


Exotic states are expected to predominantly decay into final states containing Σ , Λ , p , n , d , as an example the decay channels of the kaonic tribaryon state are:



important feature of the **detector** and the **tracking procedure** is the **reconstruction capability for Λ 's and Σ 's**

main source of background comes from classical hadronic interactions of K^- in ${}^4\text{He}$ (poorly known based on one paper from 1970)

Analysis of K^- He interactions in the KLOE drift chamber




KLOE Drift Chamber

- The **drift chamber** of KLOE contains **mainly ^4He** (90% helium, 10% isobutane)
- From analysis of KLOE Monte Carlo **0.1% of K^-** from DAΦNE should **stop in the DC volume**
- total amount of analyzed data up to a luminosity of $\approx 1.8 \text{ fb}^{-1}$ from KLOE data (K charged group)
- kaons tag system: 2-body decay and/or dE/dx signature in the DC gas.

Strategy

- search for hadronic interactions with $\Lambda(1116)$ as product $\Lambda \rightarrow p + \pi^-$ (64% BR), $\Lambda \rightarrow n + \pi^0$ (36% BR)
- construct a vertex $\Lambda + \text{other particle}$

Analysis of K^- He interactions in the KLOE drift chamber



Λ invariant mass reconstruction.

The background of the slide features a faint, light-colored musical score with two staves per system, numbered 117, 120, 123, 126, and 129. The notes are mostly eighth and sixteenth notes, some beamed together. The title 'Conclusions' is centered over the middle of the page.

Conclusions

- The AMADEUS collaboration aims to perform a complete search for deeply bound kaonic nuclear states, and study of low energy K^- light nuclei interaction
- To this end an AMADEUS dedicated setup will be implemented in KLOE (data taking to start after KLOE2)
- All charged and neutral particles involved in formation and decay processes will be detected in a 4π geometry
- The reconstruction capability for Λ 's and Σ 's was tested analyzing KLOE data

AMADEUS scientific case

Deeply bound Kaonic nuclear states:

- In presence of strong KN attractive potential were firstly suggested by Wycech

(S. Wycech, Nucl. Phys. A450 (1986) 399c)

- Y. Akaishi and T. Yamazaki 'nuclear bound states in light nuclei'

(Phys. Rev. C65 (2002) 044005)



- strong attractive $l=0$ interaction KN interaction favours discrete nuclear states, bound 100-200 Mev, narrow 20-30 Mev
- shrinkage effect of a K on core nuclei forming unusual dense nuclear medium

AMADEUS scientific case

Deeply bound Kaonic nuclear states requires the presence of a strong attractive KN interaction in the isospin $I=0$ channel.

From experimental data:

- S-wave K^- nucleon scattering length is negative at threshold
- K_{α} line shift of kaonic hydrogen is negative

Repulsive type interaction

KN potential strongly dependent on density:

- **repulsive** in **free space**
- **attractive** in **nuclear matter**

The background of the slide features a faint, repeating pattern of musical notation, including treble and bass clefs, staves, and various notes and rests, arranged in a grid-like fashion.

State of the experimental search and theoretical debate for DBKS

- Possible experimental indications of the formation of kaonic nuclear states have received alternative explanations in the framework of known processes
- Recent calculations of k_{pp} systems suggests relatively moderate bindings and large widths

N. V. Sevchenko, A. Gal, J. Mares, J. Revai, Phys. Rev. C 76, 044004 (2007)
A. Dote, T. Hyodo, W. Weise, Nucl. Phys. A 804, 197 (2008)




new complete experimental results are needed


Selection criteria for Λ

requests:


- vertex with at least two opposite charged particles
- spatial position of vertex inside DC, or in DC entrance wall
- negative tracks with $dE/dx < 95 \text{ ADC}_{\text{counts}}$



protons having right E-p relation using energy released in the calorimeter



selection with energy loss in DC



cut because treshold of calorimeter

