

ID contributo: 83 Tipo: Parallel Contribution

## Results on the $K^-_{stop}A \to \Sigma^{\pm}\pi^{\mp}A'$ reaction on light nuclei.

martedì 11 ottobre 2011 17:55 (25 minuti)

The  $K_{stop}^-A \to \Sigma^{\pm}\pi^{\mp}A'$  reaction is studied on light nuclei,  $A = ^{6,7}Li, ^9Be, ^{13}C$  and  $^{16}O$ .

The outgoing  $\Sigma$ 's and  $\pi$ 's are detected using the FINUDA spectrometer, which operated at the DA $\Phi$ NE  $e^+e^-$  facility (LNF). The  $\Sigma^\pm$  hyperons are reconstructed via the  $n\pi^\pm$  decay with the neutrons detected by TOFONE, a large volume plastic scintillator array. The two final  $\pi^\pm$  mesons are reconstructed by means of the FINUDA tracker, which consists of 5 position sensitive layers. Final  $\Sigma^\pm\pi^\mp$  pairs are selected requiring topological cuts for the  $n\pi^\pm$  correlated pairs, where the  $n\pi^\pm$  pairs are requested to have  $\Sigma^\pm$  invariant mass. \\ The  $\Sigma^\pm\pi^\mp/K_{stop}^-$  emission rates are reported as a function of A. These rates are compared to previous experimental measurements. They are also used to calculate the  $\gamma$  ratio ( $\gamma = \Sigma^+\pi^-/\Sigma^-\pi^+$ ) which strongly increases when the kaon is absorbed on an in-medium proton instead of a free proton. This effect is related to the sub-threshold behavior of the  $\bar{K}N$  interaction. \\ The momentum spectra of prompt pions and free sigmas are also discussed as well as the  $\Sigma^\pm\pi^\mp$  missing mass behavior. In this case, the  $\Sigma^\pm\pi^\mp$  channel is filled by two resonances  $\Sigma(1385)$  and  $\Lambda(1405)$  as well as by the  $\Sigma^\pm\pi^\mp$  quasi-free reaction whose phase space develops in the same region as the two resonances.

Autore principale: Prof. CAMERINI, Paolo (Univ. Trieste & INFN Trieste)

Relatore: Prof. CAMERINI, Paolo (Univ. Trieste & INFN Trieste)

Classifica Sessioni: Hadron Physics II

Classificazione della track: Hadron Physics