

WPCF 2023 - XVI Workshop on Particle Correlations and Femtoscopy & IV Resonance Workshop 2023



Contribution ID: 35

Type: **Invited**

Dineutron correlation in neutron drip-line nuclei

Tuesday, 7 November 2023 14:30 (25 minutes)

The dineutron is a hypothetical bound state of two neutrons in a nuclear medium and a spatially compact pair, different from the one realized by the BCS mechanism[1]. The dineutron correlation is expected to appear in various circumstances, such as the surface of weakly bound neutron-rich systems and the inner crust of neutron stars. It has been studied using various approaches, such as the breakup reactions[2,3]. However, previous measurements were insufficient to discuss the magnitude of the dineutron correlation and its density dependence due to the sensitivity of the probe[4].

In the present study, it is found for the first time that the dineutron in the ^{11}Li nucleus is localized on the surface of the ^{11}Li nucleus. The use of the quasi-free (p, pn) reaction was essential to extract the radial information of the dineutron and to minimize the effect of the final state interactions[5]. The published results [6] and recent updates will be presented.

References

- [1] A. B. Migdal, Soviet J. Nucl. Phys. **16**, 238 (1973).
- [2] H. Simon et al., Phys. Rev. Lett. **83**, 496 (1999).
- [3] T. Nakamura et al., Phys. Rev. Lett. **96**, 252502 (2006).
- [4] Y. Kikuchi et al., Prog. Theor. Exp. Phys. **2016**, 103D03 (2016).
- [5] Y. Kikuchi et al., Phys. Rev. C **87**, 034606 (2013).
- [6] Y. Kubota et al., Phys. Rev. Lett. **125** 252501 (2020).

Primary author: KUBOTA, Yuki (RIKEN Nishina Center)

Presenter: KUBOTA, Yuki (RIKEN Nishina Center)

Session Classification: Day 2 - Afternoon