

HOLOGRAPHY

IN ELASTIC pp SCATTERING AND BINARY STARS

T. Csörgő^{1,2}

¹ HUN-REN Wigner RCP, Budapest, Hungary

² MATE KRC, Gyöngyös, Hungary

Motivation:

**Proton Holography
phase reconstruction**

e-Print: 2004.07095 [hep-ph], EPJ Web Conf. 235 (2020) 06002



Stellar Interferometry:

Binary stars

New results, motivated by M. Lisa's and Naomi Vogel's talk at WPCF2023

Conclusion

INTRODUCTION: HOLOGRAPHY



HOYGENS
YOUNG
FRESNEL

HOYGENS
YOUNG
FRESNEL

Fig 4.
First Holographic Reconstruction, 1948

Basic idea of holography (1947): amplitude level reconstruction.
First hologram (1948) from D. Gabor's Nobel lecture (1967).

<https://www.nobelprize.org/uploads/2018/06/gabor-lecture.pdf>

Formalism: elastic pp scattering

$$\sigma_{el}(s) = \int_0^\infty d|t| \frac{d\sigma(s)}{dt}$$

$$\frac{d\sigma(s)}{dt} = \frac{1}{4\pi} |T_{el}(s, \Delta)|^2, \quad \Delta = \sqrt{|t|}.$$

$$B(s, t) = \frac{d}{dt} \ln \frac{d\sigma(s)}{dt}$$

$$B(s) \equiv B_0(s) = \lim_{t \rightarrow 0} B(s, t),$$

$$\sigma_{tot}(s) \equiv 2 \operatorname{Im} T_{el}(\Delta = 0, s)$$

$$\rho(s, t) \equiv \frac{\operatorname{Re} T_{el}(s, \Delta)}{\operatorname{Im} T_{el}(s, \Delta)}$$

$$\rho(s) \equiv \rho_0(s) = \lim_{t \rightarrow 0} \rho(s, t)$$

Basic problem: $d\sigma/dt$ measures an amplitude, *modulus squared*.
Amplitude level reconstruction??? Phase info apparently lost...

MODEL INDEPENDENT LEVY EXPANSION

$$\frac{d\sigma}{dt} = A w(z|\alpha) \left| 1 + \sum_{j=1}^{\infty} c_j l_j(z|\alpha) \right|^2,$$

$$w(z|\alpha) = \exp(-z^\alpha), \text{ non-exponential behavior (NEB) in a single parameter}$$

$$z = |t|R^2 \geq 0, \quad \alpha$$

$$c_j = a_j + ib_j, \quad \text{idea: complete set of orthonormal functions, put NEB to the weight}$$

$$l_j(z|\alpha) = D_j^{-\frac{1}{2}} D_{j+1}^{-\frac{1}{2}} L_j(z|\alpha),$$

$$D_0(\alpha) = 1,$$

$$D_1(\alpha) = \mu_{0,\alpha},$$

$$D_2(\alpha) = \det \begin{pmatrix} \mu_{0,\alpha} & \mu_{1,\alpha} \\ \mu_{1,\alpha} & \mu_{2,\alpha} \end{pmatrix},$$

$$D_3(\alpha) = \det \begin{pmatrix} \mu_{0,\alpha} & \mu_{1,\alpha} & \mu_{2,\alpha} \\ \mu_{1,\alpha} & \mu_{2,\alpha} & \mu_{3,\alpha} \\ \mu_{2,\alpha} & \mu_{3,\alpha} & \mu_{4,\alpha} \end{pmatrix},$$

$$\int_0^\infty dz \exp(-z^\alpha) l_n(z|\alpha) l_m(z|\alpha) = \delta_{n,m}$$

$$\mu_{n,\alpha} = \int_0^\infty dz z^n \exp(-z^\alpha) = \frac{1}{\alpha} \Gamma\left(\frac{n+1}{\alpha}\right)$$

T. Csörgő, R. Pasechnik, A. Ster,
[arxiv.org:1807.02897](https://arxiv.org/abs/1807.02897)

Eur.Phys.J.C 79 (2019) 1, 62


$$L_0(z|\alpha) = 1,$$

$$L_1(z|\alpha) = \det \begin{pmatrix} \mu_{0,\alpha} & \mu_{1,\alpha} \\ 1 & z \end{pmatrix},$$



$$L_2(z|\alpha) = \det \begin{pmatrix} \mu_{0,\alpha} & \mu_{1,\alpha} & \mu_{2,\alpha} \\ \mu_{1,\alpha} & \mu_{2,\alpha} & \mu_{3,\alpha} \\ 1 & z & z^2 \end{pmatrix},$$

$$L_3(z|\alpha) = \det \begin{pmatrix} \mu_{0,\alpha} & \mu_{1,\alpha} & \mu_{2,\alpha} & \mu_{3,\alpha} \\ \mu_{1,\alpha} & \mu_{2,\alpha} & \mu_{3,\alpha} & \mu_{4,\alpha} \\ \mu_{2,\alpha} & \mu_{3,\alpha} & \mu_{4,\alpha} & \mu_{5,\alpha} \\ 1 & z & z^2 & z^3 \end{pmatrix},$$



ABILITIES: CONVERGES TO pp $d\sigma/dt$ @ 13 TeV



CONVERGENCE PROPERTIES OF LEVY SERIES



Partial sum converges both in pbarp (n=3) and also in pp (n=4)




Partial sum converges to profile function $P(b)$ and slope $B(t)$

CONVERGENCE OF PHASE RECONSTRUCTION

Partial sums contributions to $\phi(t)$, 13 TeV pp

$$T_{el}(t) = |T_{el}(t)| \exp [i\phi_1(t)]$$




Levy expansion converges to the phase $\phi(t)$: Proton holography ?!
Cross-check with Coulomb-Nuclear Interference at $t=0$ successful!
Deeper level of understanding – in progress, but stop here for now.

SUMMARY 1: PROTON HOLOGRAPHY


Partial sums contributions to $\phi(t)$, 13 TeV pp

$$T_{e1}(t) = |T_{e1}(t)| \exp [i\phi_1(t)]$$



Levy expansion converges to the phase $\phi(t)$: Proton holography !


INTENSITY INTERFEROMETRY FOR BINARIES



Source	Dschubba (Delta Scorpii)
Magnitude (mag)	2.2
Spectral type	B0.3IV
System	Binary star
HBT time (h)	115.1
HBT diameter (mas)	0.45 ± 0.04
Time (h)	5.1
Diameter (mas)	470nm: 0.613 ± 0.072 375nm: 0.612 ± 0.081

Stellar interferometry results, HESS, talk of Naomi Vogel@ WPCF23:
Fits a **BINARY** star with intercept parameter $\lambda = 1$
But **this is a puzzle**, see R. Hanbury Brown et al, MNRAS (1974) **167**, 121


HBT FOR BINARY AND MULTIPLE STARS




The normalized correlation also depends upon whether a star is single or multiple. It was shown in Paper II that, if a star is binary and the angular separation of the two components is completely resolved by the interferometer at the shortest baseline, then the normalized zero-baseline correlation $\overline{c_N(0)}$ averaged over a range of position angles is reduced relative to a single star $c_N(0)$ by the factor,

$$\overline{c_N(0)}/c_N(0) = (I_1^2 + I_2^2)/(I_1 + I_2)^2 \quad (9)$$

Mon. Not. R. astr. Soc. (1974) **167**, 121-136.




(b)



THE ANGULAR DIAMETERS OF 32 STARS

R. Hanbury Brown, J. Davis and L. R. Allen



Core-halo picture:

T. Cs, B. Lörstad and J. Zimányi, *Z.Phys.* C71 (1996) 491-497

J. Bolz et al, *Phys.Rev.D* 47 (1993) 3860-3870


Fig. 7. a) Two stars, S , and T , along nearby lines of sight from the earth; b) of correlated intensity from the two stars; c) schematic of HBT measurement halo of dim stars.

Q /MeV/

Fig. 7 from G. Baym's review paper: arXiv:nucl-th/9804026

Schematics of HBT for multiple stars, but no formula. Note: intercept!

HBT FOR MARGINALLY RESOLVED STARS



projection of the baseline (Hanbury Brown et al. 1967):

$$V^2(r) = \frac{1}{(I_a + I_b)^2} \times \left[I_a^2 V_a^2(r) + I_b^2 V_b^2(r) + 2I_a I_b V_a(r) V_b(r) \cos\left(\frac{2\pi r \bar{\theta} \cos \psi}{\lambda_0}\right) \right],$$

Partial resolution of the components from W. Guerin et al,
MNRAS 480, 245–250 (2018)

Reduction of average measured intensity reduces:

R. H. Brown and R. Q. Twiss, Brown, R. Hanbury, and R. Q. Twiss: in *Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences* (1958): 291-319.

HBT EFFECT FOR „IDENTICAL TWIN“ STARS

$$\rho(x) = f_+ s(x - x_+) + f_- s(x - x_-),$$

$$f_+ + f_- = 1$$

Normalization: $\text{FT}(s|q = 0) = 1$

$$C(q) = 1 \pm |\tilde{\rho}(q)|^2 = 1 \pm \Omega(q) |\tilde{s}(q)|^2,$$

Stellar model: uniformly illuminates sphere $|\tilde{s}(q)|^2 = \left[\frac{2J_1(\pi r \theta / \lambda_0)}{\pi r \theta / \lambda_0} \right]^2$.

Normalization: $C(q = 0) = 1 + 1$ but with an oscillating prefactor

$$\Omega(q) = [(f_+^2 + f_-^2) + 2f_+ f_- \cos[q(x_+ - x_-)]]$$

12

Binary source formalism: binary sources in hydro (Cooper-Frye)

Schematics worked out for identical stars, e-Print: [hep-ph/0011320](https://arxiv.org/abs/hep-ph/0011320)

Detailed **review** including hydro results with two saddle points in [hep-ph/0001233](https://arxiv.org/abs/hep-ph/0001233)

HBT FOR BINARY AND MULTIPLE STARS

Mon. Not. R. astr. Soc. (1974) **167**, 121–136.

THE ANGULAR DIAMETERS OF 32 STARS

R. Hanbury Brown, J. Davis and L. R. Allen

depends upon whether a star is single or binary and the angular separation of the two components is completely resolved by the interferometer at the shortest baseline, then the normalized zero-baseline correlation $\overline{c_{N(0)'}}$ averaged over a range of position angles is reduced relative to a single star $\overline{c_{N(0)}}$ by the factor,

$$\overline{c_{N(0)'}}/\overline{c_{N(0)}} = (I_1^2 + I_2^2)/(I_1 + I_2)^2 \quad (9)$$

where I_1, I_2 are the brightness of the two components. It is simple to extend this analysis to a multiple star with n components and to show that, if the angular separation between all the components is resolved, the zero-baseline correlation is reduced relative to a single star by the factor,

$$\overline{c_{N(0)'}}/\overline{c_{N(0)}} = \sum_n I^2 / \left(\sum_n I \right)^2. \quad (10)$$

It follows that if a star yields a correlation which is significantly less than that expected from a single star, then it must be multiple.

R. Hanbury Brown, J. Davis and L. R. Allen, MNRAS (1974) **167** 121

Note: $f_1 = I_1/(I_1 + I_2)$, $f_2 = I_2/(I_1 + I_2)$, etc \rightarrow HEP connection

HBT FOR „NON-IDENTICAL TWIN“ STARS

Similar, but two sources have different sizes, but with

$$f_+ + f_- = 1$$

Stellar model: uniformly illuminates sphere or any unknown shape

An oscillating prefactor remains, and averages to

$$\overline{\Omega(q)} = [(f_+^2 + f_-^2) + \cancel{2f_+f_- \cos[q(x_+ - x_-)]}]$$

Binary source formalism: effective reduction of intercept
for well separated compact sources


$$\text{Effectively } \frac{1}{2} \leq \lambda = \overline{\Omega(q)} \leq 1$$

Schematics works even for out non-identical stars,
observations of N. Vogel explained

Multiple n source, well separated compact sources

$$\text{Easy to show: } \frac{1}{n} \leq \lambda = \overline{\Omega_n(q)} \leq 1$$

SUMMARY FOR BINARY STARS



Source	Dschubba (Delta Scorpii)
Magnitude (mag)	2.2
Spectral type	B0.3IV
System	Binary star
HBT time (h)	115.1
HBT diameter (mas)	0.45 ± 0.04
Time (h)	5.1
Diameter (mas)	470nm: 0.613 ± 0.072 375nm: 0.612 ± 0.081

Stellar interferometry results, HESS, from N. Vogel's talk @ WPCF23: Fits with intercept parameter $\frac{1}{2} \leq \lambda \leq 1$ possible, **puzzle resolved.**