

HIDDe 

Hunting Invisibles: Dark sectors, Dark matter and Neutrinos

# Asymmetries in Extended Dark Sectors: A Cogenesis Scenario

Juan Herrero-García, GL, Drona Vatsyayan  
[2301.13238]

**Giacomo Landini**




14/09/2023



# Dark Matter

Dark Matter is five times more abundant than baryonic matter




$$\rho_{\text{DM}} \simeq 5\rho_B$$

Cosmic coincidence


➔ A common origin for baryons and DM?

# Standard (symmetric) production

**Freeze-out:** DM is in thermal equilibrium with the SM



**Freeze-in:** DM is **not** in thermal equilibrium with the SM



Dark Matter is **symmetric** (?)

$$Y_{\chi} = Y_{\bar{\chi}}$$

# Asymmetric Dark Matter

The Baryon abundance is set by an asymmetry

$$\eta_B = Y_b - Y_{\bar{b}} = 0.88 \times 10^{-11}$$

,

The nature of DM could also be asymmetric!

$$\eta_D = Y_{\chi} - Y_{\bar{\chi}}$$

$$\frac{\rho_{\text{DM}}}{\rho_B} = \frac{m_{\text{DM}} \eta_D}{m_p \eta_B} \simeq 5$$



Attractive scenario:  $\eta_D \simeq \eta_B \longrightarrow m_{\text{DM}} \simeq 5m_p \simeq 5 \text{ GeV}$

# Asymmetric freeze-out

Asymmetry and Fractional asymmetry

$$\eta_D = Y_\chi - Y_{\bar{\chi}} \quad r = Y_{\bar{\chi}}/Y_\chi$$

Asymmetry generation






# Asymmetric freeze-out

Asymmetry and Fractional asymmetry

$$\eta_D = Y_\chi - Y_{\bar{\chi}} \quad r = Y_{\bar{\chi}}/Y_\chi$$

Annihilation of the symmetric component






$$r_\infty \propto \exp \left[ -\frac{\pi g_*}{45 x_f} M_{\text{Pl}} \sigma v \eta_D m \right]$$

Graesser, Shoemaker, Vecchi  
[1103.2771]

# Asymmetric freeze-out

Asymmetry and Fractional asymmetry

$$\eta_D = Y_\chi - Y_{\bar{\chi}} \quad r = Y_{\bar{\chi}}/Y_\chi$$



$$r_\infty \propto \exp \left[ -\frac{\pi g_*}{45 x_f} M_{\text{Pl}} \sigma v \eta_D m \right]$$




Graesser, Shoemaker, Vecchi  
[1103.2771]

# Asymmetric freeze-out

Asymmetry and Fractional asymmetry

$$\eta_D = Y_\chi - Y_{\bar{\chi}} \quad r = Y_{\bar{\chi}}/Y_\chi$$

Annihilation of the symmetric component



$$r_\infty \propto \exp \left[ -\frac{\pi g_*}{45 x_f} M_{\text{Pl}} \sigma v \eta_D m \right]$$

Graesser, Shoemaker, Vecchi  
[1103.2771]





# Asymmetric freeze-in?

Asymmetry and Fractional asymmetry

$$\eta_D = Y_\chi - Y_{\bar{\chi}} \quad r = Y_{\bar{\chi}}/Y_\chi$$

Annihilation of the symmetric component



**Asymmetric DM out of equilibrium (tiny couplings, freeze-in) ?**



(How to erase the symmetric component?)

# Asymmetric freeze-in?

Asymmetry and Fractional asymmetry



$$\eta_D = Y_\chi - Y_{\bar{\chi}} \quad r = Y_{\bar{\chi}}/Y_\chi$$

Annihilation of the symmetric component



**Idea:** late decays of an asymmetric species after symmetric population has been erased


# Early vs Late decays



Dominant production from early decays takes place around  $x \sim 1$



Asymmetric freeze-out takes place around  $x \sim 20$

Late decays take place at  $x \gg 1$




No annihilations

# Early vs Late decays




**Asymmetric FIMP DM!**



# A Cogenesis scenario

Falkowski, Ruderman, Volansky [1101.4936]

CP-violating decays of **RHNs**  
out of equilibrium



**Dark asymmetry generation**

$$\eta_\chi = \eta_S \sim \eta_B$$

$$\bar{\chi}\chi, S^\dagger S$$


annihilations erase  
symmetric components

Lepton asymmetry


Sphalerons  
↓

**Baryon asymmetry**  $\eta_B$

# A Cogenesis scenario



# A Cogenesis scenario




This scenario can explain neutrino masses, baryon asymmetry and (FIMP) Dark Matter!

# The model

Field	Spin	$U(1)_{B-L}$	$U(1)_D$	$U(1)_X$
$N_R^i$	1/2	-1	0	0
$\sigma$	0	+2	0	0
$\chi_0$	1/2	-1	1	0
$\psi_0$	1/2	0	0	+1
$S$	0	0	-1	0
$\phi$	0	+1	-1	+1

DM stability +  
Dirac Nature

$$M_{N_3}, M_{N_2} \gg M_{N_1} \gg m_\chi^0 \gg m_\psi^0, m_S > m_\phi$$



$$y_\phi \ll 1, g_X \ll 1$$

$\psi$  is **not** in thermal equilibrium

$$\mathcal{L}_{\text{int}} = -y_\nu^{\alpha i} \bar{L}^\alpha \tilde{H} N_R^i - y_\sigma^{ij} \sigma \overline{N_R^i} N_R^j - y_S^i S \bar{N}_R^i \chi_0 - y_\phi \phi \bar{\psi}_0 \chi_0 + \text{H.c.}$$

Neutrino masses

$$M_{N_1} \propto \langle \sigma \rangle = v_{B-L} \longrightarrow m_\nu = -m_D M_N^{-1} m_D^T$$



# The model

$$\begin{array}{ccccc}
 U(1)_{B-L} \otimes U(1)_D \otimes U(1)_X & \rightarrow & U(1)_D \otimes U(1)_X & \rightarrow & U(1)_{X+D} \\
 & & \downarrow & & \downarrow \\
 & & \langle \sigma \rangle = v_{B-L} > 10^{11} \text{ GeV} & & \langle \phi \rangle = v_\phi \ll v_{B-L}
 \end{array}$$


	Particle	$U(1)_{X+D}$
<u>DM candidates</u>	$\chi$	+1
	$\psi$	+1
	$S$	-1

$S \rightarrow \bar{\psi} + \bar{\nu}$  or  $\psi \rightarrow S^\dagger + \nu$  are allowed but suppressed


Both cosmologically stable

**Multicomponent DM**

# DM production (Fermion)



$$T_D \simeq 10 \text{ MeV} \left( \frac{y_\phi}{10^{-12}} \right)$$



Early decays are dominant

$$Y_\psi = Y_{\bar{\psi}}$$


Symmetric DM

Late decays are dominant


$$Y_\psi = \eta_\psi \gg Y_{\bar{\psi}}$$

Asymmetric DM


# DM production (Scalar)



# DM production (Scalar)



# DM production (Scalar)




$$R = \frac{\Gamma(\chi \rightarrow S^\dagger \nu)}{\Gamma(\chi \rightarrow \psi \phi)} \sim \frac{|y_S|^2}{y_\phi^2} \frac{m_\nu}{M_{N_1}}$$

$R \ll 1$  No extra decays, asymmetric DM (freeze-out)  
 $Y_S = \eta_S$  ,  $Y_{S^\dagger} = \frac{R}{1+R} \eta_S$

$R \gg 1$  Cancellation of the asymmetry, symmetric DM (freeze-out)


# Scenarios





Sc.	$\psi$ population	$S$ population	$10^{-10} y_\phi / \sqrt{\eta_D / \eta_B}$	$R$	$T_D^{(S)} / T_\star^{(S)}$
<b>1</b>	Asymmetric	Asymmetric	$\leq 0.06$	$\ll 1$	Any
<b>2</b>	Asymmetric	Partially Asymmetric	$\leq 0.06$	$\mathcal{O}(1)$	$< 1$
<b>1-2</b>	Asymmetric	Asymmetric	$\leq 0.06$	$\mathcal{O}(1)$	$> 1$
<b>3</b>	Partially Asymmetric	Asymmetric	$0.06 - 2$	$\ll 1$	Any
<b>4</b>	Partially Asymmetric	Partially Asymmetric	$0.06 - 2$	$\mathcal{O}(1)$	$< 1$
<b>3-4</b>	Partially Asymmetric	Asymmetric	$0.06 - 2$	$\mathcal{O}(1)$	$> 1$
<b>5</b>	Symmetric	Asymmetric	$\gtrsim 2$	$\ll 1$	Any
<b>6</b>	Negligible	Symmetric	$y_\phi \lesssim 5 \times 10^{-7}$	$\gtrsim \mathcal{O}(10)$	$< 1$

# Scenarios

## Freeze-in



## Thermal



$$\frac{\Omega_\psi}{\Omega_S} = \frac{m_\psi(\eta_D + Y_{FI})}{\eta_D m_S f(R)}, \quad \frac{\Omega_{DM}}{\Omega_B} = \frac{m_\psi(\eta_D + Y_{FI}) + \eta_D m_S f(R)}{\eta_B(1+R)m_p},$$

$$f(R) \equiv \begin{cases} 1 + 2R & \text{if } T_D^{(S)} < T_*^{(S)} \\ 1 & \text{if } T_D^{(S)} > T_*^{(S)}. \end{cases}$$

The DM abundance is reproduced for  $m_\psi, m_S \sim \text{GeV}$

# Phenomenology

Freeze-in: small couplings, suppressed DD

$$y_\phi \ll 1 \quad g_X \ll 1$$

ADM: suppressed ID + annihilations in dark sectors

Large  $B - L$  scale: no collider searches

$$v_{B-L} \gtrsim 10^{11} \text{ GeV}$$

DD of scalar DM through Higgs portal

$$\lambda_{HS}(H^\dagger H)(S^\dagger S)$$

ID of scalar DM when  $S$  is symmetric

Enhanced ID signal for **Scenario 6**

$$Y_S = Y_{S^\dagger} = \eta_S \quad \begin{array}{l} m_S = 2.5 \text{ GeV} \\ \sigma v > \sigma v_{\text{WIMP}} \end{array}$$


**+ neutrino line!**



# Smoking gun: neutrino line

$$\mathcal{O}_6 = \bar{L}\tilde{H}S\phi^\dagger\psi$$

generated at  $E \ll m_\chi \ll M_{N_1}$



(analogous process if  $m_\psi > m_S$ )

$$\Gamma(S \rightarrow \bar{\psi} + \nu) = \frac{|y_S|^2 y_\phi^2 m_S}{32\pi} \left(\frac{v_\phi}{m_\chi}\right)^2 \left(\frac{m_\nu}{M_{N_1}}\right) \left(1 - \frac{m_\psi^2}{m_S^2}\right)$$

Neutrino line peaked at  $E_\nu \simeq m_S/2 \longrightarrow \mathcal{O}(\text{GeV})$


Experimental bound

$$\tau > 10^{23} \text{ sec}$$

Future neutrino telescopes?

$$\tau \sim 10^{24-25} \text{ sec}$$


# Summary



# Summary

Symmetric 1DM with interesting features (mix Cold/Warm DM, enhanced ID signals,...)

Smoking Gun **neutrino line**: close to experimental sensitivity (neutrino experiments)



# Scenario 6: 1 DM

$$R = \frac{\Gamma(\chi \rightarrow S^\dagger \nu)}{\Gamma(\chi \rightarrow \psi \phi)} \sim \frac{|y_S|^2}{y_\phi^2} \frac{m_\nu}{M_{N_1}} \gg 1 \longrightarrow Y_S = Y_{S^\dagger} = \eta_S$$


No decays to  $\psi$   
1 DM

**Symmetric** population

$m_S = 2.5 \text{ GeV}$

**Asymmetry** sets the abundance

## Mix Cold/Warm DM



**Enhanced Indirect Detection** signals may be present  $\sigma v > \sigma v_{\text{WIMP}}$

# Scenario 6: 1 DM

$$R = \frac{\Gamma(\chi \rightarrow S^\dagger \nu)}{\Gamma(\chi \rightarrow \psi \phi)} \sim \frac{|y_S|^2}{y_\phi^2} \frac{m_\nu}{M_{N_1}} \gg 1 \longrightarrow Y_S = Y_{S^\dagger} = \eta_S$$


No decays to  $\psi$   
1 DM

**Symmetric** population

$m_S = 2.5 \text{ GeV}$

**Asymmetry** sets the abundance

**Mix Cold/Warm DM**




**Enhance** Detection signals may be present  $\sigma v > \sigma v_{\text{WIMP}}$

# Summary

- Asymmetric DM models needs large annihilations cross section: thermalization
- Asymmetric FIMP DM can be realized through late decays of asymmetric particle
- The framework naturally needs an extended dark sector: multicomponent DM, baryogenesis, neutrino masses
- Late DM decays can be constrained by neutrino experiments

# Backup

# Fermion annihilations





Sc.	$\psi$	$S$	$\Omega_{DM}/\Omega_B$	$\Omega_S/\Omega_\psi$
<b>1</b>	Asymmetric LD $\chi \rightarrow \psi\varphi$ $Y_\psi^+ = \eta_D$ $Y_\psi^- \ll Y_\psi^+$	Asymmetric FO $S^\dagger S \rightarrow \varphi\varphi$ $Y_S^+ = \eta_D$ $Y_S^- \ll Y_S^+$	$\frac{\eta_D}{\eta_B} \frac{m_\psi + m_S}{m_p}$	$\frac{m_\psi}{m_S}$
<b>2</b>	Asymmetric LD $\chi \rightarrow \psi\varphi$ $Y_\psi^+ = \eta_D/(1+R)$ $Y_\psi^- \ll Y_\psi^+$	Partially asymmetric FO $S^\dagger S \rightarrow \varphi\varphi$ + LD $\chi \rightarrow S^\dagger \nu_L$ $Y_S^+ = \eta_D$ $Y_S^- = \eta_D R/(1+R)$	$\frac{\eta_D}{\eta_B} \frac{m_\psi + (1+2R)m_S}{(1+R)m_p}$	$\frac{m_\psi}{m_S(1+2R)}$
<b>1-2</b>	Asymmetric LD $\chi \rightarrow \psi\varphi$ $Y_\psi^+ = \eta_D/(1+R)$ $Y_\psi^- \ll Y_\psi^+$	Asymmetric FO $S^\dagger S \rightarrow \varphi\varphi$ + LD $\chi \rightarrow S^\dagger \nu_L$ $Y_S^+ = \eta_D/(1+R)$ $Y_S^- \ll Y_S^+$	$\frac{\eta_D}{\eta_B} \frac{m_\psi + m_S}{(1+R)m_p}$	$\frac{m_\psi}{m_S}$
<b>3</b>	Partially asymmetric FI + LD $\chi \rightarrow \psi\varphi$ $Y_\psi^+ = Y_{FI}/2 + \eta_D$ $Y_\psi^- = Y_{FI}/2$	Asymmetric FO $S^\dagger S \rightarrow \varphi\varphi$ $Y_S^+ = \eta_D$ $Y_S^- \ll Y_S^+$	$\frac{m_\psi(\eta_D + Y_{FI}) + \eta_D m_S}{\eta_B m_p}$	$\frac{m_\psi(\eta_D + Y_{FI})}{m_S \eta_D}$
<b>4</b>	Partially Asymmetric FI + LD $\chi \rightarrow \psi\varphi$ $Y_\psi^+ = (Y_{FI}/2 + \eta_D)/(1+R)$ $Y_\psi^- = Y_{FI}/(2(1+R))$	Partially Asymmetric FO $S^\dagger S \rightarrow \varphi\varphi$ + LD $\chi \rightarrow S^\dagger \nu_L$ $Y_S^+ = \eta_D$ $Y_S^- = \eta_D R/(1+R)$	$\frac{m_\psi(\eta_D + Y_{FI}) + \eta_D(1+2R)m_S}{\eta_B(1+R)m_p}$	$\frac{m_\psi(\eta_D + Y_{FI})}{m_S \eta_D(1+2R)}$
<b>3-4</b>	Partially Asymmetric FI + LD $\chi \rightarrow \psi\varphi$ $Y_\psi^+ = (Y_{FI}/2 + \eta_D)/(1+R)$ $Y_\psi^- = Y_{FI}/(2(1+R))$	Asymmetric FO $S^\dagger S \rightarrow \varphi\varphi$ + LD $\chi \rightarrow S^\dagger \nu_L$ $Y_S^+ = \eta_D/(1+R)$ $Y_S^- \ll Y_S^+$	$\frac{m_\psi(\eta_D + Y_{FI}) + \eta_D m_S}{\eta_B(1+R)m_p}$	$\frac{m_\psi(\eta_D + Y_{FI})}{m_S \eta_D}$
<b>5</b>	Symmetric FI $\chi \rightarrow \psi\varphi$ $Y_\psi^+ = Y_{FI}/2 + \eta_D \simeq Y_{FI}/2$ $Y_\psi^- = Y_{FI}/2$	Asymmetric FO $S^\dagger S \rightarrow \varphi\varphi$ $Y_S^+ = \eta_D$ $Y_S^- \ll Y_S^+$	$\frac{\eta_D}{\eta_B} \frac{m_\psi(Y_{FI}/\eta_D) + m_S}{m_p}$	$\frac{m_\psi Y_{FI}}{m_S \eta_D}$
<b>6</b>	Negligible production	Symmetric FO $S^\dagger S \rightarrow \varphi\varphi$ + LD $\chi \rightarrow S^\dagger \nu_L$ $Y_S^+ = \eta_D$ $Y_S^- = \eta_D$	$< 1$	$\frac{\eta_D}{\eta_B} \frac{2m_S}{m_p}$


# Scalar annihilations

Field	Spin	$U(1)_{B-L}$	$U(1)_D$	$U(1)_X$
$N_R^i$	1/2	-1	0	0
$\sigma$	0	+2	0	0
$\chi_0$	1/2	-1	1	0
$\psi_0$	1/2	0	0	+1
$S$	0	0	-1	0
$\phi$	0	+1	-1	+1

$$\phi(x) = v_\phi + \varphi(x)/\sqrt{2}$$

Various terms in the scalar potential such as  $\lambda_{S\phi}|S|^2|\phi|^2$

$$\sigma v(S^\dagger S \rightarrow \varphi\varphi) \simeq \frac{\lambda_{S\phi}^2}{32\pi m_S^2} \left(1 - \frac{m_\varphi^2}{m_S^2}\right)^{1/2} \left(\frac{1 - m_\varphi^2/2m_S^2 - 2\lambda_{S\phi}v_\phi^2/m_S^2}{1 - m_\varphi^2/2m_S^2}\right)^2$$



# Asymmetric WIMP



Graesser, Shoemaker, Vecchi  
[1103.2771]

$$\frac{dr}{dx} = -\lambda \eta g_*^{1/2} x^{-n-2} \left[ r - \frac{Y_{eq}^2}{\eta^2} (1-r)^2 \right]$$

$$r = Y^- / Y^+$$

$$\eta = Y^+ - Y^- = \epsilon \eta_B$$

$$\lambda = \left( \frac{\pi}{45} \right)^{1/2} M_{Pl} m \sigma_0$$



# Partially asymmetric DM





$$\rho_{\text{DM}} = s \sum_i m_i \eta_i \left( \underset{\downarrow}{1} + 2 \frac{r_i}{\underset{\downarrow}{1-r_i}} \right)$$

Asymmetric                      Symmetric

# Cogenesis

Generation of the asymmetries through out-of-equilibrium decays

Falkowski, Ruderman, Volansky [1101.4936]




$$\varepsilon_L = \sum_{\alpha} \frac{\Gamma_{N_1 \rightarrow L_{\alpha} H} - \Gamma_{N_1 \rightarrow L_{\alpha} H^{\dagger}}}{\Gamma_{N_1}}, \quad \varepsilon_{\chi} = \frac{\Gamma_{N_1 \rightarrow \chi S} - \Gamma_{N_1 \rightarrow \bar{\chi} S^{\dagger}}}{\Gamma_{N_1}}$$

# Cogenesis



## Washout and transfer of the asymmetries

Falkowski, Ruderman, Volansky [1101.4936]




# Cogenesis

Falkowski, Ruderman, Volansky [1101.4936]



SM lepton asymmetry, neutrino spectrum and mixing angles are correctly reproduced

# Smoking gun: neutrino line



Coi, Gupta, Hambye [2104.00042]




# Dark Matter nature

Asymmetry and Fractional asymmetry

$$\eta_D = Y_\chi - Y_{\bar{\chi}} \quad r = Y_{\bar{\chi}}/Y_\chi$$

**Symmetric**


$$r_\infty > 0.9$$



$$\rho_{\text{DM}} \propto 1/\sigma v$$

**Asymmetric**


$$r_\infty < 10^{-2}$$



$$\rho_{\text{DM}} \propto m\eta_D$$

**Partially Asymmetric**

$$10^{-2} < r_\infty < 0.9$$




$$\rho_{\text{DM}} = f(m, \sigma v, \eta_D)$$

# Cold vs Warm

Small-scale structure constraint

$$m_S \gtrsim \underbrace{3.5 \text{ KeV} \langle p/T \rangle_{\text{prod}}}_{\mathcal{O}(\text{KeV})} \left( \frac{10}{g_s(T_D^S)} \right)^{1/3}$$

$\downarrow$   
 $\mathcal{O}(\text{GeV})$




# Cold vs Warm

Small-scale structure constraint

$$m_S \gtrsim 3.5 \text{ KeV} \langle p/T \rangle_{\text{prod}} \left( \frac{10}{g_s(T_D^S)} \right)^{1/3}$$

$\downarrow$   $\mathcal{O}(\text{GeV})$                        $\mathcal{O}(\text{GeV})$




$$\langle p/T \rangle_{\text{prod}} \simeq m_\chi / T_D^S \gg 1 \quad \textbf{Warm}$$

$\downarrow$

$$\mathcal{O}(\text{TeV})$$

# Gauge boson bounds

$$m_{Z_{B-L}} \geq 10^{11} \text{ GeV}$$



[Heeck, 2014]

$$g_X^{\text{eff}} A'_\mu J_{B-L}^\mu$$

$$\longrightarrow g_X \leq 10^{-8}$$

$$g_i^{\text{eff}} \sim g_i \left( \frac{v_\phi}{v_{B-L}} \right)^2 \sim 10^{-16} g_i$$

$$g_D^{\text{eff}} Z_{D,\mu} J_{B-L}^\mu$$

$$\longrightarrow g_D \leq 10^{13}$$

# A low-energy variant


Lowering the  $B - L$  scale  $\begin{cases} \rightarrow \text{Resonant leptogenesis} \\ \rightarrow \text{Inverse see-saw} \end{cases}$

Field	Spin	$U(1)_{B-L}$	$U(1)_D$	$U(1)_X$
$S_L$	1/2	0	0	0
$\sigma'$	0	+1	0	0

$$\langle \sigma' \rangle = v_{B-L} \sim \mathcal{O}(\text{TeV})$$

$$\mathcal{L}_{\text{ISS}} = \bar{S}_L i \not{\partial} S_L - \sigma' \bar{S}_L y_{\sigma'} N_R - \frac{1}{2} \bar{S}_L \mu S_L^c + \text{H.c.}$$

Neutrino masses  $m_\nu \simeq m_D M_D^{-1} \mu (M_D^{-1})^T m_D^T$



# A low-energy variant

Lowering the  $B - L$  scale  $\begin{cases} \rightarrow \text{Resonant leptogenesis} \\ \rightarrow \text{Inverse see-saw} \end{cases}$

Field	Spin	$U(1)_{B-L}$	$U(1)_D$	$U(1)_X$
$S_L$	1/2	0	0	0
$\sigma'$	0	+1	0	0


$$\langle \sigma' \rangle = v_{B-L} \sim \mathcal{O}(\text{TeV})$$

$$\mathcal{L}_{\text{ISS}} = \bar{S}_L i \not{\partial} S_L - \sigma' \bar{S}_L y_{\sigma'} N_R - \frac{1}{2} \bar{S}_L \mu S_L^c + \text{H.c.}$$

Low-scale  $m_{Z_{B-L}}$  allows for annihilations to SM fermions to erase the symmetric component

$$\bar{\chi}\chi \rightarrow Z_{B-L} \rightarrow \bar{q}q(\bar{l}l)$$

(highly suppressed in the high-scale scenario)



# A low-energy variant

Lowering the  $B - L$  scale  $\begin{cases} \rightarrow \text{Resonant leptogenesis} \\ \rightarrow \text{Inverse see-saw} \end{cases}$

Field	Spin	$U(1)_{B-L}$	$U(1)_D$	$U(1)_X$
$S_L$	1/2	0	0	0
$\sigma'$	0	+1	0	0


$$\langle \sigma' \rangle = v_{B-L} \sim \mathcal{O}(\text{TeV})$$

$$\mathcal{L}_{\text{ISS}} = \overline{S}_L i \not{\partial} S_L - \sigma' \overline{S}_L y_{\sigma'} N_R - \frac{1}{2} \overline{S}_L \mu S_L^c + \text{H.c.}$$

The ratio  $m_\nu / M_{N_1}$  is enhanced

Scenarios with  $R \gg 1$  are preferred

$S$  is the dominant DM candidate



# A low-energy variant

Lowering the  $B - L$  scale  $\begin{cases} \rightarrow \text{Resonant leptogenesis} \\ \rightarrow \text{Inverse see-saw} \end{cases}$

Field	Spin	$U(1)_{B-L}$	$U(1)_D$	$U(1)_X$
$S_L$	1/2	0	0	0
$\sigma'$	0	+1	0	0


$$\langle \sigma' \rangle = v_{B-L} \sim \mathcal{O}(\text{TeV})$$

$$\mathcal{L}_{\text{ISS}} = \overline{S}_L i \not{\partial} S_L - \sigma' \overline{S}_L y_{\sigma'} N_R - \frac{1}{2} \overline{S}_L \mu S_L^c + \text{H.c.}$$

Collider searches

$$\bar{q}q \rightarrow Z_{B-L} \rightarrow \bar{\chi}\chi$$

Looking for missing energy from  $\chi$  decays






# The model

Field	Spin	$U(1)_{B-L}$	$U(1)_D$	$U(1)_X$
$N_R^i$	1/2	-1	0	0
$\sigma$	0	+2	0	0
$\chi_0$	1/2	-1	1	0
$\psi_0$	1/2	0	0	+1
$S$	0	0	-1	0
$\phi$	0	+1	-1	+1

$$M_{N_3}, M_{N_2} \gg M_{N_1} \gg m_\chi^0 \gg m_\psi^0, m_S > m_\phi$$



$$\mathcal{L}_{\text{int}} = -y_\nu^{\alpha i} \bar{L}^\alpha \tilde{H} N_R^i - \underline{y_\sigma^{ij} \sigma \overline{N_R^i} N_R^j} - y_S^i S \bar{N}_R^i \chi_0 - y_\phi \phi \bar{\psi}_0 \chi_0 + \text{H.c.}$$


Majorana masses for RHNs from  $U(1)_{B-L}$  breaking  $v_{B-L} \gtrsim 10^{11}$  GeV

➡ neutrino masses (Type- I see-saw)  $m_\nu = -m_D M_N^{-1} m_D^T$

# The model

Field	Spin	$U(1)_{B-L}$	$U(1)_D$	$U(1)_X$
$N_R^i$	1/2	-1	0	0
$\sigma$	0	+2	0	0
$\chi_0$	1/2	-1	1	0
$\psi_0$	1/2	0	0	+1
$S$	0	0	-1	0
$\phi$	0	+1	-1	+1

$$M_{N_3}, M_{N_2} \gg M_{N_1} \gg m_\chi^0 \gg m_\psi^0, m_S > m_\phi$$



$$\mathcal{L}_{\text{int}} = -y_\nu^{\alpha i} \bar{L}^\alpha \tilde{H} N_R^i - y_\sigma^{ij} \sigma \overline{N_R^i} N_R^j - y_S^i S \bar{N}_R^i \chi_0 - y_\phi \phi \bar{\psi}_0 \chi_0 + \text{H.c.}$$


Dark gauge group  $U(1)_D \otimes U(1)_X$

➡ Assure DM stability and Dirac nature of dark fermions (necessary to have an asymmetry)

# The model

Field	Spin	$U(1)_{B-L}$	$U(1)_D$	$U(1)_X$
$N_R^i$	1/2	-1	0	0
$\sigma$	0	+2	0	0
$\chi_0$	1/2	-1	1	0
$\psi_0$	1/2	0	0	+1
$S$	0	0	-1	0
$\phi$	0	+1	-1	+1

$$M_{N_3}, M_{N_2} \gg M_{N_1} \gg m_\chi^0 \gg m_\psi^0, m_S > m_\phi$$



$$\mathcal{L}_{\text{int}} = \underline{-y_\nu^{\alpha i} \bar{L}^\alpha \tilde{H} N_R^i} - \underline{y_\sigma^{ij} \sigma \overline{N_R^i} N_R^j} - \underline{y_S^i S \bar{N}_R^i \chi_0} - y_\phi \phi \bar{\psi}_0 \chi_0 + \text{H.c.}$$


Gauge invariance allows for Yukawa operators

➡ Generation baryon and dark asymmetries  $\eta_\chi = \eta_S \sim \eta_B$

# The model

Field	Spin	$U(1)_{B-L}$	$U(1)_D$	$U(1)_X$
$N_R^i$	1/2	-1	0	0
$\sigma$	0	+2	0	0
$\chi_0$	1/2	-1	1	0
$\psi_0$	1/2	0	0	+1
$S$	0	0	-1	0
$\phi$	0	+1	-1	+1

$$M_{N_3}, M_{N_2} \gg M_{N_1} \gg m_\chi^0 \gg m_\psi^0, m_S > m_\phi$$



$$\mathcal{L}_{\text{int}} = \underline{-y_\nu^{\alpha i} \bar{L}^\alpha \tilde{H} N_R^i} - \underline{y_\sigma^{ij} \sigma \overline{N_R^i} N_R^j} - \underline{y_S^i S \bar{N}_R^i \chi_0} - y_\phi \phi \bar{\psi}_0 \chi_0 + \text{H.c.}$$


$\chi$  and  $S$  get in thermal equilibrium with the SM through gauge and scalar interactions

$$Y_\chi = Y_\chi^{\text{eq}} + \eta_\chi \quad Y_{\bar{\chi}} = Y_\chi^{\text{eq}} \quad \eta_\chi = \eta_S \sim \eta_B$$

# The model

Field	Spin	$U(1)_{B-L}$	$U(1)_D$	$U(1)_X$
$N_R^i$	1/2	-1	0	0
$\sigma$	0	+2	0	0
$\chi_0$	1/2	-1	1	0
$\psi_0$	1/2	0	0	+1
$S$	0	0	-1	0
$\phi$	0	+1	-1	+1

$$M_{N_3}, M_{N_2} \gg M_{N_1} \gg m_\chi^0 \gg m_\psi^0, m_S > m_\phi$$




$$\mathcal{L}_{\text{int}} = -y_\nu^{\alpha i} \bar{L}^\alpha \tilde{H} N_R^i - y_\sigma^{ij} \sigma \overline{N_R^i} N_R^j - y_S^i S \bar{N}_R^i \chi_0 - \underline{y_\phi \phi \bar{\psi}_0 \chi_0} + \text{H.c.}$$

We assume  $\begin{cases} y_\phi \ll 1 \\ g_X \ll 1 \end{cases}$  so that  $\psi$  is never in thermal equilibrium

# The model

Field	Spin	$U(1)_{B-L}$	$U(1)_D$	$U(1)_X$
$N_R^i$	1/2	-1	0	0
$\sigma$	0	+2	0	0
$\chi_0$	1/2	-1	1	0
$\psi_0$	1/2	0	0	+1
$S$	0	0	-1	0
$\phi$	0	+1	-1	+1

$$M_{N_3}, M_{N_2} \gg M_{N_1} \gg m_\chi^0 \gg m_\psi^0, m_S > m_\phi$$



$$\mathcal{L}_{\text{int}} = -y_\nu^{\alpha i} \bar{L}^\alpha \tilde{H} N_R^i - y_\sigma^{ij} \sigma \overline{N_R^i} N_R^j - y_S^i S \bar{N}_R^i \chi_0 - \underline{y_\phi \phi \bar{\psi}_0 \chi_0} + \text{H.c.}$$


The  $\chi$  asymmetry is transferred to  $\psi$  through late decays  $\chi \rightarrow \psi + \phi$

# Asymmetric freeze-out


## Asymmetry and Fractional asymmetry

$$\eta_D = Y_\chi - Y_{\bar{\chi}} \quad r = Y_{\bar{\chi}}/Y_\chi$$


Asymmetry generation



Annihilation of the symmetric component



Only asymmetric component survives



$$r_\infty \propto \exp \left[ -\frac{\pi g_*}{45 x_f} M_{\text{Pl}} \sigma v \eta_D m \right]$$


Graesser, Shoemaker, Vecchi  
[1103.2771]

# Asymmetric freeze-out


Asymmetry and Fractional asymmetry

$$\eta_D = Y_\chi - Y_{\bar{\chi}} \quad r = Y_{\bar{\chi}}/Y_\chi$$


Asymmetry generation



Annihilation of the symmetric component



Only asymmetric component survives



$$r_\infty \propto \exp \left[ -\frac{\pi g_*}{45 x_f} M_{\text{Pl}} (\sigma v) \eta_D m \right]$$

Graesser, Shoemaker, Vecchi  
[1103.2771]




# Asymmetric freeze-out

Asymmetry and Fractional asymmetry

$$\eta_D = Y_\chi - Y_{\bar{\chi}} \quad r = Y_{\bar{\chi}}/Y_\chi$$

Annihilation of the symmetric component



**Large cross sections  
are needed!**

$$\sigma v > \sigma v_{\text{WIMP}}$$


**Asymmetric DM out of equilibrium (tiny couplings, freeze-in) ?**

(How to erase the symmetric component?)


# Early vs Late decays

$$T_D \simeq 10 \text{ MeV} \left( \frac{y_\phi}{10^{-12}} \right)$$

$$2 \times 10^{-10} < y_\phi < 5 \times 10^{-7}$$




Late decays take place at  $x \gg 1$



Dominant production from early decays takes place around  $x \sim 1$

Asymmetric freeze-out takes place around  $x \sim 20$



No annihilations

# Early vs Late decays

$$T_D \simeq 10 \text{ MeV} \left( \frac{y_\phi}{10^{-12}} \right)$$

$$y_\phi < 6 \times 10^{-12}$$

